


Pneumococcal Vaccine Timing—For Adults

DO NOT administer PCV13 and PPSV23 at the same visit.

Age 65 Years or Older

- If PCV13 was given before age 65 years, no additional PCV13 is needed.



Age 19-64 Years With Underlying Condition(s)

- Prior doses count towards doses recommended below and do not need to be repeated.
- If PPSV23 given previously – wait one year before giving PCV13
 - for group B, wait at least five years before giving a second dose of PPSV23.
- No more than two doses of PPSV23 recommended before 65th birthday and one dose thereafter.

A. Smoker, Long-term facility resident, or Chronic conditions:

- heart disease (excluding hypertension)
- lung disease (including asthma)
- liver disease (including cirrhosis)
- diabetes
- alcoholism

PPSV 23

B. Immunocompromised (including HIV infection), Chronic renal failure, Nephrotic syndrome, or Asplenia

PCV 13

8 weeks

PPSV 23

5 years

PPSV 23

C. CSF leaks or Cochlear implants

PCV 13

8 weeks

PPSV 23



For further details, see: www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/pneumo.html

California Department of Public Health, Immunization Branch www.EZIZ.org

This publication was supported by Grant Number H23/CCH922507 from the Centers for Disease Control and Prevention (CDC)

IMM-1152 (9/15)