# Original Article Characterization of Susceptibility of Inbred Mouse Strains to Diabetic Nephropathy

Zhonghua Qi,<sup>1</sup> Hiroki Fujita,<sup>1</sup> Jianping Jin,<sup>1</sup> Linda S. Davis,<sup>1</sup> Yihan Wang,<sup>2</sup> Agnes B. Fogo,<sup>2</sup> and Matthew D. Breyer<sup>1,3,4</sup>

Differential susceptibility to diabetic nephropathy has been observed in humans, but it has not been well defined in inbred strains of mice. The present studies characterized the severity of diabetic nephropathy in six inbred mouse strains including C57BL/6J, DBA/2J, FVB/NJ, MRL/MpJ, A/J, and KK/HIJ mice. Diabetes mellitus was induced using low-dose streptozotocin injection. Progression of renal injury was evaluated by serial measurements of urinary albumin excretion, glomerular filtration rate (GFR), and terminal assessment of renal morphology over 25 weeks. Despite comparable levels of hyperglycemia, urinary albumin excretion and renal histopathological changes were dramatically different among strains. DBA/2J and KK/HIJ mice developed significantly more albuminuria than C57BL/6J, MRL/MpJ, and A/J mice. Severe glomerular mesangial expansion, nodular glomerulosclerosis, and arteriolar hyalinosis were observed in diabetic DBA/2J and KK/HIJ mice. Glomerular hyperfiltration was observed in all diabetic strains studied except A/J. The significant decline in GFR was not evident over the 25-week period of study, but diabetic DBA/2J mice exhibited a tendency for GFR to decline. Taken together, these results indicate that differential susceptibility to diabetic nephropathy exists in inbred mice. DBA/2J and KK/HIJ mice are more prone to diabetic nephropathy, whereas the most widely used C57BL/6J mice are relatively resistant to development of diabetic nephropathy. Diabetes 54:2628-2637, 2005

© 2005 by the American Diabetes Association.

iabetic nephropathy is an insidious and lethal complication of diabetes mellitus leading to renal failure in a substantial fraction of patients with diabetes mellitus. Epidemiological studies in humans suggest that the lifetime risk for development of diabetic nephropathy is  $\sim 35\%$  (1,2). Significant racial differences have been reported including increased incidence in native Americans and Asian populations, indicating differential susceptibility to diabetic nephropathy exists in humans (1.3-6). In cohorts susceptible to diabetic nephropathy, patients progressively increase urinary albumin excretion (UAE) followed by a decline of glomerular filtration rate (GFR). These functional changes are accompanied by characteristic renal pathological features including mesangial expansion, glomerulosclerosis, and tubulointerstitial fibrosis (5). In contrast, cohorts resistant to diabetic nephropathy do not develop severe renal histopathological lesions or albuminuria despite comparable levels of hyperglycemia (2,7). Genetic factors have been suggested to contribute to this differential susceptibility to diabetic nephropathy in humans (2-4,8). Despite substantial effort, mapping of susceptibility genes in people has been hindered because of the genetic heterogeneity in human populations as well as the diversity of environmental factors including treatment.

Inbred mice have been wildly used to model human diseases not only because they facilitate gene manipulation, but also because they share high genetic homology with humans (9-11). Many mouse models of diabetes have been established including type 1 diabetes due to streptozotocin (STZ) treatment and genetic models (e.g., C57BL/  $6-Ins2^{Aklta}/J$ ) (12–14). Models of type 2 diabetes include diet-induced and inherited deficiency in the leptin receptor (i.e., db/db) mice (15–17). However, nephropathy has been studied in only a limited number of strains, and none of these models develops renal lesions that fully resemble advanced human diabetic nephropathy (18). The majority of studies have been performed using C57BL/6J mice; however, this strain does not typically develop robust renal histopathological changes or marked albuminuria (18–21). These deficiencies have hindered the use of the unique genetic murine reagents to dissect the mechanistic underpinnings of diabetic nephropathy.

An inbred mouse strain is established by sibling mating for 20 or more consecutive generations (available at

From the <sup>1</sup>Division of Nephrology, Department of Medicine, Vanderbilt University, Nashville, Tennessee; the <sup>2</sup>Department of Pathology, Vanderbilt University, Nashville, Tennessee; the <sup>3</sup>Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, Tennessee; and the <sup>4</sup>Veterans Administration Medical Center, Nashville, Tennessee.

Address correspondence and reprint requests to Zhonghua Qi, MD, F-424 ACRE Building, Veterans Administration Medical Center, Nashville, TN 37212. E-mail: zhonghua.qi@vanderbilt.edu.

Received for publication 24 January 2005 and accepted in revised form 6 June 2005.

ACR, albumin-to-creatinine ratio; FITC, fluorescein isothiocyanate; GBM, glomerular basement membrane; GFR, glomerular filtration rate; STZ, strep-tozotocin; UAE, urinary albumin excretion rate.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

http://www.jax.org). Thus mice within an inbred strain are genetically identical, whereas mice from other inbred strains are genetically distinct from the index strain (22,23). As in human populations, differential susceptibility to some renal diseases has been demonstrated among inbred mice (21,24,25); however, the susceptibility of inbred mice to diabetic nephropathy has not been defined. Characterization of this susceptibility may not only help establish more robust mouse models of diabetic nephropathy, but also provide a basis for mapping of the underlying genes predisposing to the development of diabetic nephropathy. The present studies compared the development and severity of three major phenotypic features of diabetic nephropathy—albuminuria, histopathology, and GFR—in six commonly used inbred strains of mice.

#### **RESEARCH DESIGN AND METHODS**

The inbred male mice used in the present studies were purchased from The Jackson Laboratory (Bar Harbor, ME) at 6–8 weeks of age. STZ and fluorescein isothiocyanate (FTTC)-inulin were obtained from Sigma-Aldrich (St. Louis, MO). The enzymatic immunoassay kits for determining urinary albumin and creatinine were purchased from Exocell (Philadelphia, PA). All protocols were approved by the institutional animal care and use committee of Vanderbilt University and recommended by the Animal Models of Diabetic Complications Consortium (available at http://www.amdcc.org).

**Induction of diabetes in inbred mice.** At 10 weeks of age, inbred mice received daily STZ injections intraperitoneally (40 mg/kg for DBA/2J and 50 mg/kg for all other strains, made freshly in 0.1 mol/l citrate buffer, pH 4.5) for 5 consecutive days. The onset of diabetes was evaluated by measuring fasting blood glucose and HbA<sub>1c</sub> (A1C). Blood glucose was measured biweekly using a B-Glucose Analyzer (HemoCue, Lake Forest, CA) on samples obtained after a 6-h fast starting at 6:00 A.M. A1C levels were determined monthly using a commercially available analysis kit (DCA 2000; Bayer, Elkhart, IN). Blood was collected in conscious mice via the saphenous vein as described previously (26).

**Measurement of UAE.** Excretion of urinary albumin was determined using albumin-to-creatinine ratio (ACR) on morning spot urine and UAE in 24-h urine collections. Spot urine collection was conducted monthly using a custom-made mouse urine collection station that used a 96-well enzyme-linked immunosorbent assay plate as the floor. Mice were allowed to roam freely on this 96-well plate until they spontaneously urinated. Urine was obtained in the wells without contamination by feces, using a pipette. Twenty-four-hour urine was collected using metabolic cages (Braintree Scientific, Braintree, MA). The concentration of albumin and creatinine in the urine were determined using a commercially available kit (Exocell).

**Measurement of GFR in conscious mice.** Renal function in diabetic mice was evaluated by serial determination of GFR before and 5, 15, and 25 weeks after the onset of hyperglycemia. FTTC-inulin clearance was determined as described previously (26). Briefly, 3% FTTC-inulin was injected retro-orbitally, followed by collection of ~20 µl of saphenous vein blood at 3, 7, 10, 15, 35, 55, and 75 min after the FTTC-inulin bolus injection in conscious mice. Plasma fluorescence concentration at each time point was determined using a Fluoroscan Ascent FL (Labsystems, Helsinki, Finland) with 485-nm excitation and read at 538-nm emission. The decay in plasma fluorescence levels was fit to a two-phase exponential decay curve using nonlinear regression (GraphPad Prism; GraphPad Software, San Diego, CA.). GFR is calculated using the equation: GFR =  $I/(A/\alpha + B/\beta)$ , where *I* is the amount of FITC-inulin delivered in bolus injection; *A* and  $\alpha$  are the *y*-intercept and the decay constant of the rapid (initial) decay phase, respectively; and *B* are the *y*-intercept and the decay constant of the slow decay phase, respectively (26).

**Renal histopathology.** Mice were killed after hyperglycemia was established for at least 25 weeks. Under anesthesia induced with phentobarbital sodium injection (50 mg/kg body wt i.p.), the left renal artery and vein were clipped with a hemostatic forceps. The left kidney was removed, and the weight was measured. The right kidney was perfused with PBS (pH 7.0) through a butterfly 23-gauge needle inserted into left ventricle at 160–170 mmHg for ~5 min. This is followed by 4% paraformaldehyde for another 5 min. The perfused right kidney was removed and routinely processed. Four-micrometer sections were stained with periodic acid Schiff. A semiquantitative score was used to evaluate the degree and extent of glomerulosclerosis as described previously (27). Mesangial matrix expansion occupying <25, 25–50, 50–75, or >75% of tuft was scored 1, 2, 3, and 4+, respectively, and no mesangial expansion was

scored as 0. A whole kidney average sclerosis index was obtained by averaging scores from all glomeruli on one section. On average, more than 100 glomeruli were assessed per mouse. Tubules and interstitium were also evaluated by light microscopy. Electron microscopic examination was conducted in the Research Electron Microscopy Core of Vanderbilt University. Samples were fixed in 2.5% glutaraldehyde in 0.1 mol/l cacodylate buffer (pH 7.4). After fixation, samples were dehydrated through a graded series of ethanol and embedded in Spurr resin. The sections (80–100 nm) were viewed using a FEI/Philips CM12 transmission electron microscope operated at 80 KeV.

**Statistics.** All data are expressed as means  $\pm$  SE. ANOVA and t test were used for data analysis. P < 0.05 was considered significant.

## RESULTS

STZ-induced diabetes in inbred mouse strains. All strains of mice tested developed stable hyperglycemia 1 week after low-dose STZ injection (Table 1). A1C was elevated in all studied mice compared with either each baseline or age-matched controls (Table 1). Strain-dependent susceptibility to STZ was observed. KK/HlJ mice appeared relatively resistant to STZ-induced hyperglycemia with lower levels of blood glucose. Without the insulin supplementation, most diabetic mice survived more than 25 weeks. After that, the mortality increased in most strains studied, especially in DBA/2J, KK/HIJ, and A/J strains (data not shown). In contrast, C57BL/6J mice appear to tolerate persistent hyperglycemia well with a group of diabetic C57BL/6J mice surviving longer than 45 weeks despite fasting glucose levels of 300-600 mg/dl. Compared with age-matched controls, body weight was lower in diabetic mice except in FVB/NJ and MRL/MpJ strains (Table 1). Substantial loss of body weight was especially evident in diabetic DBA/2J mice.

Albuminuria in diabetic inbred mice. Albuminuria is a central manifestation of diabetic nephropathy (2,5,16,28). Spot urine ACR and 24-h UAE were examined in the diabetic and control mice in the present studies. Spot urine ACR was serially measured and 24-h UAE was determined in mice after 25 weeks of hyperglycemia. A significant correlation between spot ACR and 24-h UAE was observed in C57BL/6J, DBA/2J, and KK/HIJ mice (Figs. 1 and 2), but not in FVB/NJ mice (Fig. 2).

Despite persistent hyperglycemia, no significant increase in ACR was observed in C57BL/6J and FVB/NJ mice (Table 1 and Fig. 2A). A separate group of diabetic C57BL/6J mice were followed for 45 weeks, and these mice also failed to develop albuminuria (fasting blood glucose, 446.0  $\pm$  103.1 mg/dl; ACR, 19.3  $\pm$  5.8 µl/mg; n = 6). In contrast, diabetic DBA/2J and KK/HIJ mice exhibited a significant increase in ACR compared with their agematched controls or other strains of diabetic mice (Table 1 and Fig. 2). Hyperglycemic A/J and MRL/MpJ mice intermittently increased ACR, but the levels were significantly less than DBA/2J and KK/HIJ mice.

Consistent with the spot ACR results, 24-h UAE was significantly greater in diabetic DBA/2J and KK/HIJ mice than diabetic C57BL/6J, A/J, and MRL/MpJ mice after ~25 weeks of hyperglycemia (Fig. 2B). In contrast, diabetic FVB/NJ mice exhibited a marked increase in 24-h UAE despite an unchanged spot urine ACR (Fig. 2B). Further analysis of these mice showed that diabetic FVB/NJ mice significantly increased 24-h urinary creatinine excretion after ~25 weeks of hyperglycemia (1.52 vs. baseline at 0.37 mg/24 h, P < 0.001). In contrast, other diabetic mouse

#### TABLE 1

Fasting blood glucose and UAE in diabetic and control mice

Strain	Glycemic		Weeks after STZ or vehicle		
	status	Before STZ	5	15	25
C57BL/6J					
FBS (mg/dl)	DM	$170.0 \pm 11.2 (10)$	$424.9 \pm 33.9 \ (9)^{*\dagger}$	$502.5 \pm 31.2 (21)^{*\dagger}$	$506.6 \pm 46.0 (18)^{*\dagger}$
	CN	$170.0 \pm 3.7 (10)$	$170.4 \pm 7.1$ (5)	$164.8 \pm 4.7$ (5)	$157.4 \pm 8.1$ (5)
A1C (%)	DM	$3.8 \pm 0.2$ (3)	NA	$8.2 \pm 0.4 (21)^{*\dagger}$	$9.4 \pm 0.6 (11)^{*\dagger}$
	CN	$3.6 \pm 0.1$ (5)	NA	$4.1 \pm 0.1 (5)^*$	$3.8 \pm 0.1$ (5)
BW (g)	DM	$21.3 \pm 0.4$ (10)	$25.6 \pm 0.6(9)^*$	$25.6 \pm 0.7 (21)^{*+}$	$26.9 \pm 0.8$ (18)*
	CN	$20.3 \pm 0.2$ (5)	$26.7 \pm 1.2 (5)^*$	$28.9 \pm 0.9(5)^*$	$29.2 \pm 1.4(5)^{*}$
ACR (µg/mg)	DM	36.9 + 7.6(9)	$35.4 \pm 5.7$ (9)	$29.5 \pm 12.4 (16)^*$	$72.1 \pm 20.4$ (16)
	CN	$45.6 \pm 12.4$ (5)	$201 \pm 16(5)$	48.11 + 16.5(5)	532 + 243(5)
DBA/2.I	en	10.0 = 12.1 (0)	10.1 = 1.0 (0)	10.11 = 10.0 (0)	
FBS (mg/dl)	DM	$160.5 \pm 8.6(15)$	$485.8 \pm 31.3(19)*\pm$	485.3 + 27.4 (21)*†	$511.3 \pm 75.6(6)*$
	CN	$165.2 \pm 16.2(5)$	$1274 \pm 155(5)^{*}$	$148.6 \pm 8.5(5)$	$152.6 \pm 4.5(5)$
A1C (%)	DM	$29 \pm 01(3)$	NA	$67 \pm 0.1(21)^{*+}$	$134 \pm 0.8(6)^{*+\pm}$
	CN	$2.8 \pm 0.1$ (5)	NA	$28 \pm 01(5)$	$27 \pm 02(5)$
BW (g)	DM	$234 \pm 0.6(15)$	$22.3 \pm 0.7(19)^{\ddagger}$	$20.8 \pm 0.7 (21)*$	$18.7 \pm 0.9(10)*$
DW (g)	CN	$22.2 \pm 0.9(6)$	$282 \pm 0.8(4)^{*}$	$29.5 \pm 1.2 (4)^*$	$30.1 \pm 0.9 (5)^{*}$
ACR (µg/mg)	DM	$26.6 \pm 6.6$ (29)	4244 + 894(24)*	$608.0 \pm 220.8(15)***8$	421.4 + 167.3(11)***8
	CN	$19.7 \pm 5.1(5)$	487 + 70(9)*	71.1 + 15.2(7)*	$65.8 \pm 13.2(7)^*$
A/J	011	10.1 = 0.1(0)	10.1 = 1.0 (0)	11.1 = 10.2 (1)	00.0 = 10.2 (1)
FBS (mg/dl)	DM	$1544 \pm 44(15)$	$360.8 \pm 28.0(10)*$	505 6 + 31 4 (22)**	391.6 + 50.6 (16)**
	CN	$159.8 \pm 7.8(5)$	$146.0 \pm 13.4(4)$	$157.0 \pm 4.0(4)$	$1710 \pm 21(4)$
A1C (%)	DM	$26 \pm 0.03(9)$	$\frac{110.0}{NA}$	$51 \pm 01(22)*\pm \P$	$41 \pm 02(16)^{*}$
1110 (70)	CN	$2.0 \pm 0.03(5)$ $2.6 \pm 0.03(5)$	NA	$2.6 \pm 0.04(5)$	$26 \pm 0.04(5)$
BW (g)	DM	$23.0 \pm 0.03(0)$	$18.3 \pm 0.6(10)*$	$205 \pm 0.8(27)$ *	$232 \pm 0.01(8)$
Dw (g)	CN	$29.4 \pm 0.6(10)$	$268 \pm 0.8(4)*$	$20.9 \pm 0.0 (21)$ + $28.2 \pm 0.9 (4)*$	$20.2 \pm 0.1 (10)^{+}$ $30.6 \pm 1.3 (5)^{*}$
$ACR (\mu \sigma/m\sigma)$	DM	$26.6 \pm 12.2$ (23)	$20.0 \pm 0.0 (4)$ 1195 + 211 (27)*	174.0 + 48.2(17)*	$64.0 \pm 12.6(15)*$
non (µg/mg)	CN	$18.7 \pm 22.(5)$	$545 \pm 142(5)*$	884 + 258(4)*	81.2 + 8.8(4)*
FVB/NI	OIN	$10.7 \pm 2.2 (0)$	$04.0 \pm 14.2(0)$	$00.4 \pm 20.0$ (4)	$01.2 \pm 0.0$ (4)
FBS (mg/dl)	DM	$141.8 \pm 12.4(12)$	$576.0 \pm 43.0(8)$ **	$549.4 \pm 52.1(10)*$	$610.0 \pm 23.3(5)*$
i bo (ing/ui)	CN	$134.0 \pm 10.4(12)$	$1695 \pm 25(4)$	$1483 \pm 108(4)$	$151.0 \pm 12.1(4)$
A1C (%)	DM	$30 \pm 01(7)$	$\frac{100.0}{NA}$	$58 \pm 02(10)*$	$50 \pm 0.3(5)^{*}$
AIC (%)	CN	$0.0 \pm 0.1$ (1)	NA	$3.0 \pm 0.2$ (10) +	$0.0 \pm 0.5(0)$
$\mathbf{BW}(\mathbf{a})$	DM	$270 \pm 0.8(12)$	$268 \pm 0.2(8)$	$97.4 \pm 0.6(10)$	$33.0 \pm 0.5$ (5)*
Dw (g)	CN	$27.5 \pm 0.0 (12)$ $26.7 \pm 2.0 (4)$	$20.0 \pm 0.3 (8)$ $20.0 \pm 2.2 (4)$	$27.4 \pm 0.0$ (10) $30.0 \pm 2.1$ (4)	$33.0 \pm 0.5(5)^{\circ}$ $33.1 \pm 1.8(4)$
ACP(ug/mg)	DM	$20.7 \pm 2.0 (4)$ 61.6 + 13.2 (7)	$61 \pm 14(9)$ **	$30.0 \pm 2.1 (4)$ 88.0 + 21.3 (12)	$80.3 \pm 27.7(7)$
AON (µg/mg)	CN	$45.4 \pm 18.5(4)$	$74.1 \pm 21.7(4)$	$\frac{11.5}{12}$	$63.0 \pm 21.1(1)$ $63.0 \pm 31.0(4)$
MRI/MnI	ON	$40.4 \pm 10.0(4)$	$74.1 \pm 21.7$ (4)		$05.0 \pm 51.0$ (4)
FBS (mg/dl)	DM	$120.8 \pm 6.6(5)$	$434.8 \pm 22.7(10)*$	$403.5 \pm 21.0(17)*\pm$	428.0 + 45.1(5)*+
rbs (ing/ui)	CN	$120.0 \pm 0.0(3)$ $140.5 \pm 4.8(4)$	$165.0 \pm 3.7(4)$	$455.0 \pm 21.5(11)$	$1535 \pm 202(4)$
A1C (%)	DM	$30 \pm 01(5)$	$53 \pm 01(19)$	$60 \pm 02(8)*$	$61 \pm 0.2(4)$
AIC $(70)$		$3.0 \pm 0.1 (3)$ $3.0 \pm 0.1 (4)$	$5.5 \pm 0.1$ (13)	$0.5 \pm 0.2 (6)^{-1}$	$0.1 \pm 0.2 (0)^{+1}$ $3.3 \pm 0.4 (4)$
$\mathbf{DW}(\mathbf{a})$	DM	$3.0 \pm 0.1 (4)$ $27.1 \pm 0.4 (5)$	1NA $24.0 \pm 0.8(10)*$	$3.0 \pm 0.2$ (4) $26.7 \pm 0.6$ (17)	$3.3 \pm 0.4 (4)$ $27.0 \pm 1.4 (5)$
BW (g)		$37.1 \pm 0.4 (3)$ $35.7 \pm 1.4 (4)$	$34.0 \pm 0.8 (13)^{\circ}$ $36.5 \pm 2.6 (4)$	$30.7 \pm 0.0 (17)$ $38.6 \pm 3.7 (4)$	$31.0 \pm 1.4(0)$ $30.2 \pm 4.3(4)$
ACR (µg/mg)	DM	$55.7 \pm 1.4 (4)$ $74.4 \pm 17.0 (14)$	$05.9 \pm 42.0$ (4)	$100 \pm 0.7 (4)$ $100 5 \pm 0.7 (10) \pm$	$97.2 \pm 4.3 (4)$
		$74.4 \pm 17.0 (14)$ $20.5 \pm 14.9 (4)$	$95.2 \pm 45.4 (20)$ $72.2 \pm 14.5 (4)$	$122.0 \pm 52.7$ (10)	$67.7 \pm 22.3 (0)$
KK/HII	UN	$39.3 \pm 14.2$ (4)	$75.2 \pm 14.5 (4)$	$16.0 \pm 7.5$ (4)	$45.2 \pm 14.0(4)$
FPS (mg/dl)	DM	$160.6 \pm 7.4$ (15)	$207.5 \pm 20.6 (15) * \pm 1$	$221.2 \pm 22.0 (10)*+\parallel$	$220.0 + 22.8 (12)*+\parallel$
i Do (mg/m)		$100.0 \pm 7.4 (10)$ $142.0 \pm 5.2 (5)$	$297.5 \pm 50.0 (15)^{\circ}$	$521.5 \pm 25.0 (19)^{+}$	$550.0 \pm 52.8 (13)^{\circ}$
A1C (%)		$142.0 \pm 0.2 (0)$ $15 \pm 0.1 (15) \pm$	$107.4 \pm 9.3(10)$	$6.4 \pm 0.4 (10) **$	$6.4 \pm 0.5 (10) + 6.4 \pm 0.5 (10) + 6.4 \pm 0.5 (10) + 100 + 1$
		$4.0 \pm 0.1 (10)$ $4.7 \pm 0.2 (5)$		$0.4 \pm 0.4 (19)^{-1}$	$0.4 \pm 0.0 (10)^{++}$ $4.0 \pm 0.1 (2)*$
BW (g)		$4.7 \pm 0.2 (0)$ $21.4 \pm 1.0 (05)$	1NA 20.8 + 0.7 (15)	$4.0 \pm 0.2 (10)$ $30.0 \pm 0.6 (10)$	$4.0 \pm 0.1 (0)^{+}$ $32.2 \pm 0.2 (12)^{+}$
		$31.4 \pm 1.0 (20)$ $31.4 \pm 1.5 (7)$	$23.0 \pm 0.7 (10)$ $33.8 \pm 1.5 (5)$	$30.9 \pm 0.0 (19)^{\dagger}$ 36.6 + 1.4 (10)*	$32.3 \pm 0.0 (13)$ $35.0 \pm 1.1 (12)$ *
ACD (		$01.4 \pm 1.0(1)$ 72.0 + 12.1(12)	$33.0 \pm 1.3 (3)$ $500.0 \pm 97.6 (10) *^{\pm}$	$50.0 \pm 1.4 (10)^{-1}$	$00.9 \pm 1.1 (10)^{\circ}$ 695.9 $\pm 100.0 (10)***$
AUR (µg/mg)	CN	$72.8 \pm 12.1 (13)$ $72.8 \pm 11.4 (14)$	$177.0 \pm 50.7 (10)$	$224.4 \pm 80.8 (10)^*$	$349.1 \pm 39.7 (10)^*$

Data are means  $\pm$  SE (no. of animals). \*P < 0.05 vs. baseline at same group; †P < 0.05 vs. age-matched controls at same time point; ‡P < 0.05 vs. all other diabetic strains at same time point; \$P < 0.05 vs. diabetic C57BL/6J, A/J, FVB/NJ, and MRL/MpJ mice; ¶P < 0.001 vs. A1C in all other diabetic strains except FVB/NJ strain at same time point; ||P < 0.05 vs. FBS in diabetic C57BL/6J, DBA/2J, and FVB/NJ mice at same time point. BW, body weight; CN, control mice; DM, diabetic mice; FBS, fasting blood glucose; NA, not available.

strains exhibited no change (e.g., A/J and KK/HIJ mice) or a decrease (e.g., DBA/2J mice) in 24-h urinary creatinine excretion. Urine volume in hyperglycemic FVB/NJ mice  $(12.96 \pm 3.06 \text{ ml}/24 \text{ h}, n = 7)$  was also significantly greater than in other diabetic strains including C57BL/6J (2.74  $\pm$  0.74 ml/24 h, n = 13), DBA/2J (2.56  $\pm$  0.38 ml/24 h, n = 14), A/J (0.45  $\pm$  0.10 ml/24 h, n = 6), or KK/HIJ mice (1.95  $\pm$  0.91 ml/24 h; n = 11). In contrast to previous studies



FIG. 1. Relationship between morning spot urine ACR and 24-h UAE in diabetic mice. Data were obtained from diabetic and control C57BL/6J (n = 4), DBA/2J (n = 6), and KK/HIJ (n = 9) mice. Spot urine was collected for 3 consecutive days from each mouse and average spot ACR was used for the correlation analysis.  $R^2 = 0.69$ . P < 0.0001.

examining a transgenic diabetic model with FVB/N background (OVE26) (29), we did not observe hydronephrosis in STZ-induced hyperglycemic FVB/NJ mice, despite the presence of polyuria.

Moderate albuminuria was also observed in non-STZinjected control KK/HIJ mice (Table 1 and Fig. 2), despite lack of increased fasting blood glucose or A1C (Table 1). However, albuminuria in control KK/HIJ mice was significantly less than in STZ-injected hyperglycemic KK/HIJ mice.

**Development of albuminuria in C57BL/6-Ins2**<sup>Akita</sup>/J mice. To confirm the resistance of diabetic C57BL/6J mice, we examined the UAE in a second model of type 1 diabetes, the C57BL/6-*Ins2*<sup>Akita</sup>/J mouse. These mice express a mutation in the insulin 2 gene and develop hyper-glycemia because of misfolding of insulin and proteotoxicity of  $\beta$ -cells (14,30,31). As in STZ-injected diabetic model, male C57BL/6-*Ins2*<sup>Akita</sup>/J mice did not develop significant albuminuria despite 20 weeks of hyper-glycemia (at 30 weeks of age: fasting blood glucose, 631.6 ± 55.2 mg/dl, and ACR, 29.2 ± 4.2 µg/mg; n = 8).

**Renal morphology in diabetic inbred mice.** To examine the relationship between albuminuria and renal histopathological lesions, renal morphology was studied by light and electronic microscopy. Consistent with albuminuria, diabetic DBA/2J and KK/HIJ mice exhibited the most dramatic histopathological changes after 25–35 weeks of hyperglycemia (Fig. 3). Significantly greater mesangial expansion was observed in diabetic DBA/2J and KK/HIJ kidneys with some glomeruli developing nodular glomerular sclerosis and arteriolar hyalinosis (Figs. 3 and 4). Diabetic C57BL/6J mice also exhibited increased mesangial expansion scores compared with age-matched controls, but the scores were significantly less than in diabetic DBA/2J and KK/HIJ mice. In contrast, despite developing



FIG. 2. Development of albuminuria in STZ-induced diabetic inbred mice after ~25 weeks of hyperglycemia. A: Spot urine ACR in diabetic mice (**II**) and age-matched controls ( $\Box$ ). \*P < 0.05 (t test) vs. respective age-matched controls; †P < 0.05 (ANOVA) vs. diabetic C57BL/6J, A/J, MRL/MpJ, and FVB/NJ mice; and ‡P < 0.001 (ANOVA) vs. controls in other strains. B: 24-h UAE in diabetic mice (**II**) and age-matched controls ( $\Box$ ). Age-matched controls for MRL/MpJ and FVB/NJ strains were not available (NA) for studying. \*P < 0.05 (t test) vs. age-matched controls; \$P < 0.05 (ANOVA) vs. diabetic C57BL/6J, A/J, and MRL/MpJ strains; and ||P < 0.05 (ANOVA) vs. controls in other studied strains.



FIG. 3. Representative glomerular histopathology of perfuse-fixed diabetic mouse kidneys (periodic acid Schiff, ×400). A: C57BL/6J mice (35 weeks after STZ). B: A/J (32 weeks after STZ). C: MRL/MpJ (35 weeks after STZ). D: FVB/NJ (25 weeks after STZ). E: DBA/2J (25 weeks after STZ). F: Arteriolar hyalinosis (arrow) in diabetic DBA/2J mice. G: KK/HIJ mice (35 weeks after STZ). H: Nodular glomerulosclerosis (arrow) in diabetic KK/HIJ mice. I: Arteriolar hyalinosis (arrow) in diabetic KK/HIJ mice.

albuminuria, diabetic FVB/NJ mice exhibited significantly less mesangial expansion than diabetic DBA/2J, KK/HIJ, or C57BL/6J mice. Glomerular mesangial expansion in nonhyperglycemic KK/HIJ mice was significantly greater than control DBA/2J and C57BL/6J mice but less than STZinduced hyperglycemic KK/HIJ mice. These differential levels of glomerulosclerosis are consistent with their relative levels of albuminuria. Interstitial fibrosis or tubu-

![](_page_4_Figure_4.jpeg)

FIG. 4. Glomerular mesangial expansion scores in diabetic inbred mice. The scores were determined using light microscopy at ×400 in perfuse-fixed diabetic mouse kidneys as described in RESEARCH DESIGN AND METHODS. \*P < 0.05 (t test) vs. age-matched controls; †P < 0.05 (ANOVA) vs. diabetic C57BL/6J, A/J, MRL/MpJ, and FVB/NJ mice; and  $\ddagger P < 0.05$  vs. controls in other strains. Age-matched controls for diabetic FVB/NJ and MRL/MpJ were not available (NA).

lar atrophy was not observed in any strain of diabetic mouse within the 25 weeks of study.

Glomerular ultrastructure was further examined using an electronic microscope. Diabetic DBA/2J and KK/HIJ mice developed the most dramatic thickening of glomerular basement membrane (GBM) among the strains studied. Diabetic C57BL/6J mice also exhibited increased GBM width, but the level was significantly less than diabetic DBA/2J and KK/HIJ mice (Figs. 5 and 6). No electronic dense deposits were observed in the basement membrane. Fragmental podocyte foot process effacement was observed in some glomeruli of diabetic KK/HIJ and DBA/2J strains.

**Kidney weight in diabetic inbred mice.** Renal hypertrophy is observed in type 1 diabetic patients with early-stage diabetic nephropathy (32,33). In the present studies, we examined kidney weight-to-body weight ratio in diabetic and age-matched controls. As shown in Fig. 7, a significant increase in kidney weight was observed in all studied strains of diabetic mice except A/J strains (the age-matched controls for diabetic FVB/NJ mice were not available). Diabetic DBA/2J mice exhibited the highest kidney weight-to-body weight ratio among the studied strains of diabetic mice (Fig. 7).

**GFR in diabetic inbred mice.** GFR was serially examined in diabetic inbred mice and age-matched controls using FITC-inulin clearance (26). GFR increased in all strains of diabetic mice compared with age-matched controls except for A/J (Fig. 8). Among the strains studied, diabetic DBA/2J and FVB/NJ exhibited the highest GFR at

![](_page_5_Picture_1.jpeg)

15 and 25 weeks after hyperglycemia (P < 0.05), respectively. A significant decline in GFR was not observed in these diabetic mice within the 25-week period of study compared with their baseline. Nevertheless, a tendency for GFR to decline was observed in diabetic DBA/2J mice (Fig. 8). The GFR calculated per mouse was listed in Table 2.

## DISCUSSION

The progression of diabetic nephropathy in humans has been suggested to be significantly affected by genetic

![](_page_5_Figure_5.jpeg)

FIG. 6. GBM width in diabetic inbred mice after  $\sim 25$  weeks of hyperglycemia ( $\blacksquare$ ) and age-matched controls ( $\square$ ). \*P < 0.05 (t test) vs. age-matched controls; †P < 0.05 (ANOVA) vs. diabetic C57BL/6J mice. The age-matched controls in FVB/NJ strain were not available (NA).

FIG. 5. Electronic microscopic images of glomeruli from diabetic mice after 25 weeks (DBA/2J and FVB/NJ strains) to 35 weeks (C57BL/6J and KK/HIJ strains) of hyperglycemia. Opposed double arrows indicate GBM. E and M designate endothelial cell and mesangial cell, respectively.

factors accounting for the heterogeneous susceptibility to diabetic nephropathy seen in diabetic patients (2-4,8,34). The present studies addressed whether genetic background also significantly affects the susceptibility to diabetic nephropathy in genetically homogenous inbred mouse strains. These studies compared the severity of the three major criteria for the diagnosis of diabetic nephropathy: albuminuria, GFR, and histopathological changes (5,18).

![](_page_5_Figure_9.jpeg)

FIG. 7. Left kidney-to-body weight ratio in diabetic inbred mice after ~25 weeks of hyperglycemia (**II**) and age-matched controls ( $\Box$ ). \*P < 0.05 (t test) vs. age-matched controls; †P < 0.05 (ANOVA) vs. all studied strains of diabetic mice, and ‡P < 0.05 (ANOVA) vs. controls in other studied strains. The age-matched controls in FVB/NJ strain were not available.

![](_page_6_Figure_1.jpeg)

![](_page_6_Figure_2.jpeg)

Diabetes was induced using multiple low-dose STZ injection, an established model for generating type 1 diabetes in mice (12,18). In contrast to single high-dose STZ, which causes massive necrosis of the pancreatic  $\beta$ -cell mass and has potential toxicity to other organs, multiple low doses of STZ initiate an insulitis similar to that observed in type 1 diabetes and moderate hyperglycemia (12,35). After 5 consecutive days of low-dose STZ injection, hyperglycemia developed in all inbred strains of mice. As previously observed (36), the average fasting blood glucose and A1C in response to low-dose STZ were strain dependent. KK/HIJ mice developed relatively low levels of hyperglycemia and A1C compared with other studied strains. In contrast, DBA/2J mice appear to be susceptible to STZ-induced diabetes, developing comparable levels of hyperglycemia with only 40 mg  $\cdot$  kg<sup>-1</sup>  $\cdot$  day<sup>-1</sup>. STZ, whereas other strains required 50 mg  $\cdot$  kg<sup>-1</sup>  $\cdot$  day<sup>-1</sup>. To maintain hyperglycemia and to accelerate the development of diabetic nephropathy, we avoided the superimposed effects of insulin treatment on the progression of diabetic nephropathy (37) and did not administer exogenous insulin to these mice. Although most mouse strains survived for >25 weeks with blood glucose levels from 300 to 600 mg/dl, those mouse strains not developing significant albuminuria appeared to exhibit a greater survival rate (data not shown). For example, a group of diabetic C57BL/6J mice lived longer than 45 weeks after the onset of hyperglycemia. In contrast, the mortality was markedly

## TABLE 2

Temporal pattern of GFR in STZ-induced diabetic and control inbred mice

Strain		Weeks after STZ or vehicle			
	Before STZ	5	15	25	
C57BL/6J					
Diabetic	$217.2 \pm 13.7$ (18)	$300.6 \pm 24.2 \ (13)^*$	$275.4 \pm 31.7 (12)^*$	$345.5 \pm 24.5 \ (9)^*$	
Control	$207.5 \pm 18.7(12)$	$306.0 \pm 47.2 (8)^{*}$	$244.3 \pm 37.9 (4)$	$325.9 \pm 17.6(5)^{*}$	
DBA/2J					
Diabetic	$220.6 \pm 22.7$ (20)	$365.5 \pm 30.4 (23)^*$	$451.9 \pm 62.8$ (8)*	$369.9 \pm 43.0 (12)^{*\dagger}$	
Control	$246.3 \pm 35.1$ (8)	$457.5 \pm 23.2$ (3)*	$452.2 \pm 121.5$ (3)*	$473.9 \pm 29.4$ (12)*	
A/J					
Diabetic	$261.2 \pm 23.6$ (16)	$246.7 \pm 27.3 (15)^{\dagger}$	$288.7 \pm 20.6 (10)^{\dagger}$	$287.5 \pm 28.6 (15)$	
Control	$271.5 \pm 47.4$ (4)	$378.5 \pm 32.8$ (4)	$393.6 \pm 13.8(4)$	$301.0 \pm 80.6$ (4)	
FVB/NJ					
Diabetic	$254.4 \pm 26.3$ (16)	$479.3 \pm 64.8 (11)^*$	$562.8 \pm 40.0 \ (4)^{*\dagger}$	$822.6 \pm 84.9 (7)^{*}$	
Control	$260.2 \pm 43.9(7)$	$245.7 \pm 58.8$ (4)	$302.4 \pm 15.2$ (4)	NA	
MRL/MpJ					
Diabetic	$239.8 \pm 72.9$ (7)	$309.6 \pm 36.2 (14)$	$491.5 \pm 69.3 (8)^{*\ddagger}$	$479.1 \pm 49.9 (7)^{*\ddagger}$	
Control	$300.3 \pm 118.6(4)$	NA	$302.3 \pm 31.7$ (4)	$272.3 \pm 25.6$ (4)	
KK/HlJ					
Diabetic	$269.3 \pm 28.9$ (12)	$353.6 \pm 24.6 \ (17)^*$	$285.6 \pm 19.5 (15)$ <sup>‡†</sup>	$315.4 \pm 37.2$ (10)	
Control	$261.4 \pm 32.6$ (8)	$286.0 \pm 26.8$ (4)	$483.0 \pm 64.6 \ (6)^{*\ddagger}$	$307.7 \pm 39.3$ (17)§	

Data are means  $\pm$  SE (µl/min) (no. of animals). NA, not available. \*P < 0.05 vs. GFR before STZ injection;  $\dagger P < 0.05$  vs. controls at same time point;  $\ddagger P < 0.05$  vs. GFR at 5 weeks after STZ; \$ P < 0.05 vs. GFR at 15 weeks after STZ.

increased in diabetic DBA/2J mice after 25 weeks of hyperglycemia, with  $\sim 40\%$  of these mice dying by this time. Whether the high mortality in diabetic mice with albuminuria is due to high incidence of infection, diabetic wasting, or cardiovascular events as in humans remains to be determined (38).

Albuminuria is a hallmark of diabetic nephropathy (5,6). In the present studies, we evaluated the development of albuminuria in diabetic inbred mice using ACR (obtained on morning spot urine) and UAE (obtained from a 24-h urine collection). Daily UAE has been a gold standard for diagnosis of albuminuria. However, environmental stress imposed on mice housed in metabolic cages, together with incomplete collection due to evaporation of urine on the cage walls may substantially influence the accuracy of urine collection in this species (26,39). Thus validation of alternative markers for 24-h UAE should facilitate studies on diabetic nephropathy and other kidney diseases using mouse models. Correlation between ACR and 24-h UAE rate as demonstrated in the present studies suggest that morning spot urine ACR can be used as an alternative for 24-h UAE in C57BL/6J, DBA/2J, A/J, MRL/MpJ, and KK/HlJ strains. This does not hold true for FVB/NJ mice.

Inbred mice exhibited distinct differences in the development of diabetic albuminuria. Diabetic DBA/2J and KK/HIJ mice appear to develop significantly more albuminuria than diabetic C57BL/6J and MRL/MpJ mice (Table 1 and Fig. 2). Although diabetic DBA/2J mice might be exposed to higher levels of blood glucose at 25 weeks after hyperglycemia (as reflected by A1C), the development of albuminuria in this strain could not be solely attributed to blood glucose level because a significant increase in albuminuria was observed after 5 weeks of hyperglycemia. At that time point and in the following weeks, fasting blood glucose and A1C were comparable with other strains. Furthermore, diabetic KK/HIJ developed albuminuria despite lower levels of A1C and hyperglycemia, suggesting that genetic factors play an important role in the development of albuminuria in these two strains. A transient increase in albuminuria was also observed in A/J and MRL/MpJ mice. This finding could correspond to human cohorts that exhibit regression of microalbuminuria to normal levels (34).

It is also significant that the widely studied C57BL/6J strain is relatively resistant to development of albuminuria despite exhibiting moderate mesangial expansion and GBM thickening. The resistance of this strain to albuminuria was not only seen in low-dose STZ-induced diabetic mice, but also in type 1 diabetes resulting from mutation of insulin 2 gene, C57BL/6-Ins2<sup>Akita</sup>/J mice (14). These results are consistent with previous studies showing that C57BL/6J mice are relatively resistant to the development of albuminuria or severe glomerulosclerosis in either STZ-induced diabetes or 5/6 nephrectomy (18–21,24,40). These previously reported levels for 24-h UAE in STZinduced hyperglycemic C57BL/6J mice were uniformly less than 100 µg/24 h (18-21), similar to the present studies. Although C57BL/6J strain has been routinely used for studying the effects of gene disruption in mice, their utility as a model for severe diabetic nephropathy may be limited.

The KK strain has previously been proposed to represent a model of diabetic nephropathy (17,41-43). The present finding of moderate albuminuria in nondiabetic KK/HIJ mice (Fig. 2) is consistent with previous studies reporting that KK mice develop albuminuria after ~10 weeks of age (17). The factors contributing to albuminuria in KK/HIJ mice have not been fully elucidated, but spontaneous glucose intolerance and hyperinsulinemia have been previously reported (17,41,44). Regardless, STZ-induced diabetes produced a further increase in albuminuria in KK/HIJ mice, demonstrating that this strain is also susceptible to diabetic albuminuria. Previous studies have also reported albuminuria in KK-A<sup>y</sup> mice, a model of type 2 diabetes (17).

Diabetic DBA/2J and KK/HlJ mice exhibited significantly

more glomerulosclerosis (Figs. 3 and 4) and GBM thickening (Figs. 5 and 6) compared with other diabetic strains. Diabetic renal lesions seen in advanced stages of human diabetic nephropathy including nodular glomerulosclerosis and arteriolar hyalinosis (5) were observed in diabetic KK/HIJ and DBA/2J mice (Fig. 3). Age-matched nonhyperglycemic control KK/HIJ mice developed less severe glomerulosclerosis than their hyperglycemic counterparts. In humans, interstitial fibrosis and tubular atrophy also develop in advanced diabetic nephropathy (5,45), however, we did not detect these lesions in the present studies.

Altered GFR represents another important indicator of the progression of diabetic nephropathy (5,46). Glomerular hyperfiltration is an early sign of renal involvement in diabetic patients (5). This is followed by a late progressive loss of renal function reflected by a decline in GFR, typically occurring after 20–25 years of type 1 diabetes in humans (5). Using a recently established approach for measuring GFR in conscious mice, we now provide the first systemic survey of GFR in inbred mouse strains over time. Glomerular hyperfiltration was observed in diabetic mice developing albuminuria including DBA/2J. KK/HlJ. and FVB/NJ mice, and in strains without significant albuminuria including C57BL/6J and MRL/MpJ mice (Fig. 8). These results suggest that hyperfiltration is not invariably linked to albuminuria. Renal hyperfiltration was associated with an increased kidney-to-body weight ratio in diabetic C57BL/6J, DBA/2J, MRL/MpJ, and KK/HlJ mice (Figs. 7 and 8).

Reduced GFR is an essential, albeit late, manifestation of diabetic nephropathy in humans. Despite severe albuminuria and renal morphological changes in diabetic DBA/2J and KK/HlJ mice, we did not detect a significant decline in GFR within the 25-week period of study compared with the baseline. In type 1 diabetic patients with nephropathy, GFR decline occurs after 15-25 years of accumulated renal injury (5). The time course of diabetic nephropathy in inbred mice remains to be defined. In a recently published study, Zheng et al. (29) reported a transgenic diabetic mouse model with FVB/N background (OVE26) developing significantly decreased GFR after  $\sim 35$ weeks of hyperglycemia (9 months). Whereas those mice developed hydronephrosis, this was not observed in the present studies. Nevertheless, a tendency for GFR to decline was observed in diabetic DBA/2J mice after 25 weeks of hyperglycemia (Fig. 8 and Table 2). Thus it is conceivable that a decline in GFR might be detected in diabetic DBA/2J and KK/HlJ mice after a more prolonged period of diabetes. Unfortunately, because of limitations in the survival of these mice, studies at later time points could not be performed.

Diabetic FVB/NJ mice were atypical in several ways. First, although 24-h UAE was significantly increased, no increase in spot ACR was observed (Fig. 2). The reason behind the disassociation between spot ACR and 24-h UAE in this strain remains to be determined; however, it is notable that these mice exhibited dramatically more polyuria than other strains, with 24-h urine volume exceeding 10 ml. For these urine samples, the concentration of albumin and creatinine was relatively low even using nondiluted samples. Second, they developed significantly greater glomerular hyperfiltration after hyperglycemia than other diabetic strains. Furthermore, we did not observe marked renal morphological changes in STZinduced hyperglycemic FVB/NJ mice within this 25-week period. Taken together, these results suggest that renal hemodynamic changes may contribute to the development of albuminuria in STZ-induced diabetic FVB/NJ mice, rather than intrinsic glomerular diseases.

In summary, the present studies provide evidence for differential susceptibility to diabetic nephropathy in inbred mice. DBA/2J and KK/HlJ mice represent strains prone to diabetic nephropathy, developing more albuminuria and correspondingly more robust renal morphological changes including mesangial expansion, nodular glomerulosclerosis, and arteriolar hyalinosis. In contrast, MRL/ MpJ, A/J, and C57BL/6J mice appear to be relatively resistant to diabetic nephropathy. They did not develop significant albuminuria or more severe renal lesions over the period of this study. Although diabetic FVB/NJ mice develop robust albuminuria, they do not develop dramatic histopathological changes over 25 weeks of hyperglycemia. Renal hypertrophy and hyperfiltration were observed in diabetic strains developing albuminuria and in diabetic mice without significant albuminuria. More prolonged hyperglycemia may be required to observe the expected decline in GFR associated with diabetic nephropathy in mice. Further studies should help establish the utility of mouse models to map genes for susceptibility to diabetic nephropathy.

#### ACKNOWLEDGMENTS

This study was supported by the Animal Models of Diabetic Complications Consortium National Institutes of Health Grant UO1-DK-61018 (to M.D.B.) and facilitated by the Vanderbilt Mouse Metabolic Phenotyping Center National Institutes of Health Grant U24-DK-59637 (to D.H. Wasserman).

We appreciate the technical help from Ellen Donnert.

## REFERENCES

- Krolewski AS, Warram JH, Christlieb AR, Busick EJ, Kahn CR: The changing natural history of nephropathy in type I diabetes. Am J Med 78:785–794, 1985
- Krolewski AS: Genetics of diabetic nephropathy: evidence for major and minor gene effects. *Kidney Int* 55:1582–1596, 1999
- Rogus JJ, Warram JH, Krolewski AS: Genetic studies of late diabetic complications: the overlooked importance of diabetes duration before complication onset. *Diabetes* 51:1655–1662, 2002
- 4. Tarnow L: Diabetic nephropathy: pathogenetic aspects and cardiovascular risk factors. *Dan Med Bull* 49:19–42, 2002
- Mauer M, Mogensen CE, Friedman EA: Diabetic nephropathy. In *Diseases* of the Kidney. 6th ed. Schrier RW, Gottschalk CW, Eds. New York, Little, Brown and Company, 1997, p. 2019–2061
- Cowie CC, Port FK, Wolfe RA, Savage PJ, Moll PP, Hawthorne VM: Disparities in incidence of diabetic end-stage renal disease according to race and type of diabetes. N Engl J Med 321:1074–1079, 1989
- 7. Freedman BI: Susceptibility genes for hypertension and renal failure. JAm Soc Nephrol 14:S192–S194, 2003
- Fava S, Hattersley AT: The role of genetic susceptibility in diabetic nephropathy: evidence from family studies. *Nephrol Dial Transplant* 17:1543–1546, 2002
- Paigen B, Schork NJ, Svenson KL, Cheah YC, Mu JL, Lammert F, Wang DQ, Bouchard G, Carey MC: Quantitative trait loci mapping for cholesterol gallstones in AKR/J and C57L/J strains of mice. *Physiol Genomics* 4:59–65, 2000
- Sugiyama F, Churchill GA, Higgins DC, Johns C, Makaritsis KP, Gavras H, Paigen B: Concordance of murine quantitative trait loci for salt-induced hypertension with rat and human loci. *Genomics* 71:70–77, 2001

- Sugiyama F, Yagami K, Paigen B: Mouse models of blood pressure regulation and hypertension. Curr Hypertens Rep 3:41–48, 2001
- Like AA, Rossini AA: Streptozotocin-induced pancreatic insulitis: new model of diabetes mellitus. *Science* 193:415–417, 1976
- Tabatabaie T, Waldon AM, Jacob JM, Floyd RA, Kotake Y: COX-2 inhibition prevents insulin-dependent diabetes in low-dose streptozotocin-treated mice. *Biochem Biophys Res Commun* 273:699–704, 2000
- 14. Wang J, Takeuchi T, Tanaka S, Kubo SK, Kayo T, Lu D, Takata K, Koizumi A, Izumi T: A mutation in the insulin 2 gene induces diabetes with severe pancreatic beta-cell dysfunction in the Mody mouse. *J Clin Invest* 103:27–37, 1999
- Surwit RS, Kuhn CM, Cochrane C, McCubbin JA, Feinglos MN: Dietinduced type II diabetes in C57BL/6J mice. *Diabetes* 37:1163–1167, 1988
- Sharma K, McCue P, Dunn SR: Diabetic kidney disease in the db/db mouse. *Am J Physiol Renal Physiol* 284:F1138–F1144, 2003
- Okazaki M, Saito Y, Udaka Y, Maruyama M, Murakami H, Ota S, Kikuchi T, Oguchi K: Diabetic nephropathy in KK and KK-Ay mice. *Exp Anim* 51:191–196, 2002
- Breyer MD, Bottinger E, Brosius FC III, Coffman TM, Harris RC, Heilig CW, Sharma K: Mouse models of diabetic nephropathy. J Am Soc Nephrol 16:27–45, 2005
- Susztak K, Bottinger E, Novetsky A, Liang D, Zhu Y, Ciccone E, Wu D, Dunn S, McCue P, Sharma K: Molecular profiling of diabetic mouse kidney reveals novel genes linked to glomerular disease. *Diabetes* 53:784–794, 2004
- 20. Huang W, Gallois Y, Bouby N, Bruneval P, Heudes D, Belair MF, Krege JH, Meneton P, Marre M, Smithies O, Alhenc-Gelas F: Genetically increased angiotensin I-converting enzyme level and renal complications in the diabetic mouse. *Proc Natl Acad Sci U S A* 98:13330–13334, 2001
- Zheng F, Striker GE, Esposito C, Lupia E, Striker LJ: Strain differences rather than hyperglycemia determine the severity of glomerulosclerosis in mice. *Kidney Int* 54:1999–2007, 1998
- Glazier AM, Nadeau JH, Aitman TJ: Finding genes that underlie complex traits. Science 298:2345–2349, 2002
- 23. Churchill GA, Airey DC, Allayee H, Angel JM, Attie AD, Beatty J, Beavis WD, Belknap JK, Bennett B, Berrettini W, Bleich A, Bogue M, Broman KW, Buck KJ, Buckler E, Burmeister M, Chesler EJ, Cheverud JM, Clapcote S, Cook MN, Cox RD, Crabbe JC, Crusio WE, Darvasi A, Deschepper CF, Doerge RW, Farber CR, Forejt J, Gaile D, Garlow SJ, Geiger H, Gershenfeld H, Gordon T, Gu J, Gu W, de Haan G, Hayes NL, Heller C, Himmelbauer H, Hitzemann R, Hunter K, Hsu HC, Iraqi FA, Ivandic B, Jacob HJ, Jansen RC, Jepsen KJ, Johnson DK, Johnson TE, Kempermann G, Kendziorski C, Kotb M, Kooy RF, Llamas B, Lammert F, Lassalle JM, Lowenstein PR, Lu L, Lusis A, Manly KF, Marcucio R, Matthews D, Medrano JF, Miller DR, Mittleman G, Mock BA, Mogil JS, Montagutelli X, Morahan G, Morris DG, Mott R, Nadeau JH, Nagase H, Nowakowski RS, O'Hara BF, Osadchuk AV, Page GP, Paigen B, Paigen K, Palmer AA, Pan HJ, Peltonen-Palotie L, Peirce J, Pomp D, Pravenec M, Prows DR, Qi Z, Reeves RH, Roder J, Rosen GD, Schadt EE, Schalkwyk LC, Seltzer Z, Shimomura K, Shou S, Sillanpaa MJ, Siracusa LD, Snoeck HW, Spearow JL, Svenson K, Tarantino LM, Threadgill D, Toth LA, Valdar W, de Villena FP, Warden C, Whatley S, Williams RW, Wiltshire T, Yi N, Zhang D, Zhang M, Zou F: The Collaborative Cross, a community resource for the genetic analysis of complex traits. Nat Genet 36:1133-1137, 2004
- Ma LJ, Fogo AB: Model of robust induction of glomerulosclerosis in mice: importance of genetic background. *Kidney Int* 64:350–355, 2003
- 25. Hartner A, Cordasic N, Klanke B, Veelken R, Hilgers KF: Strain differences in the development of hypertension and glomerular lesions induced by deoxycorticosterone acetate salt in mice. *Nephrol Dial Transplant* 18: 1999–2004, 2003

- 26. Qi Z, Whitt I, Mehta A, Jin J, Zhao M, Harris RC, Fogo AB, Breyer MD: Serial determination of glomerular filtration rate in conscious mice using FITC-inulin clearance. Am J Physiol Renal Physiol 286:F590–F596, 2004
- 27. Ma LJ, Jha S, Ling H, Pozzi A, Ledbetter S, Fogo AB: Divergent effects of low versus high dose anti-TGF-beta antibody in puromycin aminonucleoside nephropathy in rats. *Kidney Int* 65:106–115, 2004
- Mattix HJ, Hsu CY, Shaykevich S, Curhan G: Use of the albumin/creatinine ratio to detect microalbuminuria: implications of sex and race. J Am Soc Nephrol 13:1034–1039, 2002
- Zheng S, Noonan WT, Metreveli NS, Coventry S, Kralik PM, Carlson EC, Epstein PN: Development of late-stage diabetic nephropathy in OVE26 diabetic mice. *Diabetes* 53:3248–3257, 2004
- 30. Ron D: Proteotoxicity in the endoplasmic reticulum: lessons from the Akita diabetic mouse. J Clin Invest 109:443–445, 2002
- Oyadomari S, Koizumi A, Takeda K, Gotoh T, Akira S, Araki E, Mori M: Targeted disruption of the Chop gene delays endoplasmic reticulum stress-mediated diabetes. J Clin Invest 109:525–532, 2002
- Mogensen CE, Andersen MJ: Increased kidney size and glomerular filtration rate in early juvenile diabetes. *Diabetes* 22:706–712, 1973
- Osterby R, Gundersen HJ: Glomerular size and structure in diabetes mellitus: I. Early abnormalities. *Diabetologia* 11:225–229, 1975
- 34. Perkins BA, Ficociello LH, Silva KH, Finkelstein DM, Warram JH, Krolewski AS: Regression of microalbuminuria in type 1 diabetes. N Engl J Med 348:2285–2293, 2003
- Rossini AA, Williams RM, Appel MC, Like AA: Complete protection from low-dose streptozotocin-induced diabetes in mice. *Nature* 276:182–184, 1978
- Iwatsuka H, Shino A, Taketomi S: Streptozotocin resistance of the genetically diabetic KK mouse. *Diabetes* 23:856–857, 1974
- 37. Ferrannini E, Galvan AQ, Gastaldelli A, Camastra S, Sironi AM, Toschi E, Baldi S, Frascerra S, Monzani F, Antonelli A, Nannipieri M, Mari A, Seghieri G, Natali A: Insulin: new roles for an ancient hormone. *Eur J Clin Invest* 29:842–852, 1999
- Yuyun MF, Adler AI, Wareham NJ: What is the evidence that microalbuminuria is a predictor of cardiovascular disease events? *Curr Opin Nephrol Hypertens* 14:271–276, 2005
- Champy MF, Selloum M, Piard L, Zeitler V, Caradec C, Chambon P, Auwerx J: Mouse functional genomics requires standardization of mouse handling and housing conditions. *Mamm Genome* 15:768–783, 2004
- 40. Fujita H, Haseyama T, Kayo T, Nozaki J, Wada Y, Ito S, Koizumi A: Increased expression of glutathione S-transferase in renal proximal tubules in the early stages of diabetes: a study of type-2 diabetes in the Akita mouse model. *Exp Nephrol* 9:380–386, 2001
- Reddi AS, Camerini-Davalos RA: Hereditary diabetes in the KK mouse: an overview. Adv Exp Med Biol 246:7–15, 1988
- 42. Reddi AS, Wehner H, Khan MY, Camerini-Davalos RA: Kidney disease in KK mice: structural, biochemical and functional relationships. Adv Exp Med Biol 246:135–145, 1988
- Wehner H: Sequence of changes in KK mouse nephropathy. Adv Exp Med Biol 246:147–156, 1988
- 44. Suto J, Matsuura S, Imamura K, Yamanaka H, Sekikawa K: Genetic analysis of non-insulin-dependent diabetes mellitus in KK and KK-Ay mice. *Eur J Endocrinol* 139:654–661, 1998
- 45. Fioretto P, Steffes MW, Sutherland DE, Mauer M: Sequential renal biopsies in insulin-dependent diabetic patients: structural factors associated with clinical progression. *Kidney Int* 48:1929–1935, 1995
- Anderson S, Brenner BM: Pathogenesis of diabetic glomerulopathy: hemodynamic considerations. *Diabete Metab Rev* 4:163–177, 1988