

MED-SURG EXAM

- 1. A patient with congestive heart failure should be carefully watched for signs of:
 - a. Fatique
 - b. Uremia
 - c. Pulmonary edema
 - d. Heart block
 - e. Bleeding
- 2. In the absence of cardiac enlargement or other symptoms, a blood pressure reading of 210/100 mm Hg would suggest a diagnosis of:
 - a. Anxiety
 - b. Sclerosis of the aorta
 - c. Hypertensive heart disease
 - d. Essential hypertension
 - e. Clinomania
- 3. In the cardiac cycle, what is the period of relaxation called?
 - a. Systole
 - b. Diastole
 - c. Ventricular pressure
 - d. Semilunar valve closure
- 4. In myocardial infarction, there is:
 - a. Necrosis of the heart muscle
 - b. Inflammation of the heart muscle
 - c. Narrowing of the coronary veins
 - d. Thickening of the endocardium
- 5. The first drug of choice for a patient with known cardiac disease who complains of chest pain is:
 - a. Nitroglycerin
 - b. Oxygen
 - c. Morphine
 - d. Lanoxin
- 6. When an anticoagulant is given to a patient, which of the following procedures must be performed daily?
 - a. Sedimentation rate
 - b. Urinalysis
 - c. Complete blood count
 - d. Prothrombin time



- 7. A patient receiving anticoagulants should be watched for:
 - a. Dyspnea
 - b. Bleeding
 - c. Hypotension
 - d. Vomiting
- 8. A complication of acute myocardial infarction is:
 - a. Pulmonary edema
 - b. Pulmonary embolism
 - c. Ventricular fibrillation
 - d. All of the above
- 9. Most respiratory tract infections are probably caused by:
 - a. Fungi
 - b. Viruses
 - c. Bacteria
 - d. Spirochetes
 - e. Protozoa
- 10. A patient with an asthmatic attack should not be given:
 - a. Epinephrine
 - b. Benadryl
 - c. Potassium iodide
 - d. Ether inhalation
 - e. Morphine
- 11. Status asthmaticus refers to:
 - a. A chronic condition
 - b. An initial asthmatic attack
 - c. A continuous asthmatic attack
 - d. An asthmatic attack caused by humidity change
- 12. Drugs used in the treatment of asthma include:
 - a. Epinephrine
 - b. Ephrine sulfate
 - c. Aminophylline
 - d. All of the above
- 13. Drugs used in the treatment of peptic ulcer will:
 - a. Decrease gastric secretions
 - b. Lower the acidity of gastric fluid
 - c. Reduce gastric motility
 - d. All of the above



- 14. Peptic ulcer disease is believed to occur when:
 - a. The patient is under stress
 - b. The patient has used alcohol
 - c. Aggressive forces overwhelm natural protective mechanisms
 - d. The patient follows a bland diet
- 15. Acute mechanical intestinal obstruction may cause all of the following EXCEPT:
 - a. Diarrhea
 - b. Nausea
 - c. Vomiting
 - d. Distension
 - e. Colicky pain
- 16. Hepatocellular disease tend to decrease the serum level of:
 - a. Albumin
 - b. Prothrombin
 - c. Fibrinogen
 - d. All of the above
- 17. Blood chemistry tests useful in evaluating renal function include:
 - a. Blood urea nitrogen (BUN)
 - b. Nonprotein nitrogen (NPN)
 - c. Serum creatinine level
 - d. All of the above
- 18. Classic symptoms of patient with diabetes include increased:
 - a. Thirst
 - b. Urinary output
 - c. Appetite
 - d. All of the above
- 19. Fractional urine specimens are collected from diabetic patients to determine:
 - a. The amount of albumin in the urine
 - b. The amount of hemoglobin in the urine
 - c. The time of day when excretion of sugar is heaviest
 - d. The amount of fat needed by the patient
- 20. The oliguric-anuric phase of acute renal failure means the patient's urine output is:
 - a. More than 400 cc/day
 - b. Less than 400 cc/day
 - c. More than 1,000 cc/day
 - d. Varies depending of fluid intake



21. Digi	talis shoul	d NOT be	e administered	to a	patient who:
----------	-------------	----------	----------------	------	--------------

- a. Has nausea and is vomiting
- b. Complains of headache
- c. Both
- d. Neither
- 22. Narcotic analgesics work by:
 - a. Inhibiting the syntheses of prostaglandins
 - b. Binding to specific opiate receptors throughout the CNS
 - c. Binding to receptor sites in the periphery
 - d. Inhibiting the synthesis fatty acid substances
- 23. Acetaminophen's most serious potential adverse effect is:
 - a. Tinnitus
 - b. Hepatotoxicity
 - c. Respiratory depression
 - d. Psychotomimetic effects
- 24. Traction is used in order to:
 - a. Reduce a fracture
 - b. Maintain the alignment of the bones
 - c. Both
 - d. Neither
- 25. The patient on continuous tube feedings still requires that his feeding tube be checked for placement and irrigated for patency at least once a shift.
 - a. True
 - b. False

Signature	 Date	 e