PYROCIDE® Concentrated Aqueous Spray 74404

[Commercial Section of Label]

[] or () Indicates alternate or optional wording

[Contains Pyrethrins, a botanical insecticide.]

A micro-emulsion, water-base spray

Kills Stable Flies, Horse Flies, Face Flies, Deer Flies, House

Flies, Mosquitoes, and Gnats

Kills Black Widow Spiders and Black Carpenter Ants

Pyrethins kills listed crawling and flying insects in the hours

schools, hospitals.

Kills Fleas and Ticks indoors and around pet areas

Kills Deer Ticks and other Ixodid species that may carry and transmit Lyme Disease

For use in Animal Quarters, Dairies, Milk Rooms, Milking

Parlors, Carving Areas, Dairy and Hog Barns

For use on Beef Cattle, Dairy Cattle, Horses, Hogs, Poultry and other listed livestock

Kills Mosquitoes including the Culex species that may carry and transmit the West Nile Virus

ACTIVE MOREDIENTS

Pyrethrins 1.00%
Piperonyl butoxide 10.00%
HER NGREDIENTS 89.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice.
	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to by a poison control center or doctor.
	Do not give anything by mouth to an unconscious person.
IF ON SKIN:	Take off contaminated clothing.
	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For	
information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.	

Manufactured by



8810 Tenth Avenue North Minneapolis, MN 55427

EPA Reg. No. 1021-1799

Net Contents _____

EPA Est. No. 1021-MN-2

74404_1021-1799_0512 Page 1 of 18

^{* (}butylcarbityl) (6-propylpiperonyl) ether and related compounds
PYROCIDE® is a registered trademark of McLaughlin Gormley King Company

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE

Mixers, loaders, applicators, and other handlers must wear:

- Long-sleeve shirt,
- Long pants,
- Shoes and socks.
- Chemical resistant gloves, selection category A.

In addition to the above PPE, applicators using high pressure handward in an en - A dust/mist filter with MSHA/NIOSH approval number prefix TC 12C or wear at least a NIOSH-approved respirator with:

- Any R, R, or HE filter.

In addition to the above PPE, applicators using hand held foggers must wear a half-face, full-face, or hood-style NIOSHapproved respirator with:

- A dust/mist filtering cartridge (MSHA/NIØSH approval number prefix TC-12C) or A canister approved for pesticides (MSHA/NIØSH approval number prefix TC-14G), or
- A cartridge or canister with any M, R, B or NE filter

SAPETY REQUIREMENTS

It no such instructions for washables exist, use detergent and hot water. Keep and Follow manufacturer's instructions to cleaning maintaining PPE. wash PPE separately from other laundly

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

USER SAFETY RECCOMMENDATIONS

- User should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product may contaminate water through runoff. This product has a potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or rinsate.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

PHYSICAL OR CHEMICAL HAZARDS

Do not use this product in or on electrical equipment because of shock hazard.

[Pyrocide® Concentrated Aqueous Spray 74404] contains the botanical insecticide, Pyrethrins, which provides effective kill of listed insects upon direct contact with an added benefit of flushing these insects to aid in a more complete kill.]. This waterbased formula can be used on listed flying and crawling insects indoors and outdoors.

[Pyrocide® Concentrated Aqueous Spray 74404] can be used to kill fleas and ticks hidden in carpets, rugs, upholstery, draperies, floor cracks and cabinets. Also Kills Deer Ticks and other Ixodid species that may carry and transmit Lyme Disease.

74404_1021-1799_0512 Page 2 of 18 [Pyrocide® Concentrated Aqueous Spray 74404] can be used in the home, schools, hospitals, restaurants and other listed areas to kill listed flying and crawling insect pests kill. Pyrethrins have quick killing ability with no persistent residue. [Pyrocide® Concentrated Aqueous Spray 74404] can be used to reduce annoyance of mosquitoes, including the Culex species, which may carry and transmit the West Nile Virus, outdoors around homes, recreational and other listed areas. Ideal for use prior to outdoor activities.

[Pyrocide® Concentrated Aqueous Spray 74404] can be used outdoors with the Lock N Spray system to kill pests in lawns and on home foundations.

[Pyrocide® Concentrated Aqueous Spray 74404] can be used in food and non-food areas of food processing plants, canneries, industrial installations, bakeries, rice and wheat mills, tobacco warehouses, grain elevators, warehouses, storage sites and other listed areas to kill, on contact, listed crawling posts and is a fast, effective product for flying insect control.

[Pyrocide® Concentrated Aqueous Spray 74404] provides rapid kill of insects without objectionable residues.

FOR THE KILL OF INSECTS:

Almond Moths
Angoumois Grain Moths

Ants
Bedbugs

Black Carpenter Ants

Black Flies

Black Widow Spiders

Blood Sucking Lice

Boxelder Bugs Brown Dog Ticks

Cadelles
Carpet Beetles
Cat Fleas

Cellar Spiders
Cereal Beetles

Cheese Mites
Cheese Skippers
Chocolate Moths

Cigarette Beetles Clothes Moths Confused Flour Beatles

Dark Mealworms
Darkling Beetles

Deen Flies Deen Ticks

Dermestid Beetles

Dog Fleas

Dried Fruit Beetles Drugstore Beetles

Earwigs
Face Flies
Fannia Flies
Firebrats

Flat Grain Beetles

Fruit Flies Gnats Grain Mites Granary Weevils

Horn Flies Hornets Hørse Flies Houseflies

Indian Meal Moths Khapra Beetles

Lesser Grain Borers Lesser Houseflies Lice

Mediterranean Flour Moths

Moths

Merchant Grain Beetles Midges

Miller Moths Millipedes

Mosquitoes – including the Culex species that

may carry and transmit West Nile Virus.

Palmetto Bugs Pillbugs Poultry Lice Red Flour Beetles Rice Flour Beetles

Rice Weevils
Rusty Grain Beetles

Saw-toothed Grain

Beetles Sheep Ticks Silverfish Skipper Flies Small Flying Moths

Sowbugs Spider Beetles Stable Flies

Ticks that may carry and transmit Lyme disease

Tobacco Moths Trogoderma Beetles Warehouse Beetles

Wasps
Waterbugs
Yellow Jackets
Yellow Mealworms

INDOOR AREAS for use in:

Animal Areas

Cockroaches

Animal Control Centers Animal Quarters Animal Shelters Apartments

Atriums
Attics
Auditoriums
Automobiles
Bakeries
Bars

Bedding Bedrooms Beef Barns Beverage Plants

Boats Bookcases

Basements

Bathrooms

Bottling Plants Breweries

Breweries
Brewery Warehouses
Buildings Housing Animals
Buildings Housing Poultry

Buses
Cabinets
Cabins
Cafeterias
Calving Barns
Campers
Canneries
Carpets

Cat Sleeping Quarters

Cattle Barns
Chests
Churches
Circus Tents
City Missions

Closed Porches

Closets

Clothes Storage Club Houses Commercial Buildings

Condominiums Crawl Spaces Credenzas Crematoriums Dairies

Dairy and Livestock Barns Day Care Centers Delicatessens

Dens

Department Stores

Diners Dog Houses

Dog Sleeping Quarters

Dormitories

Draperies Dressers

Dried Fruits in Storage

Drive-Ins Drugstores Dumpsters Dwellings

Egg Processing Plants Fabric Storage Areas

Factories
False Ceilings
Farm Structures

Floors
Food Processing Plants
Food Storage Areas
Freight Containers

Page 3 of 18

Fruit Packing Sheds Funeral Parlors

Furniture

74404_1021-1799_0512

Garages Garbage Bins Garbage Cans **Garbage Compactors**

Gas Stations Gazebos

Goat Holding Areas **Grain Elevators** Grain Handling Equip **Grain Harvesting Equipment**

Grain Mills Grain Storage Granaries **Grocery Stores** Hog Barns Homes Horse Stables

Horse Stables and Barns

Horse Trailers Hospices Hospitals Hotels

Household contents

Houses

Indoor Eating Buildings Industrial Installations **Industrial Plants**

Institutional Dining Areas Institutions

Jails Kennels **Kitchens** Laboratories Laundromats

Lavatories

Libraries Liquor Stores Livestock Barns Livestock Hauling Equipment

Livestock Holding Areas

Living Rooms Loafing Sheds Locker Rooms Marinas Mattresses Mausoleums

Meat Packing Plants Mess Halls Milk Houses

Milk Rooms < Milk Sheds

Milking Rooms Mini Storages Andayine SalidoM Mobile Hornes Morgues

Mortuaries Matels.

Movie Theaters Museums **Nursing Homes** Office Buildings Other Public Buildings **Outdoor Eating Buildings**

Pantries

Parking Garages Pet Bedding Pet Carriers

Pet Grooming Parlors

Pet Sleeping Areas

Pet Stores Play Houses Play Rooms Pool Halls

Porches Poultry Houses

Poultry Plants

Pounds Prisons Rublic Storages Quonset Huts

Rabbit Processing Plants

Rallroad Cars

Recreational Vehicles Research Animal Quarters

Restaurants Retail Stores Rice Mills Rooms Rugs Schools Seed Storage

Sheep Holding Areas

Ship Holds Ships **Smoke Shops** Solariums

Sport Stadiums Stable Bins Stables Storage Areas Stored Food, bags Stored Products

Stores

Sun Porches Sun Rooms Swine Houses Taverns Tents

Textile Mills, Warehouses

Theaters

Tobacco Factories Tobacco Warehouses

Tool Sheds Trailers Trains

Transportation Vehicles

Tree Houses

Tree Nuts Storage Areas

Truck Trailers Trucks **Tunnels** Utilities **Utility Rooms Vending Machines**

Verandas

Zoos

Veterinary Hospitals and

Clinics Voids in Attics

Warehouses Washrooms Wheat Mills Wholesale Stores Wine Cellars Wineries

OUTDOOR AREAS for uses in:

Allevs **Amphitheaters Amusement Parks**

Ant Hills **Back Yards Balconies Baseball Fields Beaches** Bridges Camp Sites Campgrounds Carwashes Circuses

Construction Sites

Corrals

Courtyards Curbs Decks

Drive-In Restaurants Drive-In Theaters Dumpsters Fair Grounds Feedlots Front Yards Game Fields Garbage Dumps Gazebos Grandstands

Junk Yards Kennels

Lanais Levees Manholes Manure Piles Marinas **Outdoor Eating** Establishments Parking Lots Parks **Patios** Playgrounds

Ranges Recreational Areas

Rest Areas Roadsides

Rooftops Sanitary Sewers Side Walks Side Yards Stadiums **Tennis Courts** Truck Stops Underpasses Utility Room

Outdoor Weigh Stations Wild Animal Parks

Wood Piles

Zoos

74404_1021-1799_0512 Page 4 of 18 FOR USE ON:

Pullets Beefalo Donkeys Lambs Birds **Ducks** Llamas Quail Buffalo Emus Mules Ratites Calves Exotics such as Ostrich Roosters European Red Deer Sheep Cattle – Beef Oxen Cattle - Dairy Geese **Pheasants** Swine Chickens Horses Durkeys Pigs

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read all directions completely before use. For best results, follow directions for specific use areas. Avoid contamination of food and feedstuffs.

USE RESTRICTIONS:

Do not enter or allow others to enter until vapors, mists, and aerosols have dispersed, and the treated area has been thoroughly ventilated.

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Only protected handlers may be in the area during applications.
- Do not remain in treated area. Exit area immediately and remain outside the treated area until aerosols, vapors, and/or mists have dispersed.
- Remove or cover exposed food and drinking water before application.
- Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces, or wash them before use.
- Do not apply space spray while food processing is in operation. Prior to space spray application, cover or remove food. Prior to space spray application, cover or remove all food processing surfaces or clean after treatment and before use.
- Do not enter or allow others to enter until sprays have dried.
- Remove pets and birds and cover fish aquariums before spraying. Turn aquarium systems off
- Do not make applications during rain
- · Do not water the treated area to the point of run-off
- Do not wet articles to point of runoff or drip.
- Do not use treated article until spray has dried.
- Do not use in aircraft.
- When used in dairy barns or facilities: Close milk bulk lids to prevent contamination from spray and from dead or falling insects.
 Remove or cover milking utensils before application. Wash teats of animals before milking.
- Except when applying to sewers or drains as instructed on this label, do not apply directly to sewers or drains, or to any area like a gutter where drainage to sewers storm drains, water bodies, or aquatic habitat can occur.

This concentrated product may be diluted with water. When making dilutions, add concentrate to the tank and add water. Agitate well while adding water and apply as directed.

0.05% Pyrethrins – add 1 part concentrate to 18 parts water or 6.5 fluid ounces concentrate / gallon water 0.10% Pyrethrins – add 1 part concentrate to 9 parts water or 13 fluid ounces concentrate / gallon water 0.30% Pyrethrins – add 1 part concentrate to 2 parts water or 38.5 fluid ounces concentrate / gallon water 0.50% Pyrethrins – add 1 part concentrate to 1 parts water or 64 fluid ounces concentrate / gallon water

Use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food must be removed or covered during treatment. All food processing surfaces must be covered during treatment or thoroughly cleaned before using.

Use Lock 'N Spray System outdoors only.

For maximum effectiveness, follow the dilution chart and use of a combination of spot, surface, and fogging treatment is recommended.

74404_1021-1799_0512 Page 5 of 18

SPACE SPRAYING: Use with conventional mechanical or compressed air equipment to kill exposed accessible stages of: Flies (including Fruit Flies), Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, Small Flying Moths, Fleas, Gnats, Wasps, Hornets, Clover Mites, Cheese Mites, Cheese Skippers, Box elder Bugs, Earwigs, Ants and the following common Stored Product Pests: Granary Weevils, Rice Weevils, Confused Flour Beetles, Saw-Toothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths, Tobacco Moths, Yellow Meal Worms, Dark Meal Worms, Grain Mites, Cadelles, Red Flour Beetles and other listed insects. Close Room and shut off air conditioning or ventilating equipment, Apply at the rate of 1 – 10 fluid ounces per 1000 cubic feet of space. Direct Spray toward upper corners of room and celling Leave room closed for ½ hour and ventilate thoroughly before re-entry.

SURFACE SPRAYING: To kill Cockroaches, Ants, Silverfish, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Saw-toothed Grain Beetles, Spiders Beetles, Cigarette Beetles and other listed insects use a sprayer adjusted to a coarse wet spray. Mix well and apply as a coarse wet spray using approximately 1 gallon of spray mixture per 750 square feet of surface. Direct the spray first into all cracks and crevices, behind all equipment, cupboards, obstructions or dark harborage areas, walls, floors, underneath sinks, behind pipes, beneath stoves and refrigerators, around garbage cans, cabinets, along baseboards, door sills, window sills, door and window frames and floors, around and on drains, pipes, plumbing, behind bookcases, storage and other utility installation areas and in all places that harbor insects to provide maximum penetration and impingement on the insects. Contact as many insects as possible. Do not allow spray to contact stored food. To kill ants, treat doors, around window trames and other areas of entry. Spray on ants whenever possible.

CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply liberally to empty chests, bureaus and other storage areas directing the spray into cracks, joints, and crevices.

CARPET BEETLES: Directly soral as many insects as possible. Spray all surfaces of upholstered furniture. Where possible, both top and bottom of rugs and carpets should be sprayed thoroughly. Also treat shelves, closets, areas under furniture, floors, walls, windowsills, baseboards and wherever these insects are seen or suspected.

BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, floors, walls, furniture and other infested areas.

FOR USE IN CANNERIES: Thoroughly clean floors, conveying equipment and walls, with special attention to cracks and crevices, prior to bringing in produce. Spray the entire space inside the cannery including walls, ceiling and floors, paying special attention to forcing the spray into all cracks and crevices for killof ants cockroaches, silverfish, crickets, spiders, fruit flies, moths, house flies, cheese mites and other listed insects.

FOR USE IN FOOD AND NONFOOD AREAS OF FOOD PROCESSING PLANTS, Industrial installations, Bakeries, Rice and Wheat Mills, Restaurants, Tobacco Warehouses, Grain Elevators and Warehouses: Spray as directed to killisted flying and crawling insects. The use of this product in food processing or food handling establishments should be confined to time periods when the plant is not in operation. Food should be covered or removed during treatment.

DRIED FRUIT PRODUCTS: In storage or being processed for kill of,: Almond Moths, Ants, Cereal Beetles, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Earwigs, Flies, Fruit Flies, Fungus Gnats, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Saw-toothed Grain Beetles, Silverfish, Dried Fruit Beetles, Warehouse Beetles, Khapra Beetles and other listed insects. Do not apply to fruit directly. Apply as a fine mist or fog in the air above and around the trays, bins, and shelves at the rate of 1 to 10 fluid ounces per 1000 cubic feet of space.

TREATMENT OF STORED FOOD AREAS: Foods in multi-walled bags, cloth bags or other packaging stored in such areas as, listed areas for the kill of accessible adult stages of: Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Dark Mealworms, Drugstore Beetles, Earwigs, Firebrats, Flies, Fruit Flies, Fungus Gnats, Grain Mites, Granary Weevils, Hide Beetles, Indian Meal Moths, Mediterranean Flour Moths, Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, Red Flour Beetles, Rice Weevils, Saw-toothed Grain Beetles, Silverfish, Small Flying Moths, Spiders, Spider Beetles, Yellow Mealworms, Dried Fruit Beetles, Warehouse Beetles, Rusty Grain Beetles, Lesser Grain Beetles, Khapra Beetle and other listed pests. Infested stored products should be destroyed, fumigated or treated by other effective methods using an approved product intended for this purpose. As noted above, disperse first by directing spray into cracks and crevices, corners, pallets and around stacks or packaged goods hitting as many insects as possible and then apply as a space spray at a dosage of 1 to 10 fluid ounces per 1000 cubic feet of room space. Repeat application as necessary for insect kill. Do not remain in treated area and ventilate before re-entry.

74404_1021-1799_0512 Page 6 of 18

ON PEANUTS, NUTS, BEANS, SEEDS IN BULK, BAGS OR OTHER PACKAGING: For the kill of accessible stages of Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Dark Mealworms, Drugstore Beetles, Earwigs, Firebrats, Flies, Fruit Flies, Fungus Gnats, Grain Mites, Granary Weevils, Hide Beetles, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Rice Weevils, Saw-toothed Grain Beetles, Silverfish, Small Flying Moths, Spiders, Spider Beetles, Yellow Mealworms, Dried Fruit Beetles, Warehouse Beetles, Rusty Grain Beetles, Lesser Grain Beetles, Kapra Beetle and other listed insects apply to the surface of stored or bagged products. Also spray walls, floors and other surfaces of bins, storage and handling areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Then apply as a space spray by directing the spray toward the ceiling and upper corners of the area and behind any obstructions.

FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS: Apply with conventional mechanical or compressed air equipment, at the rate of 1 to 10 fluid ounces per 1000 cubis feet of room space. Apply the spray into all cracks and crevices in woodwork, walls, and floors, underneath sinks, behind tipes and in all places that shelter insects. Contact as many insects as possible. Repeat as necessary. Before use, exposed food products and packaging materials must be covered or removed. After use, equipment and utensils must be washed with an effective cleaning compound and rinsed with potable water. Use as a space spray only when plant is not in operation.

FOR IN TRANSIT PROTECTION: To kill accessible stages of listed insects, clean and spray truck beds, boxcars, ships holds and other listed transit equipment before loading. The roughly treat walls and floors and spray each layer of carton or containers lightly as stacked. Apply at the vate of 1 pallow per 750 square feet.

FOR USE IN STORAGE FACILITIES AND EQUIPMENT:

SANITATION MEASURES: This product can be used as a surface spray in warehouse bins and trucks, cargo ships, mills, bin hoppers, elevators and conveying equipment as a clean up prior to use. In mills, and elevators, all grain-infested accumulations should be removed from the bin hoppers. All storage and conveying equipment should be thoroughly cleaned by sweeping out the waste grain, cobwebs and other debris from the walls and rafters as well as on the floor and door frames with special attention to material lodged in the cracks and crevices. All debris should be removed and disposed of in outside trash.

FOR FARMS, particular attention should be given to cleaning up around used feed and grain bags, grain residues from wagons, harvesting equipment and feed troughs. Newly harvested grain should not be placed in the same bin with carry-over grain and all carry-over grain stocks that are not treated with grain protectant should be fumigated or treated with EPA approved products [such as] [brand name]. These cleaning operations should be done within two or three weeks before harvest.

After above sanitation measures have been employed spray as directed.

FOR USE IN ANIMAL QUARTERS, Barns, Dairies, Milking Parlors, Milking Rooms, Poultry Houses and other listed animal areas: For rapid kill of Flies, Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, Gnats and other listed flying insects, follow dilution chart and spray at the rate of 1 to 10 fluid ounces per 1000 cubic feet of space, directing the spray toward the ceiling and upper corners and not directly towards animals or birds. For best results, close doors and windows before spraying and keep them closed for 10 to 15 minutes. Applicator should vacate the treated area and ventilate it prior to returning.

FOR USE ON LIVESTOCK follow dilution chart and apply as directed:

To protect beef cattle, dairy cattle, horses and other livestock from attacks and to kill and repel Stable Flies, Horse Flies, Deer Flies, House Flies, Horn Flies, Lesser House Flies, Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, and Gnats, spray about 3 seconds on each side of animal, being careful to spray back, withers and forelegs thoroughly.

To protect from Face Flies, spray the face and hair or spray on a cloth and rub on the face, being certain to avoid contact with eyes. Repeat treatment as necessary when flies are troublesome.

For effective kill of biting and sucking lice on cattle, horses, sheep, goats and hogs, spray to thoroughly wet hair of the animal including the head and brush of the tail. Repeat treatment in 10 days to kill newly hatched lice.

74404_1021-1799_0512 Page 7 of 18

To kill poultry lice and exposed stages of Darkling Beetles, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

To kill bedbugs and mites on poultry and in poultry houses, spray crevices of roost poles, cracks in walls and cracks in nests where the bedbugs and mites hide. This should be followed by spraying over the birds with a fine mist.

To kill sheep "tick" or ked, thoroughly wet all portions of the body by spraying directly into the wool. Treat at a rate to sufficiently wet the animal.

To kill fleas and ticks on livestock and to obtain protection against reinfestation, spray to sufficiently wet the animal.

FOR USE IN BARNS, DAIRIES, MILKING PARLORS, MILKING ROOMS, POULTRY HOUSES and other listed animal housing areas, follow dilution chart and apply as directed. Do not use in milk rooms where raw milk may be exposed. Repeat application in one to two days, if necessary.

0.10% Pyrethrins – Add 1 part Concentrate to 9 parts water 0.50% Pyrethrins – Add 1 part Concentrate to 1 part water

FOR USE WITH HANDHELD MECHANICAL FOGGERS to kill Houseflies, Horn Flies, Stable Flies, Horse Flies, Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, Gnats and other listed insects. Dilute concentrate and apply as a fine mist of fog. Direct spray toward the ceiling and upper corners of the room until the area is filled with mist, about 2 fluid ounces per 1000 cubic feet of space. For best results, close doors and windows before spraying, and keep them closed for 10 to 15 minutes after spraying. Applicator should vacate the treated area and ventilate it before reoccupying.

FOR USE THROUGH AUTOMATIC\SPRAYING SYSTEMS

Automatic spraying systems must not be programmed to release pesticide where food or feed is directly exposed.

To Kill Houseflies, Horn Flies, Stable Flies, Horse Flies, Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, Gnats and other listed insects: Add concentrate to tank and add water to obtain desired dilution. Agitate well while adding water. Dispense diluted concentrate through automatic spraying system. Set nozzles to deliver one (1) fluid ounce per minute. Locate nozzles to cover a maximum of 2000 cubic feet of space per nozzle. Set timer to operate in accordance with equipment directions.

Not for use in outdoor residential misting systems (indoor or outdoor).

Do not apply this product in barns or stables where animals intended for slaughter or human consumption will be maintained. Do not apply when food, feed, and/or water is present.

When using this product, installers and service technicians must comply with the license, certification, or registration requirements of the state(s), tribe(s), or local authority(ies) where they are installed.

When applying via a remote activation device, do not apply when people and pets are present. If possible, when applying via automatic timer, set the timing for application when people and pets are unlikely to be present.

Direct nozzles to spray towards the target area and away from areas where people are typically present.

Do not use in an evaporative cooling system.

Do not use in misters located within 3 feet of air vents, air conditioner units, or windows.

If used in a system with a reservoir tank for the end use dilution, the system reservoir tank must be locked. Securely attach the end use pesticide label and a dilution statement to the system reservoir tank in a weather protected area or plastic sleeve. The dilution statement must be phrased as follows: this container holds ___ parts [product name] to ___ parts water.

If used in a direct injection system, the pesticide container must be locked. Securely attach the end use label to the pesticide container in a weather protected area or plastic sleeve.

This product must only be used in systems that have been calibrated to apply no more than the maximum application rate of

as one (1) fluid ounce per minute.

OUTDOOR USE: Follow dilution chart and apply as directed.

0.05% Pyrethrins – add 1 part concentrate to 18 parts water or 6.5 fluid ounces concentrate / gallon water

0.10% Pyrethrins – add 1 part concentrate to 9 parts water or 13 fluid ounces concentrate / gallon water

0.30% Pyrethrins – add 1 part concentrate to 2 parts water or 38.5 fluid ounces concentrate / gallon water

0.50% Pyrethrins – add 1 part concentrate to 1 parts water or 64 fluid ounces concentrate / gallon water

74404_1021-1799_0512 Page 8 of 18

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- (1) Treatment to soil or vegetation around structures;
- (2) Applications to lawns, turf, and other vegetation;
- (3) Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.

Do not make applications in the rain.

HOW TO APPLY

Spray outdoor surfaces of buildings, porches, garages and other areas where heads have been seen or are found. Do not spray in or near fishponds or other bodies of water.

Caution should be taken to avoid spraying when wind conditions could create a mist to blow back to applicator. Do not spray near bodies of water. Do not spray directly on plants Spray the surface areas and contact as many insects as possible.

FLIES, GNATS, MOSQUITOES, including the Cuex species that may carry and transmit the West Nile Virus, Small Flying Moths, Midges, and other flying insects: Hold contained 18-24 inches from shrubs, bushes, grassy areas and other areas infested with these pests. Do not apply directly on plants. Spray with a slow sweeping motion while moving away from the treated area. To reduce annoyance from these pests, treat open porches and patios in a similar manner. Kills and repels flies, mosquitoes, midges, gnats and when used intermittently during outdoor activities, will keep the surrounding areas free from these pests. Allow spray to dissipate before occupying the treated area.

ANTS (including Black Carpenter Ants): Spray ant trails, anthills and other areas of entry and infestation. Spray on ants where possible.

SPIDERS: Spray directly on spiders and webs.

SOWBUGS, MILLIPEDES, BEETLES, EARWIGS, Pillbugs, Centipedes, Scorpions, Cockroaches, Crickets, Waterbugs, Palmetto Bugs: Spray surfaces thoroughly and into hiding places such as cracks, crevices, moist areas, openings around pipes, spigots, storage areas, sheds and other areas where these pests tend to congregate.

FOR USE AS A PET CONCENTRATE: Dilute this concentrate to a 0.50% Pyrethrins by adding 1 part concentrate to 1 part water or 64 fluid ounces concentrate/gallon water, mixing well as water is added.

FOR USE IN LISTED PET AREAS: Remove, destroy or wash pet's old bedding. To reduce infestation, thoroughly spray the animal's bedding, sleeping quarters, kennel, around baseboards, window and door frames, wall cracks, local areas of floors, and other resting places until the surface is slightly moist. Treat these areas at the rate of two seconds per linear foot at the distance of 12 to 18 inches. Also spray the entire inside surface of doghouses. Put fresh bedding in pet's quarters after spray has dried. Allow treated areas to dry thoroughly before allowing pets to re-enter.

FLEA AND TICK INFESTATION IN RUGS AND CARPETS: Prior to use, vacuum all infested areas thoroughly and dispose of vacuum bag in outside trash. Spray rugs and carpets where fleas tend to hide, usually where animals frequent. Spray thoroughly into hiding places such as cracks and crevices. In walls and floors, along baseboards, window and door frames. Spray at the rate of one square foot per second. To kill ticks and other labeled insects, apply directly to pests. Treat pets with an EPA registered [brand name] product in conjunction with this application.

FOR KENNELS AND OTHER OUTDOOR ANIMAL AREAS to kill listed insects, apply liberally to surfaces with special attention to cracks and crevices and other insect hiding places. Allow treated areas to dry thoroughly before allowing pets [animals] to re-enter.

For best results, to protect your pet against and minimize reintroduction of adult fleas from outdoors, use EPA registered flea and tick products, ([such as] [brand name] flea or flea and tick collar, [brand name] pet powder [brand name] pet spray, [brand name] pet dip, [brand name] pet shampoo, flea shampoo or flea and tick pet shampoo), in conjunction with this application.

74404_1021-1799_0512 Page 9 of 18

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

<u>PESTICIDE STORAGE</u>: Store in a cool, dry place. Keep container closed. Always store pesticides in the original container. Store away from food and pet food.

CONTAINER DISPOSAL:

Triple rinse container promptly after emptying.

Triple rinse as follows [For containers with capacities equal to or less than a gallons]. Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the Container ¼ full with water and recap. Shake for 10 seconds. Pour insate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available or reconditioning if appropriate or puncture and dispose in a sanitary landfill, or by other procedures approved by state and local authorities.

Triple rinse as follows [For containers with capacities greater than 5 gallons]: Empty the remaining contents into application equipment or a mix tank. Fill the container of full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling if available or reconditioning if appropriate or puncture and dispose in a sanitary landfill, or by other procedures approved by state and local authorities.

74404_1021-1799_0512 Page 10 of 18

PYROCIDE® Concentrated Aqueous

Spray 74404

[Residential Section of Label]

[] or () Indicates alternate or optional wording

[Contains Pyrethrins, a botanical insecticide.]

A micro-emulsion, water-base spray

Kills Stable Flies, Horse Flies, Face Flies, Deer Flies, Horse Flies, Face Flies, Face Flies, Deer Flies, Horse Flies, Face Flie

Flies, Mosquitoes, and Gnats

Kills Black Widow Spiders and Black Carpenter Ants

Kills Fleas and Ticks around pet areas

Kills Deer Ticks and other Ixodid species that may carry and transmit Lyme Disease

Kills Mosquitoes including the Culex species that may carry and transmit the West Nile Virus

Čan be used outdoors with the Lock 'N Spray System. No mess. No Mixing. Cartridges are spill-proof.

ACTIVE INGREDIENTS:

 Pyrethrins
 1.00%

 Piperoryl butoxide
 10.00%

 SREDIENTS
 89.00%

* (butylcarbityl) (6-propylpiperonyl) ether and related compounds

PYROCIDE® is a registered trademark of McLaughlin Gormley King Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice.
	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to by a poison control center or doctor.
	Do not give anything by mouth to an unconscious person.
IF ON SKIN:	Take off contaminated clothing.
	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For	
information regarding medical emergencies or pesticide incidents, call 1-888-740-8712	

Manufactured by



8810 Tenth Avenue North Minneapolis, MN 55427

EPA Reg. No. 1021-1799

Net Contents _____

EPA Est. No. 1021-MN-2

74404_1021-1799_0512 Page 11 of 18

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. Drittand runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product may contaminate water through runoff. This product has a potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

tables are more prone to produce runoff that contains this product.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide of the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use this product in or on electrical equipment because of shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- . Do not apply this product in a way that will contact other persons, either directly or through drift.
- Remove or cover exposed food and drinking water before application.
- Do not make applications during rain
- Do not water the treated area to the point of run-off
- Do not wet articles to point of runoff or drip.
- Do not use treated article until spray has dried.
- Do not apply directly to sewers or drains, or to any area like a gutter where drainage to sewers storm drains, water bodies, or aquatic habitat can occur.

OUTDOOR USE: Follow dilution chart and apply as directed.

0.05% Pyrethrins – add 1 part concentrate to 18 parts water or 6.5 fluid ounces concentrate / gallon water 0.10% Pyrethrins – add 1 part concentrate to 9 parts water or 13 fluid ounces concentrate / gallon water

0.30% Pyrethrins – add 1 part concentrate to 2 parts water or 38.5 fluid ounces concentrate / gallon water

0.50% Pyrethrins – add 1 part concentrate to 1 parts water or 64 fluid ounces concentrate / gallon water

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- (1) Treatment to soil or vegetation around structures;
- (2) Applications to lawns, turf, and other vegetation;
- (3) Applications to building foundations, up to a maximum height of 3 feet.

74404_1021-1799_0512 Page 12 of 18

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.

FOR USE WITH THE LOCK 'N SPRAY SYSTEM - Kills Lawn and Garden and nuisange pests.

	CONNECT	Connect sprayer to hose
	ATTACH	Attach cartridge to sprayer. Turn cartridge until it clicks into place.
	SPRAY	To begin spraying, rotate sprayer nozzle tip to direct spray up or down, or remove for a 20 to 25 foot jet spray. Turn on water and squeeze trigger.
	FINISH	To stop spraying, release trigger, unlock cartridge and remove from sprayer. (No need to turn of water)
		To clean sprayer, flush with water, squeezing and releasing trigger in 2 short bursts. Turn off water.
5		sprayer, then disconnect sprayer from hose.
<u> </u>	Carringe reseals is	self automatically – No spills, no leaks!

HOW TO APPLY

Spray outdoor surfaces of buildings, porches, garages and other areas where insects have been seen or are found. Do not spray in or near fishponds or other bodies of water.

Caution should be taken to avoid spraying when wind conditions could create a mist to blow back to applicator. Do not spray near bodies of water. Do not spray directly on plants. Spray the surface areas and contact as many insects as possible.

FLIES, GNATS, MOSQUITOES, including the Culex species that may carry and transmit the West Nile Virus, Small Flying Moths, Midges, and other flying insects: Hold container 18-24 inches from shrubs, bushes, grassy areas and other areas infested with these pests. Do not apply directly on plants. Spray with a slow sweeping motion while moving away from the treated area. To reduce annoyance from these pests, treat open porches and patios in a similar manner. Kills and repels flies, mosquitoes, midges, gnats and when used intermittently during outdoor activities, will keep the surrounding areas free from these pests. Allow spray to dissipate before occupying the treated area.

ANTS (including Black Carpenter Ants): Spray ant trails, anthills and other areas of entry and infestation. Spray on ants where possible.

SPIDERS: Spray directly on spiders and webs.

SOWBUGS, MILLIPEDES, BEETLES, EARWIGS, Pillbugs, Centipedes, Scorpions, Cockroaches, Crickets, Waterbugs, Palmetto Bugs: Spray surfaces thoroughly and into hiding places such as cracks, crevices, moist areas, openings around pipes, spigots, storage areas, sheds and other areas where these pests tend to congregate.

FOR USE AS A PET CONCENTRATE: Dilute this concentrate to a 0.50% Pyrethrins by adding 1 part concentrate to 1 part water or 64 fluid ounces concentrate/gallon water, mixing well as water is added.

FOR USE IN LISTED PET AREAS: Remove, destroy or wash pet's old bedding. To reduce infestation, thoroughly spray the animal's bedding, sleeping quarters, kennel, around baseboards, window and door frames, wall cracks, local areas of floors, and other resting places until the surface is slightly moist. Treat these areas at the rate of two seconds per linear foot at the distance of 12 to 18 inches. Also spray the entire inside surface of doghouses. Put fresh bedding in pet's quarters after spray has dried. Allow treated areas to dry thoroughly before allowing pets to re-enter.

FLEA AND TICK INFESTATION IN RUGS AND CARPETS: Prior to use, vacuum all infested areas thoroughly and dispose of vacuum bag in outside trash. Spray rugs and carpets where fleas tend to hide, usually where animals frequent. Spray thoroughly into hiding places such as cracks and crevices. In walls and floors, along baseboards, window and door frames.

74404_1021-1799_0512 Page 13 of 18

Spray at the rate of one square foot per second. To kill ticks and other labeled insects, apply directly to pests. Treat pets with an EPA registered [brand name] product in conjunction with this application.

FOR KENNELS AND OTHER OUTDOOR ANIMAL AREAS to kill listed insects, apply liberally to surfaces with special attention to cracks and crevices and other insect hiding places. Allow treated areas to dry thoroughly before allowing pets [animals] to reenter.

For best results, to protect your pet against and minimize reintroduction of adult items from outdoors, use EPA registered flea and tick products, ([such as] [brand name] flea or flea and tick collar, [brand name] pet powder [brand name] pet spray, [brand name] pet dip, [brand name] pet shampoo, flea shampoo or flea and tick pet shampoo, in oprigination with this application.

STORAGE AND DISPOSĂI

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE:

Store in a cool, dry place. Keep container closed. Aways store pesticides in the original container. Store away from food and pet food.

CONTAINER DISPOSAL:

If empty: Non refillable container. Op not refill this container. Offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

74404_1021-1799_0512 Page 14 of 18

74404-0512

(Consumer label, Outdoor only)

(Front Panel)

PYROCIDE® Concentrated Aqueous Spray 74404

An outdoor mosquito killing product for use in outdoor areas including - backwards, patios decks, campsites and picnic areas.

Contains Pyrethrins, a botanical insecticide.

A micro-emulsion, water-base spray

Water-based, Non-staining, No oily residue

Kills Mosquitoes, including those which may transmit West Vile Virus

Kills Mosquitoes, including Culex species that may carry and transmit the West Nile Virus

Quick Knockdown

Kills Mosquitoes, Gnats, and Biting Flies

Kills on Contact

8 fluid ounces treats up to 5000 square feet

Easy to Use

Lock & Load Cartridge

No Mixing, No Handling, No Mes

Protects your yard from biting adult mosquitoes

Convenient Lock & Load Cartridge

Low Odor

Spray [Pyrocide Concentrated Aqueous Spray 74404] before your event to kill flies, mosquitoes and gnats

Do-It-Yourself Mosquito Killing

For use in compatible [optional equipment brand name] handheld mechanical foggers [sprayers]

For Outdoor Use Only

Optional Marketing Text on Package:

[Mosquitoes are not only a nuisance, they can transmit diseases that are a serious health threat to people and pets. With just one bite, mosquitoes can transmit West Nile Virus and other viruses that cause encephalitis. They can also transmit heartworm disease to dogs and cats.]

ACTIVE INGREDIENTS:

Pyrethrins	1.00%
Piperonyl butoxide, Technical *	10.00%
OTHER INGREDIENTS	89.00%
-	100.00%

* Equivalent to min. 8.0% (butylcarbityl)(6-propylpiperonyl) ether and 2.0% related compounds PYROCIDE® - Registered trademark of McLaughlin Gormley King Company

EPA Reg. No. 1021-1799 EPA Est. No.: 1021-MN-2

KEEP OUT OF REACH OF CHILDREN CAUTION

See Side Panel for additional Precautionary Statements

Net Contents: 8 fl.oz.

74404_1021-1799_0512 Page 15 of 18

PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS AND DOMESTIC ANIMALS – CAUTION. Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove and wash contaminated clothing before reuse.

	FIRST AID
If Swallowed:	 Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on Skin or Clothing:	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center of the treatment advice.
doctor, or going	t container or label with you when calling a poison control center or for treatment. For information regarding medical emergencies or ts, call the International Poison Center at 1-888-740-8712.

ENVIRONMENTAL HAZARDS—This product is toxic to fish and aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product may contaminate water through runoff. This product has a potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated are will help avoid run off to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS - Do not use this product in or on electrical equipment because of shock hazard.

[Pyrocide® Concentrated Aqueous Spray 74404] contains the botanical insecticide, Pyrethrins, which provides rapid effective kill of insects upon direct contact with an added benefit of flushing these insects to aid in a more complete kill.

[Pyrocide® Concentrated Aqueous Spray 74404] can be used to reduce annoyance of mosquitoes, including the Culex species that may carry and transmit the West Nile Virus, outdoors around homes, recreational and other listed areas. Ideal for use prior to outdoor activities.

[Pyrocide® Concentrated Aqueous Spray 74404] provides rapid kill of insects without objectionable residues.

OUTDOOR AREAS for use in and around, including:

- Amphitheaters
- Back Yards
- Balconies
- Baseball Fields
- Campgrounds
- Corrals
- Courtyards
- Front Yards
- Gazebos
- Kennels

- Marinas
- Parks
- Patios/Decks
- Playgrounds
- Recreational Areas

74404_1021-1799_0512 Page 16 of 18

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read all directions completely before use. For best results, follow directions for specific use areas.

GENERAL DIRECTIONS

DO NOT open cartridge. This product's packaging is designed for easy lock & foad use in [optional equipment brand name] handheld mechanical foggers [sprayers]. Peel off protective strip to expose cartridge in. Do not puncture tip. Follow equipment manufacturer's directions for loading the cartridge into the spray unit.

USE RESTRICTIONS:

- Do not apply this product in a way that will contact other persons, either directly or through drift.
- Remove or cover exposed food and drinking water before application.
- Do not make applications during rain
- Do not water the treated area to the point of run-off
- Do not wet articles to point of runoff or drip.
- Do not use treated article until spray has dried
- Do not apply directly to sewers or drains, or to any area like a gutter where drainage to sewers storm drains, water bodies, or aquatic habitat can occur.


All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

- (1) Treatment to soil or vegetation around structures;
- (2) Applications to lawns, turk and other vegetation;
- (3) Applications to building foundations up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.

To kill Mosquitoes, Gnats and Biting Flies: Caution should be taken to avoid spraying when wind conditions could create a mist to blow back to the applicator. Do not apply this product in or around bodies of water. When spraying plants, do not operate closer than 18 inches. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90 F. For best results, spray when mosquitoes are most active – application during the early evening or early morning hours is usually preferable.

Applications of [Pyrocide Aqueous Spray 74404] using a handheld mechanical fogger [sprayer] provide quick knockdown of adult mosquitoes that are present in the treatment area during application. Applications target open outdoor and mosquito harborage areas and provide quick relief from biting mosquitoes.



74404_1021-1799_0512 Page 17 of 18

USE DIRECTIONS: Mechanical sprayer [optional equipment brand name] has a fixed rate of 1.2 to 1.6 fl.oz./min. No adjustment is necessary. One cartridge treats up to 5,000 sq. ft. Begin spraying at furthest down-wind edge of the treatment area. While walking at a slow, comfortable pace, spray perpendicular to the wind pointing the nozzle downwind, until reaching the opposite edge of treatment area. Continue this motion upwind every 10 feet until entire treatment area has been sprayed (see treatment diagram).

Pay special attention to standing weeds, bushy non-crop areas, grasses and other areas of dense vegetation where mosquitoes may harbor. Use a sweeping motion.

To reduce annoyance from these pests, treat open porches and patios in a similar manner. Allow spray to dissipate before occupying the treated area.

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place. Keep cartridge sealed. Always Store pesticides in the original container.

Store away from food and pet food

Container Disposal: If empty: Non refillable container. Do not reuse or refill this container. Offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured by



8810 Tenth Avenue North Minneapolis, MN 55427

74404_1021-1799_0512 Page 18 of 18