

Medication Guide

BUTRANS® (BYOO-trans)

(buprenorphine) Transdermal System, CIII

BUTRANS is:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require daily, around-the-clock, long-term treatment with an opioid, when other pain treatments such as non-opioid pain medicines or immediate-release opioid medicines do not treat your pain well enough or you cannot tolerate them.
- A long-acting (extended-release) opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to death.
- Not for use to treat pain that is not around-the-clock.

Important information about BUTRANS:

- **Get emergency help right away if you take too much BUTRANS (overdose).** When you first start taking BUTRANS, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur.
- Never give anyone else your BUTRANS. They could die from taking it. Store BUTRANS away from children and in a safe place to prevent stealing or abuse. Selling or giving away BUTRANS is against the law.

Do not use BUTRANS if you have:

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.

Before applying BUTRANS, tell your healthcare provider if you have a history of:

- head injury, seizures
- liver, kidney, thyroid problems
- problems urinating
- heart rhythm problems (Long QT syndrome)
- pancreas or gallbladder problems
- abuse of street or prescription drugs, alcohol addiction, or mental health problems.

Tell your healthcare provider if you:

- have a fever
- **are pregnant or planning to become pregnant.** Prolonged use of BUTRANS during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **are breastfeeding.** BUTRANS passes into breast milk and may harm your baby.
- are taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking BUTRANS with certain other medicines can cause serious side effects.

When using BUTRANS:

- Do not change your dose. Apply BUTRANS exactly as prescribed by your healthcare provider.
- See the detailed Instructions for Use for information about how to apply the BUTRANS patch.
- Do not apply a BUTRANS patch if the pouch seal is broken, or the patch is cut, damaged, or changed in any way.
- Do not apply more than 1 patch at the same time unless your healthcare provider tells you to.
- You should wear 1 BUTRANS patch continuously for 7 days.
- **Call your healthcare provider if the dose you are using does not control your pain.**
- **Do not stop using BUTRANS without talking to your healthcare provider.**
- **To properly dispose of used and unused patches, use the Patch-Disposal Unit or fold in half and flush down the toilet. See the detailed Instructions for Use.**

While using BUTRANS DO NOT:

- Take hot baths or sunbathe, use hot tubs, saunas, heating pads, electric blankets, heated waterbeds, or tanning lamps. These can cause an overdose that can lead to death.
- Drive or operate heavy machinery, until you know how BUTRANS affects you. BUTRANS can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines containing alcohol. Using products containing alcohol during treatment with BUTRANS may cause you to overdose and die.

The possible side effects of BUTRANS are:

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, itching, redness or rash where the patch is applied. Call your healthcare provider if you have any of these symptoms and they are severe.

Get emergency medical help if you have:

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue or throat, extreme drowsiness, light-headedness when changing positions, or you are feeling faint.

These are not all the possible side effects of BUTRANS. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. **For more information go to dailymed.nlm.nih.gov**

Distributed by: Purdue Pharma L.P., Stamford, CT 06901-3431, www.purduepharma.com or call 1-888-726-7535

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised: April 2014

Reference ID: 3612149

Butrans®
(buprenorphine) Transdermal System