

## Antenatal Vaccination Card Influenza and Whooping Cough (Pertussis)

The Antenatal Vaccination Card has been provided for you to maintain an accurate vaccination record for the whooping cough (pertussis) and influenza (flu) vaccines received during your pregnancy.

Please keep the completed record card with your antenatal documents and take it to all your appointments.

Receiving both vaccinations is recommended for pregnant women. However for greatest protection against flu it is recommended that you receive the flu vaccination as early as possible in pregnancy. This will also depend the time of year and vaccine availability.

It is recommended the whooping cough (pertussis) vaccine is given at 28 weeks pregnant or as soon as possible afterwards.

Reactions to the vaccine can be localised pain, redness and swelling at injection site and fever. Any serious or unexpected reactions can be reported to Health Protection Service, Immunisation Unit on 62052300

To be completed by person administering vaccine				
Full Name:	D.0	О.В:	//	/
Do you identify as being: Aboriginal 🗖 Torres	Strait Islander 🗖	Both 🗖	Neither	· 🗖
Estimated date of delivery:///				
Vaccine(s) administered today:///////_				
Influenza Vaccine (recommended for women at any stage of pregnancy)				
Vaccine Brand: 🗖 Vaxigrip 🛛 Fluarix 🛛	🛛 Fluvax 🗖 🖸	Other:		
Batch Number:	_			
Clinic or Surgery or Name: Signature of person administering vaccine(s): Name of person administering: Date://				
Vaccine(s) administered today:///////_				
Pertussis Vaccine (recommended for women delivery)	n at 28 weeks or A	SAP afterv	vards. Ca	an be given until
Vaccine Brand: 🗖 Adacel® 🛛 🗖 Boostrix® 🕻	<b>]</b> Other:			
Batch Number:	_			
Clinic or Surgery or Name: Signature of person administering vaccine(s): Name of person administering: Date://				