#4) A material will float on the surface of a liquid if the material has a density less than that of the liquid. Given that the density of water is approximately 1.00 g/ mL, will a block of material having a value of 1.2  $\times$  10<sup>4</sup> in<sup>3</sup> weighing 350 lb. float or sink when placed in a reservoir of water? (1 inch = 2.54 cm)

#5) A rectangular block has dimensions  $2.9 \text{ cm} \times 3.5 \text{ cm} \times 10.0 \text{ cm}$ . The mass of the block is 615.0 g. What are the volume and density of the block?

#6) Diamonds are measured in carats, and 1 carat = $0.200  \text{g}$ . The density of diamond is $3.51  \text{g/cm}^3$ . What are the volume and density of the block? The sample is $4.5  \text{carats}$ .
#7) The density of pure silver is 10.5 g/cm³ at 20°C. If 5.25 g of pure silver pellets is added to a graduated cylinder containing 11.2 mL of water, to what volume level will the water in the cylinder rise?
#8) The density of osmium (the densest metal) is 22.57 g/cm <sup>3</sup> . If a 1.00 kg rectangular block of osmium has two dimensions of 4.00 cm x 4.00 cm, calculate the third dimension of the block.
#9) In the opening scene of the movie <i>Raiders of the Lost Ark</i> , the hero Indiana Jones tries to remove a gold idol from a booby-trapped pedestal. He replaces the idol with a bag of sand of approximately equal volume. (Density of gold = $19.3 \text{g/cm}^3$ ; density of sand $\approx 2.00 \text{ g/cm}^3$ .)  a) Did he have a reasonable chance of not activating the mass sensitive booby trap?
b) In a later scene he and an unscrupulous guide play catch with the idol. Assume that the volume of the idol is about 1.0 L. If it were solid gold, what mass would the idol have? Is playing catch with it plausible?