

Igor V. Skrypnik; Dmitry V. Larin

On weighted estimates of solutions of nonlinear elliptic problems

*Mathematica Bohemica*, Vol. 124 (1999), No. 2-3, 173–184

Persistent URL: <http://dml.cz/dmlcz/126242>

## Terms of use:

© Institute of Mathematics AS CR, 1999

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON WEIGHTED ESTIMATES OF SOLUTIONS OF  
NONLINEAR ELLIPTIC PROBLEMS

IGOR V. SKRYPNIK, DMITRY V. LARIN, Donetsk

(Received November 24, 1998)

*Dedicated to Professor Alois Kufner on the occasion of his 65th birthday*

*Abstract.* The paper is devoted to the estimate

$$|u(x, k)| \leq K |k| \left\{ \operatorname{cap}_{p,w}(F) \frac{\varrho^p}{w(B(x, \varrho))} \right\}^{\frac{1}{p-1}},$$

$2 \leq p < n$  for a solution of a degenerate nonlinear elliptic equation in a domain  $B(x_0, 1) \setminus F$ ,  $F \subset B(x_0, d) = \{x \in \mathbb{R}^n : |x_0 - x| < d\}$ ,  $d < \frac{1}{2}$ , under the boundary-value conditions  $u(x, k) = k$  for  $x \in \partial F$ ,  $u(x, k) = 0$  for  $x \in \partial B(x_0, 1)$  and where  $0 < \varrho \leq \operatorname{dist}(x, F)$ ,  $w(x)$  is a weighted function from some Muckenhoupt class, and  $\operatorname{cap}_{p,w}(F)$ ,  $w(B(x, \varrho))$  are weighted capacity and measure of the corresponding sets.

*Keywords:* degeneracy, Muckenhoupt class, pointwise estimate, nonlinear elliptic equation, capacity, a-priori estimate

*MSC 1991:* 35J70, 35B45

In the study of behaviour of solutions of nonlinear elliptic and parabolic equations an important role is played by special estimates of model problems in domains with small holes (see [1, 2]). In many cases this role is analogous to that of estimates of singular solutions of linear equations. By using these estimates the following problems were studied: asymptotical behaviour and the construction of correctors for nonlinear elliptic and parabolic problems in perforated domains, a necessary condition for the regularity of boundary points, the stability of solutions of nonlinear problems with respect to the variation of domains. The proof and applications of these estimates for elliptic equations are given in [1]. This paper is devoted to the

extension of the method of obtaining of pointwise estimates for degenerate nonlinear elliptic equations.

#### 1. AUXILIARY LEMMAS AND STATEMENT OF THE RESULT

Let  $w$  be a locally integrable nonnegative function in  $\mathbb{R}^n$  and assume that  $0 < w(x) < \infty$  almost everywhere. We say that  $w$  belongs to the Muckenhoupt class  $A_t$ ,  $1 < t < \infty$ , if there exists a constant  $c_{t,w}$  such that

$$(1) \quad \frac{1}{|B|} \int_B w \, dx \leq c_{t,w} \left( \frac{1}{|B|} \int_B w^{1-t} \, dx \right)^{1-t}$$

for all balls  $B$  in  $\mathbb{R}^n$ . By  $|E|$  we denote the Lebesgue  $n$ -measure of a measurable set  $E \subset \mathbb{R}^n$ .

We shall note only certain properties of functions from Muckenhoupt class.

**Lemma 1.** *If  $w \in A_t$ , then*

$$(2) \quad \left( \frac{|E|}{|B|} \right)^t \leq c_{t,w} \frac{w(E)}{w(B)},$$

where  $B$  is an arbitrary ball in  $\mathbb{R}^n$ ,  $E$  is a measurable subset of  $B$  and

$$w(E) = \int_E w(x) \, dx.$$

**Lemma 2.** *If  $w \in A_t$ ,  $t > 1$ , then  $w \in A_{t-\varepsilon}$  for some  $\varepsilon$ ,  $0 < \varepsilon < t - 1$ . Moreover,  $\varepsilon$  and  $c_{t-\varepsilon,w}$  depend only on  $n$ ,  $t$ ,  $c_{t,w}$ .*

For the proofs of Lemmas 1, 2 see [3], Chapter 15.

**Lemma 3.** *Suppose  $w \in A_t$  and  $s > t$ . Then  $w \in A_s$ .*

This statement immediately follows from the Hölder inequality and (1).

**Lemma 4. ( $A_t$ -weighted Poincaré inequality)** *Suppose  $w \in A_t$  and let for arbitrary  $x, s, h, 0 < s \leq h$  an inequality*

$$\frac{s}{h} \left( \frac{w(B(x,s))}{w(B(x,h))} \right)^{\frac{1}{q}} \leq c \left( \frac{w(B(x,s))}{w(B(x,h))} \right)^{\frac{1}{t}}, \quad q > t,$$

hold with a constant  $c$  independent of  $x, s, h$ . Then

$$\left( \frac{1}{w(B)} \int_B |v(x) - \frac{1}{|B|} \int_B v(x) dx|^q w(x) dx \right)^{\frac{1}{q}} \leq C r \left( \frac{1}{w(B)} \int_B \left| \frac{\partial v(x)}{\partial x} \right|^t w(x) dx \right)^{\frac{1}{t}},$$

where  $B = B(x_0, r)$ ,  $v(x) \in C^\infty(B)$  and  $C$  is independent of  $x_0, r, v$ .

**Lemma 5.** ( $A_t$ -weighted Sobolev inequality) With the same hypotheses as in Lemma 4 we have

$$\left( \frac{1}{w(B)} \int_B |v(x)|^q w(x) dx \right)^{\frac{1}{q}} \leq C r \left( \frac{1}{w(B)} \int_B \left| \frac{\partial v(x)}{\partial x} \right|^t w(x) dx \right)^{\frac{1}{t}},$$

where  $B = B(x_0, r)$ ,  $v(x) \in C_0^\infty(B)$  and  $C$  is independent of  $x_0, r, v$ .

For the proofs of Lemmas 4, 5 see [4].

Definition and basic properties of the Muckenhoupt class  $A_t$  were explicitly studied in [3].

Let  $F$  be an arbitrary compact set in  $\mathbb{R}^n$ . Let us denote by  $d$  the minimum of the radii of balls containing  $F$ , and let  $x_0$  be the center of such a ball with radius  $d$ , satisfying  $F \subset \overline{B}(x_0, d)$ . Here and in the sequel  $B(x, r)$  denotes the ball with radius  $r$  and center at  $x$ .

Let  $\psi(x)$  be a function from the class  $C_0^\infty(B(x_0, 1))$ , equal to one in  $B(x_0, \frac{1}{2})$ . If  $d < \frac{1}{2}$ , then for an arbitrary real  $k$  we consider a nonlinear elliptic boundary value problem

$$(3) \quad \sum_{i=1}^n \frac{d}{dx_i} a_i(x, \frac{\partial u}{\partial x}) = 0, \quad x \in D,$$

$$(4) \quad u(x) = k\psi(x), \quad x \in \partial D.$$

Here  $D = B(x_0, 1) \setminus F$ .

We assume that the functions  $a_i(x, g)$ ,  $i = 1, \dots, n$ , are defined for  $x \in \overline{B}$  (here and in the sequel  $B = B(x_0, 1)$ ) and  $g \in \mathbb{R}^n$ , and satisfy the following conditions:

A<sub>1</sub>) functions  $a_i(x, g)$  are continuous in  $g$  for almost every  $x \in \overline{B}$ , measurable in  $x$  for all  $g \in \mathbb{R}^n$ ;

$A_2$ ) there are positive constants  $\nu_1, \nu_2$  such that for  $2 \leq p < n$  and  $x \in \overline{B}$ ,  $g, q \in \mathbb{R}^n$  the inequalities

$$\begin{aligned} |a_i(x, g)| &\leq \nu_1 |g|^{p-1} w(x), \\ \sum_{i=1}^n [a_i(x, g) - a_i(x, q)](g_i - q_i) &\geq 0, \\ \sum_{i=1}^n a_i(x, g) g_i &\geq \nu_2 |g|^p w(x) \end{aligned}$$

hold, where  $w(x) \in A_{(p-1)+\frac{1}{n}}(\mathbb{R}^n)$ ,  $[w(x)]^{-\frac{1}{p-1}} \in A_{\frac{p-1}{p-1}(1-\frac{1}{n})}(\mathbb{R}^n)$ .

**Remark 1.** Let us choose  $w(x) = |x|^\alpha$ ,  $-n + p < \alpha < n(p-1) - (n-p)$  and  $p \geq 2$ . In this case  $w(x)$  satisfies  $A_2$ . This is easily verified by a direct computation.

A solution of the boundary value problem (1), (2) is a function  $u(x, k) \in W_p^1(D, w)$  such that  $u(x, k) - k\psi(x) \in W_p^{\circ 1}(D, w)$  and the integral identity

$$(5) \quad \sum_{i=1}^n \int_D u_i \left( x, \frac{\partial u}{\partial x} \right) \frac{\partial \varphi}{\partial x_i} dx = 0$$

holds for arbitrary function  $\varphi(x) \in \overset{\circ}{W}_p^1(D, w)$ .

The definitions and properties of weighted Sobolev spaces  $W_p^1(\Omega, w)$ ,  $W_p^{\circ 1}(\Omega, w)$  were studied in [3, 4, 5] (here  $\Omega \subset \mathbb{R}^n$ ).

The existence and uniqueness of the function  $u(x, k)$  follows from the global theory of monotone operators (for example, see [1]). The function  $u(x, k)$  is assumed to be extended to  $F$  by the constant  $k$ .

For the purpose of formulation of our main result let us introduce the notion of weighted  $(p, w)$ -capacity  $\text{cap}_{p,w}$  (see [3]).

The number

$$(6) \quad \text{cap}_{p,w}(E) = \inf \int_B \left| \frac{\partial v(x)}{\partial x} \right|^p w(x) dx$$

is called the  $(p, w)$ -capacity of the closed set  $E \subset B(x_0, \frac{1}{2})$ . The infimum in (6) is taken over all functions  $v(x) \in C_0^\infty(B)$  satisfying the equality  $v(x) = 1$  for  $x \in E$ .

Further, we shall prove the following

**Theorem.** *Let us assume that conditions  $A_1, A_2$ , are satisfied. Then there exists a constant  $K$  depending only on  $n, p, \nu_1, \nu_2$  and the Muckenhoupt constant  $c_{p,w}$  of*

$w$  such that, for a solution  $u(x, k)$  of the problem (1), (2) and for an arbitrary point  $x \in D$ ,

$$(7) \quad |u(x, k)| \leq K|k| \left\{ \text{cap}_{p,w}(F) \frac{\varrho^p}{w(B(x, \varrho))} \right\}^{\frac{1}{p-1}},$$

where  $0 < \varrho \leq \varrho(x, F)$ .

**Remark 2.** In case  $w(x) \equiv 1$  the estimate (7) coincides with the pointwise estimate of the solution of nonlinear Dirichlet problem obtained by the first author in [2]. Exactness of (7) follows also from

$$G(x, \xi) \approx \frac{|x - \xi|^2}{w(B(x, |x - \xi|))}$$

for the fundamental solution  $G(x, \xi)$  of the operator

$$L = \sum_{i,j=1}^n D_{x_i} (a_{ij}(x) D_{x_j}),$$

where  $a_{i,j}(x)$  are real-valued, symmetric and

$$\lambda w(x) |\xi|^2 \leq \sum_{i,j=1}^n a_{ij}(x) \xi_i \xi_j \leq \frac{1}{\lambda} w(x) |\xi|^2,$$

whenever  $\lambda > 0$ ,  $\xi = (\xi_1, \dots, \xi_n)$ , and  $w(x)$  is the same as in condition A<sub>2</sub> (case  $p = 2$ ). This estimate was obtained in [6].

## 2. PROOF OF THE MAIN RESULT

Let us assume  $k > 0$ .

**Lemma 6.** Let us assume that conditions A<sub>1</sub>, A<sub>2</sub> are satisfied and let  $u(x, k)$  be the solution of the problem (1), (2). Then for  $k \neq 0$

$$0 \leq \frac{1}{k} u(x, k) \leq 1.$$

**P r o o f.** Let us take the test-function  $\varphi_1(x) = \min\{u(x, k), 0\}$  in the integral identity (5) and use the condition A<sub>2</sub>. We obtain

$$\int_{D_1} \left| \frac{\partial u(x, k)}{\partial x} \right|^p w(x) dx \leq 0,$$

where  $D_1 = \{x \in D : u(x, k) < 0\}$ . From this inequality it follows that  $u(x, k) \geq 0$ . Similarly, replacing  $\varphi(x)$  in (5) by  $\varphi_2(x) = \max\{u(x, k) - k, 0\}$  the inequality  $u(x, k) \leq k$  is established.  $\square$

**Lemma 7.** Assume that conditions A<sub>1</sub>, A<sub>2</sub> are satisfied. Then there exists a constant c<sub>1</sub>, depending only on n, p, ν<sub>1</sub>, ν<sub>2</sub>, c<sub>p,w</sub>, such that

$$(8) \quad \int_D \left| \frac{\partial u(x, k)}{\partial x} \right|^p w(x) dx \leq c_1 k^p \operatorname{cap}_{p,w}(F).$$

**P r o o f.** Let us take the test-function  $\varphi = u(x, k) - k\psi(x)$  in the integral identity (5), where  $\psi(x)$  is from the class  $C_0^\infty(B)$ , and  $\psi$  is equal to one in F. Using the condition A<sub>2</sub> and Young's inequality we estimate the terms of the obtained equality and get

$$(9) \quad \int_D \left| \frac{\partial u(x, k)}{\partial x} \right|^p w(x) dx \leq c_2 k^p \int_B \left| \frac{\partial \psi(x)}{\partial x} \right|^p w(x) dx.$$

Here and in the sequel we denote by  $c_i$  constants depending only on the same parameters as the constant K in the formulation of the Theorem.

By virtue of definition (6), inequality (9) proves the estimate (8).  $\square$

Let us denote for  $0 < \mu < k$

$$E_\mu = \{x \in D : 0 \leq u(x, k) \leq \mu\}.$$

**Lemma 8.** Let us assume that conditions A<sub>1</sub>, A<sub>2</sub> are satisfied. Then there exists a constant c<sub>3</sub> such that

$$(10) \quad \int_{E_\mu} \left| \frac{\partial u(x, k)}{\partial x} \right|^p w(x) dx \leq c_3 \mu k^{p-1} \operatorname{cap}_{p,w}(F).$$

**P r o o f.** We substitute  $\varphi(x) = u_\mu(x, k) - \frac{\mu}{k} u(x, k)$  in (5), where  $u_\mu(x, k) = \min\{u(x, k), \mu\}$ . By standard computations and (8) we obtain (10).  $\square$

In order to prove Theorem we need some auxiliary results.

**Lemma 9.** Let  $2 \leq p < n$  and  $w \in A_{(p-1)+\frac{1}{n}}$ ,  $[w]^{-\frac{1}{p-1}} \in A_{\frac{p}{p-1}(1-\frac{1}{n})}$ . For any function  $v(x) \in W_p^{2,1}(B(0, R), w)$  and any numbers r, R, satisfying the conditions  $0 < r \leq R$  the inequality

$$(11) \quad \int_{B(0, r)} |v(x)|^p w(x) dx \leq K_1 r^p \int_{B(0, R)} \left| \frac{\partial v(x)}{\partial x} \right|^p w(x) dx$$

holds with a constant  $K_1$  depending only on  $n, p, c_{p,w}$ .

**P r o o f.** Without loss of generality we may assume that  $v(x) \in C_0^\infty(B(0, R))$ . From  $A_p$ -weighted Poincaré inequality we have

$$\begin{aligned}
& \frac{1}{w(B(0, r))} \int_{B(0, r)} |v(x)|^p w(x) dx \\
& \leq \frac{2^{p-1}}{w(B(0, r))} \int_{B(0, r)} \left| v(x) - \frac{1}{|B(0, r)|} \int_{B(0, r)} v(y) dy \right|^p w(x) dx \\
& \quad + 2^{p-1} \left( \frac{1}{|B(0, r)|} \int_{B(0, r)} |v(y)| dy \right)^p \\
& \leq c_4 \frac{r^p}{w(B(0, r))} \int_{B(0, r)} \left| \frac{\partial v(x)}{\partial x} \right|^p w(x) dx + c_4 \left( \frac{1}{|B(0, r)|} \int_{B(0, r)} |v(y)| dy \right)^p.
\end{aligned} \tag{12}$$

Now we only need to estimate the last term on the right-hand side of (12).

Let  $\omega = \frac{x}{|x|}$ . A straightforward calculation yields

$$|v(x)| = \left| \int_{|x|}^R \frac{d}{dt} v(\omega t) dt \right| \leq \left| \int_{|x|}^R \left| \frac{\partial v}{\partial x}(\omega t) \right| dt \right|. \tag{13}$$

Transforming the last integral on the right-hand side of (12) into spherical coordinates with respect to the variables  $|x| \in [0, r]$ ,  $\omega = \frac{x}{|x|} \in S_1(0)$ , using (13) and Hölder inequality, we obtain

$$\begin{aligned}
I_1 &= \left\{ \frac{1}{|B(0, r)|} \int_{B(0, r)} |v(y)| dy \right\}^p = \left\{ \frac{1}{|B(0, r)|} \int_0^r \int_{S_1(0)} |v(|x|\omega)| |x|^{n-1} d\omega dx \right\}^p \\
&\leq \left\{ \frac{1}{|B(0, r)|} \int_0^r \int_{S_1(0)} \int_{|x|}^R \left| \frac{\partial v}{\partial x}(\omega t) \right| dt |x|^{n-1} d\omega dx \right\}^p \\
&\leq c_5 \left\{ \frac{1}{|B(0, r)|^p} \int_0^r \int_{|x|=S_1(0)} \int_{|x|}^R \left| \frac{\partial v}{\partial x}(\omega t) \right|^p w(\omega t) t^{n-1} d\omega dt |x|^{n-1} dx \right\}^p \\
&\quad \times \left\{ \int_0^r \int_{|x|=S_1(0)} \int_{|x|}^R [w(\omega t)]^{-\frac{1}{p-1}} t^{-\frac{n-1}{p-1}} d\omega dt |x|^{n-1} dx \right\}^{p-1}
\end{aligned} \tag{14}$$

$$\leq c_6 \left\{ \frac{1}{r^{n(p-1)}} \int_{B(0,R)} \left| \frac{\partial v}{\partial x}(x) \right|^p w(x) dx \right\} \\ \times \left\{ \int_0^r |x|^{n-1} \int_{|\omega| \leq S_1(0)}^R \int_{\mathbb{R}} [w(\omega t)]^{-\frac{1}{p-1}} t^{-\frac{n-1}{p-1}} d\omega dt d|x| \right\}^{p-1}.$$

Now we estimate separately the integral

$$(15) \quad I_2 = \int_{|z| \leq S_1(0)}^R \int_{\mathbb{R}} [w(\omega t)]^{-\frac{1}{p-1}} t^{-\frac{n-1}{p-1}} d\omega dt = \int_{|z| \leq |x| \leq R} |z|^{-\frac{n(p+1)}{p-1}} [w(z)]^{-\frac{1}{p-1}} dz \\ \leq \sum_{j=1}^{\infty} \int_{|x| \leq |z| \leq 2^j |x|} |z|^{-\frac{n(p+1)}{p-1}} [w(z)]^{-\frac{1}{p-1}} dz \\ \leq c_7 \sum_{j=1}^{\infty} (2^j |x|)^{-\frac{n(p+1)}{p-1}} \int_{|z| \leq 2^j |x|} [w(z)]^{-\frac{1}{p-1}} dz.$$

Since  $w^{-\frac{1}{p-1}} \in A_{\frac{p}{p-1}(1-\frac{1}{n})}$ , by Lemma 2,  $w^{-\frac{1}{p-1}} \in A_{\frac{p}{p-1}(1-\frac{1}{n})-\varepsilon_1}$ , where  $\varepsilon_1 > 0$ . Now, using (2), from (15) we obtain

$$(16) \quad I_2 \leq c_8 \sum_{j=1}^{\infty} (2^j |x|)^{-\frac{n(p+1)}{p-1}} (2^j)^{\frac{n(p-1)}{p-1}} (2^j)^{-n\varepsilon_1} \int_{|z| \leq |x|} [w(z)]^{-\frac{1}{p-1}} dz \\ \leq c_9 |x|^{-\frac{n(p+1)}{p-1}} \int_{|z| \leq |x|} [w(z)]^{-\frac{1}{p-1}} dz.$$

By virtue of Lemma 3,  $w \in A_p$ , and estimating the integral on the right-hand side of (16) with the help of (1) we have

$$(17) \quad I_2 \leq c_{10} |x|^{-\frac{n(p+1)}{p-1}} |x|^{\frac{n}{p-1}} [w(B(0,|x|))]^{-\frac{1}{p-1}} = c_{10} |x|^{\frac{p}{p-1}} [w(B(0,|x|))]^{-\frac{1}{p-1}}.$$

Since  $w \in A_{p-1+\frac{\kappa}{n}}$ , by Lemma 2,  $w \in A_{p-1+\frac{\kappa}{n}-\varepsilon_2}$ , where  $\varepsilon_2 > 0$ . Using this, we obtain from Lemma 1

$$(18) \quad [w(B(0,|x|))]^{-\frac{1}{p-1}} \leq c_{11} \left[ \frac{r}{|x|} \right]^{n+\frac{p-\varepsilon_2 n}{p-1}} [w(B(0,r))]^{-\frac{1}{p-1}}.$$

Thus from (14), (17), (18) we have

$$\begin{aligned} I_1 &\leq c_{12} [w(B(0, r))]^{-1} r^{p - \varepsilon_2 n} \\ &\times \left\{ \int_{B(0, R)} \left| \frac{\partial v}{\partial x}(x) \right|^p w(x) dx \right\} \left\{ \int_0^r |x|^{n-1} |x|^{\frac{p}{n-1}} |x|^{-n - \frac{p}{n-1} + \frac{\varepsilon_2 n}{n-1}} d|x| \right\}^{p-1} \\ &\leq c_{13} \frac{r^p}{w(B(0, r))} \int_{B(0, R)} \left| \frac{\partial v}{\partial x}(x) \right|^p w(x) dx. \end{aligned}$$

Now the desired estimate follows from (12) and the last inequality.  $\square$

**Remark 3.** In case  $p = 2$  the statement of Lemma 9 coincides with the statement of Lemma 2.2 in [7].

**Proof of Theorem.** Let  $\xi$  be an arbitrary point of  $D$  and for  $0 < \varrho \leq \varrho(\xi, F)$  we define the numerical sequence

$$\varrho_j = \frac{\varrho}{4}[3 - 2^{-j}], \quad j = 1, 2, \dots$$

Let functions  $\psi_j(x)$  be equal to one on  $B_j = B(\xi, \varrho_j)$  and to zero outside  $B_{j+1}$ , and such that  $0 \leq \psi_j(x) \leq 1$ ,  $\left| \frac{\partial \psi_j(x)}{\partial x} \right| \leq \frac{2^{j+1}}{\varrho}$ .

Substitute  $\varphi(x) = [u(x, k)]^{\sigma+1} [\psi_j(x)]^{\tau+p}$  into (5), where  $\sigma, \tau$  are arbitrary positive numbers. Using A<sub>2</sub> and Young's inequality, we obtain

$$(19) \quad \int_D \left| \frac{\partial u}{\partial x} \right|^p u^\sigma \psi_j^{\tau+p} w dx \leq c_{14} (\tau + p)^p \frac{2^{jp}}{\varrho^p} [m_{j+1}]^{p-1} \int_D u^{\sigma+1} \psi_j^\tau w dx,$$

where  $m_j = \max\{u(x, k) : x \in \overline{B_j}\}$ .

Let  $t, s$  be arbitrary positive numbers satisfying the inequalities

$$t + p > \frac{pnp_0}{np_0 - p}, \quad s + p > \frac{pnp_0}{np_0 - p},$$

where  $1 < p_0 < p - 1 + \frac{p}{n}$  and  $p_0$  depends only on  $n, p, c_{p,w}$ . Then

$$\{u^{t+p}(x, k) \psi_j^{s+p}(x)\}^{\frac{nnp-p}{npnp_0}} \in \overset{\circ}{W}_p^1(B(\xi, \frac{3}{4}\varrho), w).$$

By virtue of  $A_p$ -weighted Sobolev inequality and (19) we have

$$\begin{aligned}
& \left[ \frac{1}{w(B(\xi, \frac{3}{4}\varrho))} \int_{B(\xi, \frac{3}{4}\varrho)} \left( \{u^{t+p} \psi_j^{s+p}\}_{\frac{n p_0 - p}{n p_0}}^{\frac{p n p_0}{n p_0 - p}} w dx \right)^{\frac{n p_0 - p}{p n p_0}} \right. \\
& \quad \leq c_{15} \varrho [w(B(\xi, \frac{3}{4}\varrho))]^{-\frac{1}{p}} \\
& \quad \times \left\{ (t+p)^p \int_{B(\xi, \frac{3}{4}\varrho)} \left| \frac{\partial u}{\partial x} \right|^p u^{(t+p)\frac{n p_0 - p}{n p_0} - p} \psi_j^{(s+p)\frac{n p_0 - p}{n p_0}} w dx \right. \\
& \quad \left. + (s+p)^p \int_{B(\xi, \frac{3}{4}\varrho)} u^{(t+p)\frac{n p_0 - p}{n p_0}} \left| \frac{2^j}{\varrho} \right|^p \psi_j^{(s+p)\frac{n p_0 - p}{n p_0} - p} w dx \right\}^{\frac{1}{p}} \\
& \leq c_{16} (t+s+p)^2 [w(B(\xi, \frac{3}{4}\varrho))]^{-\frac{1}{p}} 2^j [m_{j+1}]^{\frac{p-1}{p}} \\
& \quad \times \left\{ \int_{B(\xi, \frac{3}{4}\varrho)} u^{(t+p)\frac{n p_0 - p}{n p_0} - p+1} \psi_j^{(s+p)\frac{n p_0 - p}{n p_0} - p} w dx \right\}^{\frac{1}{p}}.
\end{aligned}$$

Using Lemma 1, we obtain from the last inequality

$$\begin{aligned}
& \int_{B_{j+1}} u^{t+p} \psi_j^{s+p} w dx \leq c_{17} (t+s+p)^{\frac{2 p n p_0}{n p_0 - p}} [w(B(\xi, \varrho))]^{-\frac{p}{n p_0 - p}} \\
& \quad \times \left[ 2^{j p} m_{j+1}^{p-1} \right]^{\frac{n p_0}{n p_0 - p}} \left[ \int_{B_{j+1}} u^{(t+p)\frac{n p_0 - p}{n p_0} - p+1} \psi_j^{(s+p)\frac{n p_0 - p}{n p_0} - p} w dx \right]^{\frac{n p_0}{n p_0 - p}}.
\end{aligned}$$

Choosing

$$\begin{aligned}
t = t_i &= \left[ p + \frac{n p_0}{p} (p-1) \right] \left( \frac{n p_0}{n p_0 - p} \right)^i - \frac{n p_0}{p} (p-1) - p, \\
s = s_i &= [p + n p_0] \left( \frac{n p_0}{n p_0 - p} \right)^i - n p_0 - p,
\end{aligned}$$

we rewrite the last inequality in the form

$$(20) \quad J_i \leq c_{18} \left( \frac{n p_0}{n p_0 - p} \right)^{2 i \frac{p n p_0}{n p_0 - p}} [w(B(\xi, \varrho))]^{-\frac{p}{n p_0 - p}} \left[ 2^{j p} m_{j+1}^{p-1} \right]^{\frac{n p_0}{n p_0 - p}} [J_{i-1}]^{\frac{n p_0}{n p_0 - p}},$$

where  $J_i = \int_{B_{j+1}} u^{t_i+p} \psi_j^{s_i+p} w dx$ .

By iterating we arrive at

$$(21) \quad [J_i]^{\left(\frac{n p_0 - p}{n p_0}\right)^i} \leq \left\{ c_{19} \left[ 2^{j_p} m_{j+1}^{\frac{n p_0}{n p_0 - p}} \left[ w(B(\xi, \varrho)) \right]^{\frac{-p}{n p_0 - p}} \right] \right\}^{\frac{n p_0 - p}{n p_0} + \left(\frac{n p_0 - p}{n p_0}\right)^2 + \dots + \left(\frac{n p_0 - p}{n p_0}\right)^i} \\ \times \left\{ \left( \frac{n p_0}{n p_0 - p} \right)^{2p} \right\}^{1+2\frac{n p_0 - p}{n p_0} + \dots + i \left( \frac{n p_0 - p}{n p_0} \right)^{i-1}} \times J_0.$$

When  $i$  tends to infinity, then (21) yields

$$(22) \quad [m_j]^{p+\frac{n p_0}{p}(p-1)} \leq c_{20} 2^{j n p_0} [w(B(\xi, \varrho))]^{-1} [m_{j+1}]^{\frac{n p_0}{p}(p-1)} \int_{B_{j+1}} u^p \psi_j^p w \, dx.$$

Now we estimate the integral on the right-hand side of (22) by Lemma 8 and Lemma 9:

$$(23) \quad \begin{aligned} \int_{B_{j+1}} u^p \psi_j^p w \, dx &\leq \int_{B_{j+1}} [u_{m_{j+1}}]^p w \, dx \\ &\leq c_{21} \varrho^p \int_{E_{m_{j+1}}} \left| \frac{\partial u}{\partial x} \right|^p w \, dx \\ &\leq c_{22} \varrho^p m_{j+1} k^{p-1} \text{cap}_{p,w}(F). \end{aligned}$$

By virtue of (22), (23) implies

$$(24) \quad [m_j]^{p+\frac{n p_0}{p}(p-1)} \leq c_{23} 2^{j n p_0} \frac{\varrho^p}{w(B(\xi, \varrho))} [m_{j+1}]^{1+\frac{n p_0}{p}(p-1)} k^{p-1} \text{cap}_{p,w}(F),$$

Further we shall use the following:

**Lemma 10.** *Let  $\{\alpha_i\}$  be a bounded number sequence satisfying*

$$\alpha_i \leq A \alpha_{i+1}^\sigma, \quad i = 1, 2, \dots$$

*with positive constants  $A, a, \sigma \in (0, 1)$ . Then we have*

$$\alpha_1 \leq c A^{\frac{1}{1-\sigma}}$$

*with a constant  $c$  depending only on  $\sigma$  and  $a$ .*

For the proof of Lemma 10 see [1], Chapter 5.

Finally, from (24) and Lemma 10 we have

$$m_1 \leq c_{24} k \left\{ \text{cap}_{p,w}(F) \frac{\varrho^p}{w(B(\xi, \varrho))} \right\}^{\frac{1}{p-1}},$$

and so the inequality (7) is established.

This completes the proof of Theorem.  $\square$

#### References

- [1] Skrypnik I. V.: Nonlinear elliptic boundary value problems. B. G. Teubner Verlag, Leipzig, 1986.
- [2] Skrypnik I. V.: New conditions of homogenization of nonlinear Dirichlet problems in perforated domains. Ukrainian Math. J. 48 (1996), no. 5, 675–694.
- [3] Heinonen J., Kilpelainen T., Martio O.: Nonlinear potential theory of degenerate elliptic equations. Clarendon Press, Oxford, 1993.
- [4] Chanillo S., Wheeden R. L.: Weighted Poincaré and Sobolev inequalities and estimates for weighted Peano maximal functions. Amer. J. Math. 107 (1985), 1191–1226.
- [5] Kufner A.: Weighted Sobolev spaces. B. G. Teubner Verlag, Leipzig, 1980.
- [6] Gutierrez C. E., Nelson G. S.: Bounds for the fundamental solution of degenerate parabolic equations. Commun. Partial Differential Equations 13 (1988), no. 5, 635–649.
- [7] Leonardi S., Skrypnik I. I.: Necessary condition for regularity of a boundary point for a degenerate quasilinear parabolic equations. Catania Univ., Catania, 1995, preprint.

Authors' addresses: Igor V. Skrypnik, Dmitry V. Larin, Institute Appl. Math. & Mech. NAS Ukraine, R. Luxemburg Str. 74, 340114 Donetsk, Ukraine, e-mail: skrypnik @iamm.ac.donetsk.ua.