

# Competition for Electrons

## Aim

- write equations for oxidation and reduction half reactions

## Notes

### Atoms compete for each other's electrons

- When chemical bonds form, electrons are either lost, gained or shared
- Oxidation-Reduction reactions (Redox reactions)
  - Metals
    - lose electrons (OXIDATION)[NOTE: as when metals combine with oxygen]
    - are oxidized
    - are reducing agents
  - Nonmetals
    - gain electrons reducing their oxidation states (REDUCTION)
    - are reduced
    - are oxidizing agents

**Oxidation**  
**Is**  
**Loss**  
  
**Reduction**  
**Is**  
**Gain**

- Example 1 -  $2\text{Mg(s)} + \text{O}_2\text{(g)} \rightarrow 2\text{MgO(s)}$

| Mg   | O <sub>2</sub>                    |
|--|-----------------------------------|
| ★ loses electrons                          | ★ gains electrons                 |
| ★ gets oxidized to Mg <sup>2+</sup>        | ★ gets reduced to O <sup>2-</sup> |
| ★ is the reducing agent for O <sub>2</sub> | ★ is the oxidizing agent for Mg   |

- Half reactions — reaction showing either a gain or loss of electrons
  - $2\text{Mg}^0 \rightarrow 2\text{Mg}^{2+} + 4\text{e}^-$
  - $\text{O}_2^0 + 4\text{e}^- \rightarrow 2\text{O}^{2-}$
- Net equation (REDOX REACTION)— combination of the half reactions such that the number of electrons lost equals the number of electrons gained
 
$$2\text{Mg(s)} + \text{O}_2\text{(g)} \rightarrow 2\text{MgO(s)}$$
- Example 2 - More active metals replace less active metals in compounds by transferring electrons to them
  - Sample Reaction:
 
$$\text{Zn(s)} + \text{Cu(NO}_3)_2\text{(aq)} \rightarrow \text{Zn(NO}_3)_2\text{(aq)} + \text{Cu(s)}$$
  - Half reactions — reaction showing either a gain or loss of electrons
    - $\text{Zn}^0 \rightarrow \text{Zn}^{2+} + 2\text{e}^-$
    - $\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu}^0$
  - Net equation — combination of the half reactions such that the number of electrons lost equals the number of electrons gained
 
$$\text{Cu}^{2+} + \text{Zn}^0 \rightarrow \text{Zn}^{2+} + \text{Cu}^0$$
  - Spectator ions — ions that are present during a reaction but do not participate in the reaction:
 
$$2\text{NO}_3^-$$

**Oxidation number (Oxidation state)** - number assigned to keep track of electrons based on the arbitrary assumption that shared electrons belong to the more electronegative element

- Rules for assigning oxidation numbers
  - Oxidation numbers for atoms that are free elements are always zero
  - The oxidation numbers of ions are the same as the charge on the ion
  - Some elements have only one oxidation state
    - group 1 metals always form 1+ ions and always have a +1 oxidation state
    - group 2 metals always form 2+ ions and always have a +2 oxidation state
  - Some elements usually have a particular oxidation state
    - oxygen has a -2 oxidation state except in peroxides where it is -1 and in compounds with fluorine (OF<sub>2</sub>) where it is +2
    - hydrogen has a +1 oxidation state except in hydrides with group 1 and group 2 metals
  - the sum of the oxidation numbers
    - in a compound it is always zero
    - in a polyatomic ion it is equal to the charge on the ion
- Finding oxidation numbers
  - apply the rules
  - construct a table if necessary

### Sample Problem

Find the oxidation state of the elements in K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>.

| Element                 | K  | Cr | O   | TOTAL |
|-------------------------|----|----|-----|-------|
| Subscript               | 2  | 2  | 7   |       |
| Oxidation state         | +1 | ?  | -2  |       |
| Sum of oxidation states | +2 | ?? | -14 | 0     |

- [a] potassium is a group one metal; its oxidation state is always +1
- [b] oxygen usually has an oxidation state of -2
- [c] the sum of oxidation states of each element is the product of the subscript and the oxidation state
- [d] find the -sum of the oxidation states of chromium (??) by setting the sum of all the oxidation states to zero
 
$$\begin{array}{rcl}
 (+2) + ?? + (-14) & = & 0 \\
 ?? & = & +12
 \end{array}$$
- [f] find the oxidation state of chromium (?) by dividing the sum (+12) by the subscript (2)
 
$$+12 \div 2 = +6$$

Answer the questions below by circling the number of the correct response

- In this reaction, the oxidation number (oxidation state) of C changes from:  $2\text{CO}_2 \rightarrow 2\text{CO} + \text{O}_2$   
(1) 0 to +4 (2) +2 to +4 (3) +3 to 0 (4) +4 to +2
- In the reaction:  
 $2\text{KMnO}_4 + 3\text{H}_2\text{SO}_4 + 5\text{H}_2\text{S} \rightarrow 5\text{S} + 2\text{MnSO}_4 + \text{K}_2\text{SO}_4 + 8\text{H}_2\text{O}$   
the oxidation number of sulfur changes from  
(1) +5 to -5 (2) -5 to +5 (3) 0 to -2 (4) -2 to 0
- What is the oxidation number of Cr in  $\text{Na}_2\text{CrO}_4$ ?  
(1) +1 (2) +2 (3) +3 (4) +6
- What is the oxidation state of the chromium in  $\text{K}_2\text{Cr}_2\text{O}_7$ ?  
(1) +5 (2) +6 (3) +3 (4) +12
- In the reaction  $\text{Pb} + 2\text{Ag}^+ \rightarrow \text{Pb}^{2+} + 2\text{Ag}$ , the reducing agent is  
(1) Ag (2)  $\text{Ag}^+$  (3) Pb (4)  $\text{Pb}^{2+}$
- Which is not an oxidation-reduction reaction?  
(1)  $4\text{Na} + \text{O}_2 \rightarrow 2\text{Na}_2\text{O}$   
(2)  $\text{Fe} + 2\text{HCl} \rightarrow \text{FeCl}_2 + \text{H}_2$   
(3)  $\text{CaCl}_2(\text{aq}) + 2\text{AgNO}_3(\text{aq}) \rightarrow 2\text{AgCl}(\text{s}) + \text{Ca}(\text{NO}_3)_2(\text{aq})$   
(4)  $2\text{H}_2\text{O} \rightarrow 2\text{H}_2 + \text{O}_2$
- Given:  $2\text{Al} + 3\text{Zn}^{+2} \rightarrow 2\text{Al}^{+3} + 3\text{Zn}$ . In this reaction, the oxidizing agent is (1) Al (2)  $\text{Al}^{+3}$  (3) Zn (4)  $\text{Zn}^{+2}$
- Given:  $2\text{Al} + 3\text{Zn}^{+2} \rightarrow 2\text{Al}^{+3} + 3\text{Zn}$ . In this reaction, electrons are transferred from (1) Al to  $\text{Al}^{+3}$  (2)  $\text{Zn}^{+2}$  to Zn (3) Al to  $\text{Zn}^{+2}$  (4)  $\text{Zn}^{+2}$  to Al
- What is the oxidation number of nitrogen in  $\text{N}_2\text{O}_3$ ? (1) +1 (2) +2 (3) +3 (4) +6
- In the reaction  $3\text{CO} + \text{Fe}_2\text{O}_3 \rightarrow 3\text{CO}_2 + 2\text{Fe}$ , the oxidation number of the iron changes from (1) +2 to 0 (2) +2 to +3 (3) +3 to +2 (4) +3 to 0
- What is the oxidation number of Br in  $\text{BrO}_3^{-2}$ ?  
(1) +1 (2) +6 (3) +5 (4) +4
- Which is the reducing agent in the following reaction?  
 $\text{Cl}_2(\text{aq}) + 2\text{KBr}(\text{aq}) \rightarrow 2\text{KCl}(\text{aq}) + \text{Br}_2(\text{aq})$   
(1)  $\text{Cl}_2$  (2)  $\text{H}_2\text{O}$  (3)  $\text{K}^+$  (4)  $\text{Br}^-$
- What is the oxidation number of carbon in  $\text{C}_2\text{O}_4^{-2}$ ?  
(1) +1 (2) +2 (3) +3 (4) +4
- Which is an oxidation-reduction reaction?  
(1)  $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$   
(2)  $\text{KOH} + \text{HBr} \rightarrow \text{KBr} + \text{H}_2\text{O}$   
(3)  $\text{AgNO}_3 + \text{NaCl} \rightarrow \text{AgCl} + \text{NaNO}_3$   
(4)  $\text{Mg} + \text{Cl}_2 \rightarrow \text{MgCl}_2$
- $\text{MnSO}_4$  is a product in a reaction that contained  $\text{KMnO}_4$  as a reactant. The oxidation number of the manganese changed from (1) -2 to +5 (2) +7 to +2 (3) +5 to -2 (4) -7 to +2
- Given the balanced equation:  
 $2\text{HNO}_3 + 3\text{H}_2\text{S} \rightarrow 4\text{H}_2\text{O} + 2\text{NO} + 3\text{S}$   
Which is reduced? (1) S (2) S-2 (3) N+2 (4) N+5
- During the reaction  $\text{Ca} + \text{H}_2 \rightarrow \text{CaH}_2$ , the oxidation number of the hydrogen changes from  
(1) 0 to +1 (2) +1 to 0 (3) 0 to -1 (4) -1 to 0
- In the reaction  $\text{Sn}^{+4} + \text{H}_2(\text{g}) \rightarrow \text{Sn}^{+2} + 2\text{H}^+$ , the reducing agent is  
(1)  $\text{Sn}^{+4}$  (2)  $\text{H}_2$  (3)  $\text{Sn}^{+2}$  (4)  $\text{H}^+$
- Given:  $3\text{Ag} + 4\text{HNO}_3 \rightarrow \text{NO} + 3\text{AgNO}_3 + 2\text{H}_2\text{O}$ . The reducing agent in this reaction is  
(1) Ag (2)  $\text{Ag}^{+1}$  (3)  $\text{H}^{+1}$  (4)  $\text{N}^{+2}$
- The reaction  $\text{NaCl}(\text{s}) \rightarrow \text{Na}^+(\text{aq}) + \text{Cl}^-(\text{aq})$  is an example of  
(1) an oxidation reaction, only  
(2) a reduction reaction, only  
(3) both an oxidation and a reduction reaction  
(4) neither an oxidation nor a reduction reaction
- The oxidation number of manganese in  $\text{KMnO}_4$  is  
(1) +1 (2) +7 (3) +3 (4) +4
- In the reaction  $\text{Sn}^{+2} + 2\text{Fe}^{+3} \rightarrow \text{Sn}^{+4} + 2\text{Fe}^{+2}$ , the reducing agent is  
(1)  $\text{Fe}^{+2}$  (2)  $\text{Fe}^{+3}$  (3)  $\text{Sn}^{+2}$  (4) Sn
- An oxidizing agent will always  
(1) lose electrons (3) be reduced  
(2) increase in oxidation number (4) increase in mass