

I, \_\_\_\_\_ understand the following list of risks for taking Isotretinoin and still strongly desire to pursue treatment.

\_\_\_\_\_ **Body as a whole:** allergic reactions, including vasculitis, systemic hypersensitivity, edema, fatigue, lymphadenopathy, weight loss

\_\_\_\_\_ **Cardiovascular:** palpitation, tachycardia, vascular thrombotic disease, stroke

\_\_\_\_\_ **Endocrine/Metabolic:** hypertriglyceridemia, alterations in blood sugar level

\_\_\_\_\_ **Gastrointestinal:** Inflammatory bowel disease, hepatitis, pancreatitis, bleeding and inflammation of the gums, colitis, ileitis, nausea, other nonspecific gastrointestinal symptoms

\_\_\_\_\_ **Hematologic:** allergic reactions, anemia, thrombocytopenia, neutropenia, agranulocytosis

**Musculoskeletal:** skeletal hyperostosis, calcification of tendons and ligaments, premature epiphyseal closure, mild to moderate musculoskeletal symptoms including arthralgia, transient pain in the chest, elevations of CPK, arthritis, tendonitis, other types of bone abnormalities

\_\_\_\_\_ **Neurological:** pseudotumor cerebri, dizziness, drowsiness, headache, insomnia, lethargy, malaise, nervousness, paresthesias, seizures, stroke, syncope, weakness

\_\_\_\_\_ **Psychiatric:** suicidal ideation, suicide attempts, suicide, depression, psychosis, emotional instability

\_\_\_\_\_ **Reproductive System:** abnormal menses, birth defects

\_\_\_\_\_ **Respiratory:** bronchospasms (with or without history of asthma), respiratory infection, voice alteration

\_\_\_\_\_ **Skin and Appendages:** acne fulminans, alopecia (hair loss), bruising, cheilitis (dry lips), dry mouth, dry nose, dry skin, epistaxis, eruptive xanthomas, flushing, fragility of skin, hair abnormalities, hirsutism, hyperpigmentation and hypopigmentation, infections (including disseminated herpes simplex), nail dystrophy, paronychia, peeling of palms and soles, photoallergic/photosensitizing reactions, pruritus, pyogenic granuloma, rash (including facial erythema, seborrhea, and eczema), sunburn susceptibility increased, sweating, urticaria, vasculitis (including Wegener's granulomatosis, abnormal wound healing

\_\_\_\_\_ **Special Senses:** hearing impairment, corneal opacities, decreased night vision, cataracts, color vision disorder, conjunctivitis, dry eyes, eyelid inflammation, keratitis, optic neuritis, photophobia, visual disturbances

\_\_\_\_\_ **Urinary System:** glomerulonephritis, nonspecific urogenital findings, elevation of plasma triglycerides, decrease in serum high-density lipoprotein levels, elevations of serum cholesterol during treatment, increased alkaline phosphatase, SGOT, SGPT, GGTP or LDH, elevation of fasting blood sugar, elevations of CPK, hyperuricemia, decreases in RBC parameters, decreases in white blood cell counts, elevated sedimentation rates, elevated platelet counts, thrombocytopenia

\_\_\_\_\_  
Printed Name

\_\_\_\_\_  
Signature

\_\_\_\_\_  
Date

I have also discussed with \_\_\_\_\_ all attendant risks to this treatment which include, but are not limited to: allergic reactions, including vasculitis, systemic edema, fatigue, lymphadenopathy, weight loss, palpitation, tachycardia, vascular thrombotic disease, stroke, hypertriglyceridemia, alterations in blood sugar levels, inflammatory bowel disease, hepatitis, pancreatitis, bleeding and inflammation of the gums, colitis, ileitis, nausea, other nonspecific gastrointestinal symptoms, allergic reactions, anemia, thrombocytopenia, neutropenia, rare reports of agranulocytosis, skeletal hyperostosis, calcification of tendons and ligaments, premature epiphyseal, mild to moderate musculoskeletal, symptoms including arthralgia, transient pain in the chest, elevations of CPK, arthritis, tendonitis, other types of bone abnormalities, pseudotumor cerebri, dizziness, drowsiness, headache, insomnia, lethargy, malaise, nervousness, paresthesias, seizures, stroke, syncope, weakness, suicidal ideation, suicide attempts, suicide, depression, psychosis, emotional instability, abnormal menses, bronchospasms (with or without a history of asthma), respiratory infection, voice alteration, acne fulminans, alopecia (which in some cases persists), bruising, cheilitis (dry lips), dry mouth, dry nose, dry skin, epistaxis, eruptive xanthomas, flushing, fragility of skin, hair abnormalities, hirsutism, hyperpigmentation and hypopigmentation, infections (including disseminated herpes simplex), nail dystrophy, paronychia, peeling of palms and soles, photoallergic/photosensitizing reactions, pruritus, pyogenic granuloma, rash (including facial erythema, seborrhea, and eczema), sunburn susceptibility increased, sweating, urticaria, vasculitis (including Wegener's granulomatosis), abnormal wound healing, hearing impairment, tinnitus, corneal opacities, decreased night vision which may persist, cataracts, color vision disorder, conjunctivitis, dry eyes, eyelid inflammation, keratitis, optic neuritis, photophobia, visual disturbance, glomerulonephritis, nonspecific urogenital finding, Elevation of plasma triglycerides, decrease in serum high-density lipoprotein (HDL) levels, elevations of serum cholesterol during treatment, Increased alkaline phosphatase, SGOT (AST), SGPT (ALT), GGTP or Elevation of fasting blood sugar, elevations of CPK, hyperuricemia, Decreases in red blood cell parameters, decreases in white blood cell counts (including severe neutropenia and rare reports of agranulocytosis), elevated sedimentation rates, elevated platelet counts, thrombocytopenia, and possibly death. The patient has verbalized her comprehension of all of these risks and strongly desires to proceed.

---

Provider's Name

---

Provider's Signature

Date