[Home] [Simple] [Français] [Good Food] [Bad Food] [Vitamins +] [Books & Links] [Email] [Diabetes & Weight] [Where] [Cholesterol] [Why] [More on Causes] [31 Tips & Info] [Index]

Nutrition, Health & Heart Disease




Cholesterol and its drugs: info to share with your Dr. and friends. Is low-cholesterol really healthy? Does lowering it extend life?




Causés muscle & joint pain, "senior moments"

C27·H45·OH - High cholesterol was more likely a 'defense factor' than a 'risk factor' prior to refined foods. The famed Framingham study found better thinking at high cholesterol while decreasing levels after age 50 predicted more heart disease deaths!



Cholesterol Pills, statins
like Lipitor, Zocor,
Pravachol, Lescol, Mevacor
or Crestor, (killer) fibrates
and niacin (mega vitamin
B3) have side effects, good,
bad and unknown. Studies
show that lowering
cholesterol with statin,
fibrates or omega-6 oils
(but not niacin or fiber)
promotes cancer [general
effects and seniors].

The *more* statins cut cholesterol, the *more* they cut energy to nerves, muscle and heart by lowering CoQ10, 'battery acid' and anti-oxidant ('catalytic converter') of *every* cell. Here's a study with *best* cholesterol at the not exactly low old "normal" range of 180-240 (4.6-6.0) -also: 1, 2. Great sites: thincs.org and Cholesterol Myths.



Drug and food interests create cholesterol fear but statins don't work by lowering it and don't save women: BMJ Feb., JAMA May, JACC Sept. 2004, CMAJ Nov. 2005.

More cancer and no survival benefit (±0.1%) in high risk older Europeans: <u>Lancet</u> 2002 (PROSPER study: 3 million \$3 pills taken).

Next, in 6 years, nobody saved (±0.07%) in this younger group of Americans (ALLHAT study: graph below).


PS-1 In heart failure: lowest 20% for cholesterol: double the death rate!


PS-2 TNT trial implodes: 5000 heart patients for 5 years on 80 mg top-dose *Lipitor* suffer 2 *more* deaths than patients on only 10 mg. On either dose, the same $26 \pm 1\%$ progression in artery calcium in 12 months anyhow! *Lipitor* does NOT save lives in studies.

Out of 1/2 million U.S. men at 'prime heart attack age', the top for cholesterol [292 mg/dL (7.5 mM)] on anti-cholesterol absorbing drug for 7.4 years (and 1/8th less "bad LDL" compared with dummy drug) but no difference in survivors; count them: 3. And, oeps: in 65,000 men with 2700 deaths, those with <u>least</u> cholesterol [below 187 (4.8)] had <u>most</u> deaths!


PS-4 The American College of Cardiology in 2004: "... there is no evidence for a total mortality benefit in women from dyslipidemia [statin] treatment." Women: 3 more deaths on Zocor and 2 more heart 'events' on Lipitor in large studies (4S, ASCOT). In 24,000 women over age 50, those in the lowest 25% for cholesterol had the same risk for death (+60%) as those smoking!


Some general data about cholesterol. Later, more about drugs.







those taking the placebo (ASPEN study)









Cholesterol lowering: no mortality benefit in women, elderly, but in men?



Crestor's massive JUPITER study found no cardiovascular mortality benefit .. and per 10 NON fatal strokes or heart attacks prevented, 8 new diabetics were reported. Most benefit was in non life-saving procedures like balloon angioplasties.

So much for having 50% less 'bad' LDL cholesterol click for results New Engl J Med 11-2008

CHOLESTEROL PILLS, statins,
-Pravachol, Lipitor, Zocor,
Mevacor, Lescol, Crestor- '..if
diet and exercise alone are not
enough..' make part of a lab
report look 'better'. This makes
families poorer and less
concerned about food choices
while most doctors are not yet
blamed for prescribing them.



They hardly raise HDL and

<u>slash</u> the production of CoQ10 by the same % as LDL, forcing heart, in fact all cells, to work with less energy and more 'free radical' damage. Their limited effect is <u>clearly not</u> from cholesterol lowering but most probably from <u>imitating (cheap and usually safe)</u> <u>nitroglycerin</u> (click on picture to the left) and possibly by reducing inflammation, as would <u>(cheap but not so safe) aspirin</u>, omega-3 fish oil in pills, and many vitamins in a multivitamin, including E).

Those doctors aware of the fact that the statin studies can be explained by their powerful nitroglycerin mimicking effect (and

thus less non-fatal angina [heart pain] and angina driven non-fatal benefit) find the concept that they may have been wrong about cholesterol too terrible to contemplate, yet that is what most studies show, including *all Lipitor* and *Crestor* (atorva- and rosuva-statin) studies.

<u>Stanol-sterol</u> margarines, policosanol, fibrates, gugulipid, dextrothyroxine, estrogen, <u>ezetimibe [Zetia, Ezetrol, part of Vytorin/Enigy -click for warning]</u> and <u>'killer torcetrapib</u> all 'manage cholesterol' in various ways but have never shown to save lives, just like *Lipitor*.

Medscape and MedlinePlus list 100 side effects but not a most vital one, CoQ10 reduction. Simply put:

1. Good: Statins imitate nitroglycerin, resulting in less non-fatal chest pain ('stable' angina) and thus hospital visit and thus about 20-45% fewer stents. Example: *Lipitor* in the ASCOT study, and now the 2008 JUPITER study with fewer stents but zero cardiac deaths prevented after 17,000 years of *Crestor* use!! BETTER: To avoid 90% stents and bypasses, not just 20 or 45%, when having a heart attack, first go to a close and small hospital that cannot "go in" and statistically survival at 6 months is better. 'Planned' angioplasty (stents) also does not prevent heart attacks, make you live longer or make your life quality better;

- 2. No effect: from changes to cholesterol [lipids]; 'familial' (genetically high) patients may be different but the science is incomplete;
- 3. Bad: pain (joints, weakness), 'senior moments'/memory (try niacin instead!), less CoQ10, more Lp(a), birth defects and bleeding strokes, nerve repair problems and likely cancer;
- 4. <u>Scary ads</u> & faulty science [*Zocor*, *Vytorin* and the *Pfizer* feet in the morgue fear of death campaign called 'unethical' by the World Health Organization];
 - 5. Don't prolong survival in most high-risk groups and not in women:



Massive benefits proclaims Oxford University about its 2002 Zocor -Heart
Protection Study. 75% of heart attacks still happen and 300 on Zocor for 1 year to
postpone one (1) death. Massive drug use, few 'men' saved. The next study
claiming benefit in high-risk elderly, Pravachol's PROSPER had 28 fewer heart but 24
more cancer deaths and more 'new' cancers in each of 4 years in a group with 52 fewer smokers.
Next, zero 'anything' was found in Pravachol's ALLHAT in high-risk North Americans: 1 death
postponed per 1.1 million \$3 pills taken! Next was Lipitor's ASCOT also without mortality benefit
(graph below). From 10 years ago: cholesterol lowering by any means caused 150 more deaths per
100,000 patient-years of intervention. Cheers. No statin benefit in "primary" prevention in 2003,
click on the right picture: the University of British Columbia.

Q 1 2 CoQ10 9 10

medline 8241697

Lipitor, Zocor, etc. lower CoQ10 by the same % as LDL-cholesterol. No Q10 and a cell dies from lack of energy. Q10 is also the best heart failure drug, by far.

"Ask your doctor if "Rx" is right for you" prompt the ads but vital information is hidden from you and your doctor. This paves the way for after-the-damage criminal investigations and <u>lawsuits</u>.

<u>Statins</u>, fibrates and high intakes of the common

cholesterol-lowering polyunsaturated omega-6 'vegetable' oils (like corn) are linked to more cancer, as is just plain low cholesterol. Canadian ads but not U.S. ads warn that statins lower CoQ10; there's your

muscle and nerve pain, <u>and lack of energy</u>. Fewer studies were done on women and elderly, groups where high LDL has no evident link to mortality [low HDL does have such link, but HDL is best raised by, for example, exercise, moderate alcohol use or mega-niacin *with* -or it will raise homocysteine- a multivitamin]. No mortality studies were done in people taking whole foods, omega-3 oil and a

NO COQ10 YES
statin drug ads vary

good multi.

Many B-vitamin (homocysteine) trials are now being done but some are in people where prevention is rather (too) late, some were destined for failure (VISP) and many are set-up to have statins share the credit. The latter paves the way for *combination therapy*: statins + <u>niacin</u> and/or aspirin and/or <u>fish oil</u> and/or multi-B-vitamins ... if statins alone are not enough ...

A TV ad exclaims: 'Lipitor did it, the lower numbers you're looking for.' Then, a small banner: "Lipitor has not been shown to prevent heart disease ..." Isn't that

what people expect when taking the drug? Now, in 2005, Lipitor ads no longer mention heart disease, it

simply 'treats' (bad) cholesterol, a concept created by the drug and food industry. Another TV couple: 'Oh, Jessie, I love you too! ... Zocor, Be There! Up to 87% of Baycol* users drop to 'target levels' but 'effects on disease and death have not been established.' Why not real wellness goals such as less illness, fewer doctors visits or more time with friends?

Eight authors of the U.S. ATP III 2004 treatment guideline 'update' got money from average 10 drug companies; one, a NIH gov't employee, got \$114,000 + 'stock options' from the statin industry. The 9th author earns a living as NCEP Coordinator with the job to '.. increase the proportion [.. of Americans] who adhere to their cholesterol-lowering regimen.' And, oeps, they forgot the largest statin study ever, J-LIT (graphs at end). More in BMJ on study conflicts of interest. Could the brilliant beauty of a drug (industry) blind a well meaning doctor? Is it fear of not following 'guidelines'? *)Baycol (Lipobay) was withdrawn for causing unexpected deaths: here's an important comment about all statins. Also, StatinAlert.org or StoppedOurStatins.



How about *Pravachol*? Two analysis by the Journal Club here and here; some high risk participant would have to be treated for over 200 years at a drug cost of \$200 000 to prevent one cardiac 'event'. The WOSCOPS study selected 6600 men out of 160 000 and treated half. After 5 years and 30% giving up on the drug total deaths were not quite statistically different. The CARE study found, in patients with prior heart attacks '.. no significant differences in overall mortality

..' [1.6 per 1000 years of drug use, a statistical fluke.] *NEJM*; 99-4-8:1115 about a 3rd study (LIPID): '[Pravachol] has no particular advantage over placebo [dummy-pill].' Finally the truly massive ALLHAT study found 'zip' in any health department!



Also more dead on nasty Lopid -fibrates kill people-

(WHO & FIELD trials)

-EXCELdummy pill 3 deaths Mevacor per group 11 months: studv killed

The Mevacor (lovastatin) EXCEL trial had with 89% probability 2.75x more deaths (97%) cardio-vascular) after 2 million pills were swallowed (11% fewer heart attacks, 40% more cancer); then, dumping 97% of the placebo group, Merck continued but without the risk of the drug proving conclusively to be more fatal than a dummy pill -- and dead patients kill drugs! Then, after 3 times more *Mevacor* pills, the AFCAPS / TexCAPS trial ended with 3 more drug deaths (also 40% more cancer). Next, Merck's 1st Zocor study (4S) killed 3 more women but saved men, yet with unexplained anomalies in the mortality curves after 18

months. incidentally: statistics & data entry by Merck.



Scandal at Oxford: Sir Peto and Dr. Collins refuse to publish the Heart Protection Study mortality curves in men, women, diabetic or not. In 1992 both wanted 'total mortality' trials [they then thought this was relevant]; they did the study and now hide the mortality curves. In 2004 Dr. Collins suggested they would release these data but did not. In April 2007, author Dr. Baigent said it would be 'inappropriate' to publish these data and author Dr. Sleight: 'It is not my decision, but it looks like that.' [not releasing data about deaths]. Study sponsor Merck is also mum about deaths but as always, women did not benefit. If you're a diabetic women with a 75% chance to die from blood vessel diseases, would you not like to know how many days of life you might statistically gain or lose taking Zocor for 5 years? That is not an 'inappropriate' question but a real

one and even more so when balancing money for food or drugs.

Twenty percent of 1st time *Zocor* users were motivated by cholesterol fear instilled by a <u>football</u> coach (*Merck* annual report). <u>Pfizer</u> uses golfers and, <u>before Feb. 28 2008</u>, <u>a</u> doctor who never practiced.



A shocking statement on the <u>ALLHAT</u> website (2007): "..trials demonstrating a reduction in coronary heart disease from cholesterol lowering have not [sic] demonstrated a net reduction in mortality." <u>MedlinePlus</u> about ALLHAT: "..people taking pravastatin were no less



likely to die or develop heart disease than people receiving usual care." *Lipitor*'s ASCOT 3.3 year mortality result is no different: 5168

people on statin, 5 million pills swallowed and the curves touch, and cancers, quality of life and muscle pain not reported. The 5 year SPARCL *Lipitor* trial ended with 5 *fewer* deaths on dummy pills while the ASPEN study ended with 2 fewer deaths on placebo (2nd graph from top).

Statins aren't cholesterol but *mevalonate* lowering 'therapy'. This mother-molecule also makes CoQ10 and other vital stuff with names like <u>isoprene</u> [it takes 10 to make Q10, 6 to make cholesterol], farnesyl, geranyl, dolichol and squalene. Similarly, *aspirin* is not simply "anti-clot therapy", it rearranges all fatbased "house keeping", pain and signal machinery. Statins and aspirin hamper (*inhibit*) fundamental body processes. Whatever

aspirin hamper (*inhibit*) fundamental body processes. Whatever good necessarily comes with the bad. On the other hand, niacin *promotes* over 300 reactions and has few harmful effects since it *stimulates* body processes, like all vitamins. Niacin *must* be taken with a multi-B-vitamin or it will raise homocysteine. It is just a partial answer to heart disease and has other benefits but <u>niacin did</u> <u>prevent</u>

heart attacks and *did* save lives; 2006 update.





Bottom Line: At 10 - 36x the price of generic niacin (1.2 kg for \$50 or 80¢ per week here)*, in some types of (mainly male and noncongestive) heart patients, some statins may reduce the risk of "events" by one fifth [probably not "deaths", and you have to scrape the statistical barrel]. This leaves no less than 4/5ths of the risk. To deal with the larger under lying problem, remember that fat substitutions have proven of no value, apart from those involving omega-3 oils, and that whole-foods and supplements cheaply lower most risk factors -and survival chances. *(tel. 1.800.544.4440 or 1.954.766.8433; product 94.)

A doctor prescribing statin and not first niacin plus a good multi vitamin, without tracking homocysteine and Lp(a), and without replacement CoQ10, canola and fish omega-3 (pills) and likely a magnesium supplement is practicing incomplete medicine.

Mega-niacin [about 1/3rd of 2 grams/day after meals] does take some medical guidance and for the first few days it will give a harmless but frightening hot-flush. It is the only drug suggested by the AHA to raise HDL and it's 1 of 3 to lower LDL [including fibrates, a drug family increasing deaths -see below]. Best: plain niacin (not 'no-flush') and it may make your doctor feel better about law suits since you're doing something about 'your' cholesterol. Always take with a multi otherwise it raises homocysteine and, bonus: it lowers clotting Lp(a) and fibrinogen Why not share this info with the Dr. signing for your drugs?

Clinical results (oeps):





Note: this 'cholesterol summary' PDF is extracted from the body of www.health-heart.org 2009.5.29