

## Vascular Surgery Stent/Graft Plan

[R] = will be ordered

T= Today; N = Now (date and time ordered)

Height: \_\_\_\_\_ cm Weight: \_\_\_\_\_ kg

**Allergies:** ☐ No known allergies

☐ Medication allergy(s): \_\_\_\_\_

☐ Latex allergy ☐ Other: \_\_\_\_\_

### Admission/Transfer/Discharge

<input type="checkbox"/>	Patient Status Initial <b>Inpatient</b>	Attending Physician: _____
	Bed Type: <input type="checkbox"/> Med Surg <input type="checkbox"/> Critical Care <input type="checkbox"/> Stepdown <input type="checkbox"/> Obstetrics <input type="checkbox"/> Other	
<input type="checkbox"/>	Patient Status Initial <b>Outpatient</b>	Attending Physician: _____
	Outpatient Status/Service: <input type="checkbox"/> OP-Ambulatory <input type="checkbox"/> OP-Diagnostic Procedure <input type="checkbox"/> OP-Observation Services	
	<b>NOTE to MD:</b>	
	<b>Initial status – inpatient</b> --- For a condition/dx with severity of illness or co-morbid conditions indicating a hospital stay greater than 24 hours is required.	
	<b>Initial Status Outpatient – Ambulatory surgery</b> – Outpatient surgery/procedure with discharge anticipated after a routine or, in some cases, extended recovery. <ul style="list-style-type: none"> <li>• Routine recovery after outpatient surgery is estimated at 6-8 hours.</li> <li>• “Extended” routine recovery may be required for a patient to stay longer (could be overnight) to recover from anticipated sequela of surgery including effects of anesthesia, nausea, pain.</li> <li>• For unanticipated sequela of surgery or a complicated post operative course, the patient may require a status change to inpatient. Please consult with a case manager before making this choice of “status change”.</li> <li>• Examples: Initial status outpatient is generally selected for patients undergoing PCI, diagnostic caths, EP studies, ablations, pacemaker implantations, other routine surgeries.</li> </ul>	
	<b>Initial status Outpatient -Observation Services</b> – Short term treatment, assessment and reassessment - estimate discharge within 24 hours <ul style="list-style-type: none"> <li>• In some cases (for Medicare patients), this can be extended to 48 hours.</li> <li>• Observation Services can also be utilized when it is unclear (without additional assessment) whether the patient will require an inpatient stay.</li> </ul>	
<input type="checkbox"/>	Notify physician once _____	T;N, of room number on arrival to unit

Primary Diagnosis: \_\_\_\_\_

Secondary Diagnosis: \_\_\_\_\_

### Laboratory

<input type="checkbox"/>	Creatinine	T;N, Routine, Once, Type: Blood
<input type="checkbox"/>	BUN	T;N, Routine, Once, Type: Blood

### Diagnostic Tests

<input type="checkbox"/>	Chest 2VW Frontal & Frontal & Lat	T;N, Reason for Exam: Other, Enter in Comments, Routine, Order Comments: follow-up thoracic aortic stent grafting for aortic transection
<input type="checkbox"/>	KUB	T;N, Routine
<input type="checkbox"/>	EKG	T;N, Routine
<input type="checkbox"/>	CT Abdomen Triple Phase	T;N, Reason for Exam: Other, Enter in Comments, Order comments: For Aortic Stent Graft, 2.5 mm cuts - no contrast, then 2.5 mm cuts - arterial phase of contrast, then 2.5 mm cuts, venous phase. NO Oral contrast.
<input type="checkbox"/>	CT Pelvis W Cont	T;N, Reason for Exam: Other, Enter in Comments, Order comments: For Aortic Stent Graft, 2.5 mm cuts - no contrast, then 2.5 mm cuts - arterial phase of contrast, then 2.5 mm cuts, venous phase. NO Oral contrast.
<input type="checkbox"/>	CT Chest W Cont	T;N, Reason for Exam: Other, Enter in Comments, Order comments: For Aortic Stent Graft, 2.5 mm cuts - no contrast, then 2.5 mm cuts - arterial phase of contrast, then 2.5 mm cuts, venous phase. NO Oral contrast.

Date

Time

Physician's Signature

MD Number

