

IDENTIFICATION OF PUTATIVE VIRULENCE FACTORS FROM *CAMPYLOBACTER* spp.  
ISOLATED IN ICELAND

By

EDITH DEANN AKINS

(Under the Direction of Mark Harrison)

ABSTRACT

Two studies were conducted to determine putative virulence factors of *Campylobacter* spp. that were isolated in Iceland. The first study investigated capacitance monitoring using a simplified medium for efficient and reproducible construction of growth curves for *Campylobacter* spp., which can be a time consuming and labor intensive process. When invasion assays are performed, it is required that *Campylobacter* spp. isolates be grown to a density of  $10^6$  to  $10^8$  CFU/ml. This investigation optimized conditions for use with the Bactometer® such that the determination of growth curves was achieved in a simple medium. Results suggested that isolates should be grown on Mueller Hinton plates under a microaerobic atmosphere (37°C; 24 h), then transferred to Mueller Hinton biphasic cultures for 6 h (37°C; microaerobic atmosphere). Serial dilutions should be used for inoculation of Bactometer® wells containing 1 mL Mueller Hinton broth plus 0.1M sodium pyruvate for obtaining growth curves. In the second study, putative virulence factors of *Campylobacter* spp. were investigated. *Campylobacter* spp. exhibited a wide distribution of adhesion and invasion ability, which was determined to be unrelated to *flaA* short variable region allele type. The second part of this study investigated the most invasive isolate 14118, the least invasive isolate, 13262, and two in

between to further understanding of the molecular basis of genetic diversity among these 4 *C. jejuni* isolates. DNA-DNA microarray hybridizations identified genes absent relative to *C. jejuni* 11168 (PMSRU). Several absent genes were located in 1 of 7 previously described plasticity regions. There were 372 genes determined to be present in *C. jejuni* isolates 14118, 5116, 8557 and 13262 as well as *C. jejuni* 11168 (PMSRU). DNA suppressive subtractive hybridizations identified genes not in common with *C. jejuni* 11168 (PMSRU). *C. jejuni* 14118 contained a gene from *C. doylei* 269.97 that encoded for a motility accessory factor. *C. jejuni* 13262 contained a cytolethal distending toxin operon from *C. lari*. as well as a type II restriction modification enzyme unlike isolate 14118, 5116 and 8557 which includes a type I restriction modification enzyme.

INDEX WORDS: *Campylobacter*, virulence, capacitance, Bactometer, cell invasion, microarray hybridization, suppressive subtractive hybridization

IDENTIFICATION OF PUTATIVE VIRULENCE FACTORS FROM *CAMPYLOBACTER* spp.  
ISOLATED IN ICELAND

By

EDITH DEANN AKINS

B.S.A., University of Georgia, 2002

M.S., University of Georgia, 2005

A Dissertation Submitted to the Graduate Faculty of The University of Georgia in Partial  
Fulfillment of the Requirements for the Degree

DOCTOR OF PHILOSOPHY

ATHENS, GEORGIA

2008

© 2008

Edith Deann Akins

All Rights Reserved

IDENTIFICATION OF PUTATIVE VIRULENCE FACTORS FROM *CAMPYLOBACTER* spp.  
ISOLATED IN ICELAND

By

EDITH DEANN AKINS

Major Professor: Mark Harrison

Committee: Kelli Hiatt  
Joseph Frank  
Mark Berrang  
Jinru Chen

Electronic Version Approved:

Maureen Grasso  
Dean of the Graduate School  
The University of Georgia  
August 2008

## DEDICATION

This dissertation is dedicated to

My family

For their prayers, constant support, and unconditional love

## ACKNOWLEDGEMENTS

I would like to thank my major professor, Dr. Mark Harrison and co-advisor Dr. Kelli Hiett for their guidance, constant encouragement, and financial support throughout my Ph.D. program. They both have taught me so much about science but more importantly about life. I would also like to thank my committee members: Dr. Joseph Frank, Dr. Mark Berrang and Dr. Jinru Chen for their knowledge and willingness to help me in this program. Lastly I would like to thank all my friends and co-workers for their help, support, and encouragement.

## TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS.....	v
LIST OF TABLES.....	vii
LIST OF FIGURES.....	ix
CHAPTER	
1 INTRODUCTION.....	1
2 LITERATURE REVIEW.....	6
3 SIMPLIFIED CAPACITANCE MONITORING FOR THE DETERMINATION OF <i>CAMPYLOBACTER</i> spp. GROWTH RATES.....	28
4 IDENTIFICATION OF PUTATIVE VIRULENCE FACTORS UTILIZING INVASION ASSAYS AND WHOLE GENOME COMPARISONS OF <i>CAMPYLOBACTER</i> spp. ISOLATES RECOVERED FROM ICELAND.....	38
5 SUMMARY AND CONCLUSIONS.....	119



## LIST OF TABLES

	Page
Table 3.1: Time required for different strains of <i>Campylobacter</i> spp. to reach a detection limit of approximately 10 <sup>6</sup> CFU/mL by capacitance measurement using the Bactometer.....	35
Table 3.2: Experimental <i>Campylobacter</i> spp. growth conditions evaluated for detection of changes in capacitance using the Bactometer®.....	36
Table 4.1: <i>flagellinA</i> short variable region (SVR) allele, sample number, and sample origin of <i>Campylobacter</i> spp. isolates employed in this investigation.....	63
Table 4.2: Genes identified as absent from <i>C. jejuni</i> isolate 14118 (high invasion) based on microarray hybridization analysis with <i>C. jejuni</i> 11168 (PMSRU).....	65
Table 4.3: Genes identified as absent from <i>C. jejuni</i> isolate 5116 (mid invasion) based on microarray hybridization analysis with <i>C. jejuni</i> 11168 (PMSRU).....	66
Table 4.4: Genes identified as absent from <i>C. jejuni</i> isolate 8557 (mid invasion) based on microarray hybridization analysis with <i>C. jejuni</i> 11168 (PMSRU).....	67
Table 4.5: Genes identified as absent from <i>C. jejuni</i> isolate 13262 (low invasion) based on microarray hybridization analysis with <i>C. jejuni</i> 11168 (PMSRU).....	68
Table 4.6: Genes determined to be commonly distributed throughout <i>C. jejuni</i> isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with <i>C. jejuni</i> 11168 (PMSRU).....	70
Table 4.7: Total number of inserts, and clones provided for sequence analysis along with percentage of clones with sequences absent from <i>C. jejuni</i> 11168 (PMSRU) based on suppressive subtractive hybridizations.....	85
Table 4.8: Isolate 14118 (Caco-2 high invasion) unique clones relative to <i>C. jejuni</i> 11168 (PMSRU).....	86

Table 4.9: Unique clones, recovered from suppressive subtractive hybridizations, of <i>C. jejuni</i> isolate 14118 (Caco-2 high invasion) determined to possess significant similarity to <i>Campylobacter</i> spp. other than <i>C. jejuni</i> 11168 (PMSRU).....	92
Table 4.10: Isolate 5116 (Caco-2 mid invasion) unique clones relative to <i>C. jejuni</i> 11168 (PMSRU).....	93
Table 4.11: Unique clones, recovered from suppressive subtractive hybridizations of <i>C. jejuni</i> isolate 5116 (Caco-2 mid invasion) determined to possess significant similarity to <i>Campylobacter</i> spp. other than <i>C. jejuni</i> 11168 (PMSRU). ....	96
Table 4.12: Isolate 8557 (Caco-2 mid invasion) unique clones relative to <i>C. jejuni</i> 11168 (PMSRU).....	97
Table 4.13: Unique clones, recovered from suppressive subtractive hybridizations, of <i>C. jejuni</i> isolate 8557 (Caco-2 mid invasion) determined to possess significant similarity to <i>Campylobacter</i> spp. other than <i>C. jejuni</i> 11168 (PMSRU). ....	105
Table 4.14: Isolate 13262 (Caco-2 low invasion) unique clones relative to <i>C. jejuni</i> 11168 (PMSRU). ....	106
Table 4.15: Unique clones, recovered from suppressive subtractive hybridizations, of <i>C. jejuni</i> isolate 13262 (Caco-2 low invasion) determined to possess significant similarity to <i>Campylobacter</i> spp. other than <i>C. jejuni</i> 11168 (PMSRU) .....	112

## LIST OF FIGURES

	PAGE
FIGURE 3.1: Typical mean capacitance response for <i>Campylobacter</i> spp. 4568 inoculated into Bactometer® wells at 10 <sup>1</sup> CFU/mL.....	37
FIGURE 4.1: Dendrogram of representative <i>Campylobacter</i> spp. <i>flaA</i> SVR alleles recovered during the Icelandic Epidemiology Investigation.....	113
FIGURE 4.2: Distribution of <i>Campylobacter</i> spp. <i>flaA</i> SVR allele groups by percent adhesion to Caco-2 Cells. ....	114
FIGURE 4.3: Figure 4.3. Distribution of <i>Campylobacter</i> spp. percent adhesion of Caco-2 Cells relative to original host. ....	115
FIGURE 4.4: Distribution of <i>Campylobacter</i> spp. <i>flaA</i> SVR allele groups by percent invasion of Caco-2 Cells. ....	116
FIGURE 4.5: Distribution of <i>Campylobacter</i> spp. host type by percent invasion of Caco-2 Cells. ....	117
FIGURE 4.6: Functional categories of clones recovered using suppressive subtractive hybridization on all 4 <i>C. jejuni</i> isolates. ....	118

## CHAPTER 1

### INTRODUCTION

*Campylobacter* spp. are curved rod, gram-negative bacteria with polar flagella (2).

Campylobacteriosis is one of the most common bacterial intestinal disorders of humans in many industrialized countries (5). Human campylobacteriosis generally presents itself in 3-5 days after exposure with watery or bloody diarrhea, abdominal pain and nausea. Although normally self-limiting, other systemic infections can occur such as Guillain-Barre Syndrome and Reiter's Syndrome (6).

The consumption of poultry is considered the most likely route of infection. Broilers frequently carry large numbers of *Campylobacter* spp. in their intestinal contents. This carriage is asymptomatic and spillage of gut contents during processing can contaminate the retail poultry products and the abattoir environment (1). In the United States, retail chicken carcasses have estimated contamination rates of 60-80% with *Campylobacter* spp. populations averaging  $10^6$  for fresh chicken and  $10^4$  for frozen carcasses (3).

Reducing and eliminating foodborne human pathogens associated with poultry has received attention in the last years. The number of poultry being commercially processed grew from approximately 7.3 to 8.4 billion from 1994-2001, a 15% increase. Due to the increase of production and consumption of poultry, the national goal in the U.S, set by governmental agencies, is to reduce the incidence of *Campylobacter* infection from 13.37 per 100,000 people in 2002 to 12.30 per 100,000 people in 2010 (4). The Center for Disease Control and Prevention (CDC) reported that *Campylobacter* infection decreased 31% between 1996-1998. Since this decrease the estimated incidence of *Campylobacter* has not changed significantly. In 2007 the number of *Campylobacter* cases and incidence per 100,000 population was 12.79 (4). Since the Healthy People 2010 national health target has not been reached, there needs to be improved

understanding of the transmission of *Campylobacter* spp. from potential sources to human illness.

This dissertation used *Campylobacter* spp. isolates recovered during a comprehensive epidemiologic investigation conducted in Iceland. The original objectives were to investigate the ultimate sources and risk factors for transmission of *Campylobacter* spp. broiler flocks and their relevance to human exposure, utilizing multi-disciplinary international expertise. The study centered on the sampling of an entire broiler production population, which provides the sole source of broiler meat to the human population in Iceland. The multidisciplinary analysis of the results provide a perspective to describe sources and risk factors for exposure and infection of broilers, which demonstrate the greatest potential for reducing *Campylobacter* prevalence and concentration in poultry. It has also directed research efforts toward intervention technologies for reducing *Campylobacter* exposure in poultry and humans.

Iceland was targeted for this study due to its comparability to North American broiler production. All broiler chicken production in Iceland are derived from hatching eggs imported from grandparent breeder flocks in Sweden. Parent breeder chicks are distributed to three vertically integrated production lines, each with its own broiler hatchery. Based on visits and reviews of all on farm production, slaughter and processing, similarities to North American production are more remarkable than the differences. Poultry house construction though is more similar to Canadian poultry houses while the equipment and technology is consistent with both the U.S. and Canada. Production in Iceland is on a smaller scale with flock sizes ranging from <1,000 to 13,500. The limited scale of production and the fact that no broiler meat products are imported into Iceland, enabled a total population based epidemiological study. To compliment the closed system, molecular typing was employed and access to all human disease isolates and

available human case data was given. It would not be feasible for such an all inclusive population study in the U.S. due to scale of production, complex market distribution and disperse consuming population present numerous confounding factors, which can limit the inferential value of epidemiologic studies.

The purpose of this research is to better understand the molecular basis and biological consequences of genetic diversity among *Campylobacter* spp. for disease control.

*Campylobacter* spp. isolates, recovered from poultry only and from poultry and humans, were surveyed to determine if “markers” might exist to allow us to determine if a *Campylobacter* isolate is possibly more likely to infect or cause disease in humans. The results from these experiments could lead to further investigations which may aid in better understanding of *Campylobacter* virulence and potentially lead to developing mitigation strategies to reduce the amount of human campylobacteriosis.

## References

1. Ahmed, I. F., G. Manning, T. M. Wassenaar, S. Cawthraw, and D. G. Newell. 2002. Identification of genetic differences between two *Campylobacter jejuni* strains with different colonization potentials. *Microbiol.* 148:1203-1212.
2. Altekruze, S. F., and D. L. Swerdlow. 2002. *Campylobacter jejuni* and related organisms, p. 103-112. In D. O. Cliver and H. P. Riemann (ed.), *Foodborne Diseases*, 2 ed. Academic Press, San Diego.
3. Altekruze, S. F., D. L. Swerdlow, and N. J. Stern. 1998. Microbial food borne pathogens. *Campylobacter jejuni*. *Vet Clin. Am. Food. Anim. Pract.* 14:31-40.
4. CDC. 2004, posting date. Healthy People 2010. [www.healthypeople.gov](http://www.healthypeople.gov). Accessed Sept. 1, 2006.
5. Knudsen, K. N., D. D. Bang, L. O. Andresen, and M. Madsen. 2006. *Campylobacter jejuni* strains of human and chicken origin are invasive in chickens after oral challenge. *Avian Dis.* 50:10-14.
6. Lee, M. D., and D. G. Newell. 2006. *Campylobacter* in Poultry: Filling an Ecological Niche. *Avian Dis.* 50:1-9.



CHAPTER 2  
LITERATURE REVIEW

## ***Campylobacter* History and Characteristics**

In 1886, Theodore Escherich observed stool samples of children with diarrhea that had organisms he believed resembled *Campylobacter*. In 1913, McFaydean and Stockman identified campylobacters, at the time called related *Vibrio*, in fetal tissues of aborted sheep. In 1957, King described *Campylobacter* from blood samples of children with diarrhea, and in 1972, Belgian clinical microbiologists first isolated *Campylobacter* from stool samples from patients with diarrhea. Due to the development in the 1970s of selective growth media more laboratories were able to test stool specimens for *Campylobacter* spp. Soon, *Campylobacter* spp. were established as common human pathogens (6) and are recognized as the leading cause of bacterial foodborne diarrheal disease throughout the developed world (46).

## **The Family *Campylobacteraceae***

This family is comprised of gram negative, nonsaccharolytic bacteria with microaerobic growth requirements (3-5% carbon dioxide, 3-15% oxygen) (8, 62) and have a small genome of approximately 1.6-1.7 Mbp of AT-rich DNA; the GC ratio is approximately 30% (35). Members of the family occur primarily as commensals or parasites in humans and domestic animals. Cells in the family *Campylobacteraceae* are curved, S-shaped, or spiral rods that are 0.2 to 0.8  $\mu\text{m}$  wide, 0.5 to 5  $\mu\text{m}$  long and non-sporeforming (48). Cells in old cultures may form spherical or coccoid bodies. They are typically motile with a characteristic corkscrew motion by means of a single polar unsheathed flagellum at one or both ends of the cell. Cells grow under microaerobic conditions (42) and have a respiratory and chemoorganotrophic type of metabolism. However, some species of *Campylobacter*, such as *C. sputorum*, *C. concisus*, *C. mucosalis*, *C. curvus*, *C. rectus*, and *C. hyointestinalis* require an atmosphere containing an increased concentration of hydrogen to be isolated (42). Enzymes such as superoxide dismutase (SOD), catalase,

peroxidase, glutathione synthetase, and glutathione reductase in *C. jejuni* are believed to play an important role in providing oxygen toxicity protection. *Campylobacter* spp. are fastidious organisms that require complex growth media (61). Energy is obtained from amino acids or tricarboxylic acid cycle intermediates, not carbohydrates (62).

*Campylobacter* spp. are referred to as thermophilic *Campylobacters* and grow best at 37 to 42°C, with an optimal temperature of 42°C, which reflects the adaptation to the intestines of birds (29, 64). Typical biochemical characteristics are reduction of fumarate to succinate; negative methyl red reaction and acetoin and indole production; and for most species, reduction of nitrate, absence of hippurate hydrolysis, and presence of oxidase activity (62).

### **Clinical Features**

*Campylobacter* enteritis is considered to be a foodborne disease, with infection often being derived from a range of foods and also water-based environmental sources (35). It has been reported that every year in the United States there are about 20-150 cases per 100,000 people. There is a high isolation rate among young adults (15-24 years old), approximately 8 per 100,000 per year. Under-reporting is significant and the true incidence could be five to ten times higher (38). The vast majority of cases appear to be sporadic; unlike with *Salmonella* and *Escherichia coli*, outbreaks of campylobacteriosis are rarely observed (6). Sporadic illnesses peak during summer months and are associated with mishandling or consumption of undercooked poultry or cross- contamination of other foods by raw poultry (7). Outbreaks of *Campylobacter* spp. have resulted from cross-contaminated chicken, raw milk, and untreated water (6, 41).

Under certain conditions, for example on exposure to atmospheric oxygen, bacteria can become spherical or coccoid in shape. This change in shape has been associated from a viable

culturable form to a viable but not cultural form (VNC) and was first proposed by Colwell et al. following a study on survival of *Salmonella* in aquatic systems (53). It has been suggested that this VNC state acts as an adaptation for survival in adverse environments such as low nutrient availability or upon entry into stationary phase (41). Rollins and Colwell were the first to report a VNC form of *C. jejuni* and since then there has been debate as to whether the form for *Campylobacter* truly exists (52). There has been evidence that shows that VNC *Campylobacter* spp. are infectious in neonatal mice but in a chicken model the evidence is more contradictory. Such investigations are difficult to perform and interpret because not all coccoid cells may progress to a VNC state or VNC development may advance through several stages in a coccoid cell (35).

Human campylobacteriosis generally presents as 3-5 days of acute watery or bloody diarrhea, usually with severe abdominal pain, fever and general malaise (15). Most cases of infection are due to *C. jejuni*, with only 10% due to *C. coli* and less than 1% *C. lari* (38). In a volunteer study, *C. jejuni* infection occurred after ingestion of as few as 800 organisms (6). In another volunteer study, inoculations with large doses ( $10^5$  to  $10^8$  colony forming units) were required for infection. Most infections are due to exposure to relatively low numbers of organisms that must multiply in the host to achieve a clinically apparent outcome. Examination of colonic biopsies shows an acute inflammatory response with infiltration of the epithelium and lamina propria with neutrophils and mononuclear cells. Among infected persons in developed countries, both leukocytes and erythrocytes are nearly always present in stools, indicating the universality of the inflammatory process, even when stools are watery and not grossly bloody. Thus, *Campylobacter* colitis or enteritis must be considered an inflammatory disease (65).

Although campylobacteriosis is generally self-limiting, incapacity may last several weeks and up to 10% of reported cases may require medical intervention.

More serious systemic infections are well recognized and occur most commonly in the immunocompromised. Chronic sequelae, like arthropathies, are not uncommon and an association of campylobacteriosis with postinfectious neuropathies such as Guillain-Barré syndrome have been identified (38). Guillain-Barré syndrome is an acute inflammatory demyelinating neuropathy that occurs from a cross reaction with Schwann-cells or myelin. It results in neuromuscular paralysis. An estimated one case of GBS occurs for every 1,000 cases of campylobacteriosis (5). *Campylobacter* spp. are also associated with Reiter syndrome, a reactive arthropathy that affects the ankles, knees, wrists, and the small joints of the hands and feet. The duration of arthritis ranges from several weeks to several months. Although the arthritis can be incapacitating, full recovery is generally seen. The pathogenesis of GBS and Reiter syndrome is not completely understood (6).

Treatment for campylobacteriosis usually involves rehydration, but antimicrobial therapy may be required for patients who have high fever, bloody diarrhea, or more than 8 stools in 24 h; immunosuppressed patients, patients with bloodstream infections, and those whose symptoms worsen or persist for more than 1 week from the time of diagnosis (6). Erythromycin is the drug of choice, but ciprofloxacin is now frequently used in adults (38) and fluoroquinolones and tetracyclines are used as alternatives (15).

### **Pathogenesis**

Many pathogen-specific virulence determinants may contribute to the pathogenesis of *C. jejuni* infection in humans, but none has a proven role. Potential determinants of pathogenicity include chemotaxis, motility, and flagella, which are required for attachment and colonization of

the gut epithelium (6). *Campylobacter* enters the host intestine by way of the stomach acid barrier and colonize the distal ileum and colon. Following colonization of the mucus and adhesion to intestinal cell surfaces, *Campylobacter* perturbs the normal absorptive capacity of the intestine by damaging epithelial cell function either directly, by cell invasion or the production of toxin(s), or indirectly following the initiation of an inflammatory response. These possibilities are not mutually exclusive; any combination may have a role depending on the host status and attributes of the infecting strain (35).

The ability of *C. jejuni* to colonize the human gastrointestinal tract is essential for disease. Binding to epithelial cells prevents the colonizing bacteria from being swept away by mechanical cleansing forces such as peristalsis and fluid flow. Binding is prerequisite for entry into host cells. The ability of *C. jejuni* to invade cultured cells appears to be strain dependent (36). Recent advances include the identification and characterization of virulence factors required for *C. jejuni* binding, entry, and survival within host cells.

### **Chemotaxis and Motility**

Effective colonization requires chemotaxis. *Campylobacter* spp. have mechanisms to detect chemical gradients and linked motility functions that enable the cell to move up or down the gradient. Non-chemotactic mutants failed to colonize the suckling mouse intestine. Studies have revealed various chemoattractants including mucin, L-serine and L-fucose, while several bile acids have chemorepellant effects. Motility of *Campylobacter* spp. necessitates the production of the flagellum which is an important virulence factor (35). Functional flagella presumably help the bacteria overcome the clearing movement of peristalsis and enable them to enter and cross the mucus layer overlaying the epithelium (65). They are able to move through viscous environments at speeds up to 75  $\mu\text{m/s}$  (18). *C. jejuni* contains one or two polar flagella

that cause the typical darting motility observed by microscopy and the moist appearance of colonies on agar plates. The flagellar filament consists of multimers of the protein flagellin and is attached by the hook protein to a basal structure, which is embedded in the membrane and serves as a motor for rotation (65). The flagella are complex and are composed of two related subunits, FlaA and FlaB (37). They are >93% homologous to each other and are encoded by genes which are adjacent on the chromosome but regulated by distinct promoters. The *flaA* gene is regulated by the flagellar promoter  $\sigma^{28}$  and the *flaB* gene is regulated by a  $\sigma^{54}$  promoter (27). Early studies with genetically undefined mutants indicated that the flagellum was needed for adhesion and for colonization in a range of animals (35). Mutation of *flaA* resulted in the synthesis of a truncated flagellar filament, composed of *flaB*. This flagellar filament greatly reduced motility compared to that of the wild type flagellum. Mutation of *flaB* resulted in a flagellar filament which was indistinguishable from the wild type filament in length and with motility that was reduced slightly compared to the wild type (27).

### **Adhesion and Invasion**

*C. jejuni* is increasingly associated with episodes of gastrointestinal illness in developed countries. Although little is known about the pathogenic events required for illness, *C. jejuni* strains differ in their ability to adhere to and invade intestinal epithelial cells and it has been suggested that this is important to the pathogenic process (51). Grant et al. found that flagella are not involved in *C. jejuni* adherence to epithelial cells but that they do play a role in internalization (25). Another study by Konkel et al. showed that not only was either the FlaA or FlaB gene required for invasion but that *Campylobacter* invasion antigens (Cia proteins) were also needed for maximal invasion. This data also showed that the *C. jejuni* Cia proteins are secreted from the flagellar export apparatus (37). Hanel et al. showed that there was correlation

between invasion of Caco-2 cells (human colon cells) and colonization in the chick gut. Isolates that did not colonize the chick gut were not invasive with the Caco-2 cells. Strains that showed to be strong colonizers had high invasion rates with the Caco-2 cells (28). Monteville et al. (40) published that the CadF outer membrane protein appears to promote the binding of *C. jejuni* to fibronectin, thereby stimulating the host cell signaling events associated with bacterial uptake. Fibronectin is a glycoprotein that is present at regions of cell to cell contact in the gastrointestinal epithelium, providing a potential binding site for pathogens (40). Zheng et al. (68) and Datta et al. (22) reported for an isolate to be invasive it needs the *flaA* gene, *cad*, and other genes such as *ciaB*, *pldA*. Genes of the pVir plasmid are also involved in host cell invasion. The genes *cdtA*, *cdtB*, and *cdtC* are responsible for the expression of *Campylobacter* cytolethal distending toxin, which induces the proinflammatory cytokine production of epithelial cells and causes host cell cycle arrest, cell distention and eventually cell death (68).

### **Poultry Production**

In humans, *Campylobacter* infections are primarily the result of the ingestion of contaminated foods of animal origin such as poultry (46, 55). In poultry, *Campylobacter* is a commensal organism and colonizes the mucus overlying the epithelial cells primarily in the caeca and small intestine, but may be recovered from other places in the gastrointestinal tract and from the liver and spleen.

*Campylobacter* contamination of poultry flocks, horizontal transmission and/or vertical transmission, remain unclear. Horizontal sources that could be vectors of infection include environment of the poultry house, hatchery pads, litter, feed, water, personnel, small animals on the farm, flies and rodents (21, 34, 55). However, none of these suspected sources has been identified conclusively as the formal source of infection on broiler farms. In many cases there is



no comparison of isolates from broilers and the environment by phenotypic and genotypic typing methods, leading to significance of these putative sources of infection. *C. jejuni* was probably detected in suspect sources after the broilers had come infected, suggesting that broilers, instead of being infected from environmental sources, might be the source of environmental contamination. It is very difficult, in many situations, to determine which contamination came first (55).

*C. jejuni* is very sensitive to oxygen and drying so the organism is generally unable to grow in feed, litter or water under normal ambient conditions (32). *Campylobacter* has been isolated from water lines and reservoirs of bird houses, and these strains may be phenotypically and genotypically identical to what is found in feces. However the water contamination usually follows rather than precedes the colonization of a flock (44).

Flies, mice and other pests can act as a vector for the transmission of *C. jejuni* (31). Identical serotypes and genotypes have been isolated from both broilers and insects within broiler houses; however the direction of spread has not been determined (55). Stern et al. (58) found 25% of insects caught outside of poultry houses to be *Campylobacter* positive four weeks prior to detection of *Campylobacter* in flocks. The bacteria though, can only survive on or within these insects for a few days (44).

*Campylobacter* has also been shown to be present and possibly transferred by workers as well as equipment. The organism has been isolated from footbath water, farmer's boots and transport crates (58). Therefore it is reasonable to believe transmission of *C. jejuni* may be spread between flocks and houses by personnel. Nesbit et al. (43) showed two adjacent broiler houses that lacked biosecurity measures were colonized with different genotypes of *C. jejuni* even though the two houses shared equipment and the same farmer worked both houses.

Vertical transmission has been debated as a theory on the mode of transmission. Sahin et al. were not able to recover *Campylobacter* from eggs that had been inoculated with the bacterium. They sampled for the presence of *C. jejuni* in eggs placed in an incubator to hatch and in newly hatched chicks (54). In another study though, Van de Giessen and others have shown vertical transmission as a means of contamination of a breeder flock. *C. jejuni* isolates from a parent flock were found to be from the same clonal origin as those from the offspring in the broiler flock (34). *Campylobacter* has also been isolated from various segments of the reproductive tract of the chicken (16). *C. jejuni* can be recovered from the oviduct, which suggests a possibility of egg contamination and it also has been found in semen samples from breeder cockerels (31).

### **Poultry Processing**

When a flock of broiler chickens becomes positive for *Campylobacter*, the prevalence of infection is high, often reaching 100% of the birds tested (26). The high number of *Campylobacter* positive birds can still be found in the first stages of processing. In a poultry processing plant, there are typically six basic functions: pre-scalding, scalding, defeathering, evisceration, washing, and chilling. Poultry are eviscerated, and as the skin is not normally removed, many contaminants are found on the skin (20). Son et al. (57) reported that *Campylobacter* was isolated from 78.5% of the carcasses sampled from three sample sites (pre-scald, pre-chill, and post-chill). The pre-scald and pre-chilled sites had the highest *Campylobacter* contamination at 92% and 100%, respectively (57). Berrang and Dickens (12) found that *Campylobacter* was the highest when sampled pre-scald ( $4.7 \log_{10}$ ) and the counts dropped significantly after the carcasses were scalded ( $1.8 \log_{10}$ ). When the carcasses exited the chill tank, the incidence of *Campylobacter* was almost the same as what was recorded post-scald.

The potential for cross-contamination during scalding could occur due to the follicles remaining open throughout the processing until the carcass is chilled. Once the follicles close during chilling, the organisms may become trapped (60). Contamination levels could be directly related to bacterial levels in the processing water. Water samples from both bird washers and chlorinated chiller water have been found *Campylobacter* positive (10).

Several mitigation steps have been incorporated in poultry processing to help control *Campylobacter* contamination. Chlorine has been used for more than 40 years in poultry processing to reduce spoilage bacteria, control the spread of pathogens, and prevent buildup of microorganisms on surfaces and equipment (34). Berrang et al. (11) reported that application of chlorine in the chill tank was significantly related to a larger reduction in *Campylobacter* numbers ( $P=0.0003$ ). However, the difference with the overall reduction was small. Oyarzabal et al. (45) found that the post-chill application of acidified sodium chlorite to chicken carcasses caused a significant reduction in *Campylobacter* numbers. Rapid freezing of carcasses offers additional control measures. An Icelandic study suggested that frozen poultry poses a lower risk to health than fresh meat (59). A method used in European countries that will kill *Campylobacter* and other infectious bacteria is the use of irradiation using electron beams or high energy electromagnetic radiation. Studies have shown that *Campylobacter* are more susceptible to radiation than *Salmonella* and *Listeria monocytogenes* (31).

Other mitigation steps focus on preventing contamination from fecal or cecal sources. One study discovered the odds of having a positive skin sample were 35 times greater when the same carcasses had *Campylobacter* positive intestines (33). Berrang et al. (13) tested the effect of placing vinegar in the cloaca prior to defeathering, which is a processing step that has been

known to increase *Campylobacter* spp. levels. They found the use of vinegar resulted in a significantly lesser increase of *Campylobacter* levels.

Despite mitigation steps, as much as 98% (17) of retail broilers have been reported as being positive for *Campylobacter* spp. and levels of contamination may vary between  $10^2$  and  $10^5$  CFU per carcass (46). It is likely that many consumers purchase poultry products that are contaminated. How consumers handle products at home will ultimately affect the cross-contamination and infection rate.

### **Microarray Technology and Suppression Subtractive Hybridization**

A wealth of nucleotide sequence data from the genomes of bacterial pathogens has become readily available. Comparative analysis of bacterial genomes provides information on the physiology and evolution of bacteria and allows for detailed comparisons between related bacteria. The availability of whole genome sequences has led to the development of microarray and suppression subtractive hybridization technology. These two methods provide researchers tools to continue the investigation into why some bacterial strains are more virulent than others.

#### Microarrays

Microarray technology has rapidly advanced and gained in popularity over the years. A microarray comprises a large number of genes deposited onto a glass slide, which are used for a multiplex reaction, essentially a large dot blot (66). The identification of genes present in a cellular sample as well as genes expressed by hybridizing cDNA from reverse transcribed RNA can be determined through microarrays. It should be noted, however, that microarrays are limited to pre-determined genes of interest, which are printed on a slide. Also, some genes may be expressed and are regulated at the translational or post-translational level, so this type of expression would not be detected by microarrays.

Microarrays began in the mid-1990s with the use of glass microscope slides for printing DNA (56). Glass is a good choice as a microarray substrate due to its low fluorescence, low cost, high heat resistance, and rigidity. Slides must be cleaned and coated for DNA immobilization/spotting to occur on glass (1). Glass can be coated either adsorptively, as with poly-l-lysine, or covalently, as with functionalized silanes. Functionalized silanes are most commonly used since they can bind directly to nucleic acids (56). The polyanionic DNA interacts with the polycationic surface by way of coulombic attraction. After printing of the slide, the DNA is locked to the surface by ultraviolet irradiation or baking. Although non-covalent mechanisms are typically used, covalent binding has advantages. The DNA can be oriented in a certain way to increase probe availability for hybridization, allow less non-specific binding, allow for the use of more stringent washes, and offers a potential for stripping and re-hybridizing arrays (9). An example of covalent binding is the use of amino modified DNA to bind to epoxide activated glass slides. Amino alcohols form when the epoxides react through nucleophilic displacement (1).

Fluorescently labeled cDNA is hybridized to slides using either direct or indirect labeling. Direct labeling of cDNA is accomplished by a reverse transcriptase reaction with mRNA as a template, dye conjugated nucleotides, and oligo (59)n as primers (14). Indirect labeling incorporates amino-allyl dUTP into the cDNA and the primary amine groups are subsequently conjugated to succinimidyl ester of Cy3. This labeling method increases labeling density, however, it is also known to decrease the intensity of fluorescent signal (49).

Microarray experiments must be replicated since chips can fail or be distorted. If a study is to find large differences, then a design with 3 samples per group is adequate (50). After data is

collected, it must be normalized to remove technical differences. Local regression is typically used for normalizing microarrays (24).

#### Microarray Experiments on *Campylobacter*

Pearson et al. (47) investigated diversity in 18 *C. jejuni* strains from diverse sources using microarrays. In total, 16.3% of the genes present in the sequenced strain NCTC11168 were either absent or highly variable in sequence among the strains of *C. jejuni* examined in this study. Seven major plasticity regions (PR) were also identified in the genome and they comprised 50% of the variable gene pool. PR 1 contained genes important in the utilization of alternative electron acceptors for respiration and may confer a selective advantage to strains in restricted oxygen environments. PR 2, 3, and 7 contain many outer membrane and periplasmic proteins and hypothetical ones of unknown function. PR 4, 5, and 6 contain genes involved in the production of surface structures including LOS, flagellum, and post-translational glycosylation of the flagellum. These variable regions identified in this study highlight genetic factors that might be linked to phenotypic variation and adaptation to different ecological niches. Another study used oligonucleotide microarrays for a rapid and accurate simultaneous differentiation among *C. jejuni*, *C. coli*, *C. lari*, and *C. upsaliensis*. The array contained species-specific oligonucleotide probes developed by using specific regions of five genes (*fur*, *glyA*, *ceuB-C*, and *fliY*) (63). Champion et al. (19) used microarrays to examine 111 *C. jejuni* strains isolated from humans, chickens, bovine, ovine, and the environment. NCTC 11168 was used as the control in the experiment. The *C. jejuni* functional core was determined by calculating the number of genes that were present in each of the test strains as well as the control strain. The species-specific functional core consisted of 979 predicted coding sequences comprising 59.2% of the genome and was involved with regulatory, metabolic, cellular, and biosynthetic processes.

DNA microarray analysis has also been used to determine if isolates from patients with Guillain-Barré syndrome (GBS) differ from isolates from patients with uncomplicated gastrointestinal infection (39). Microarray analysis did not identify discrete groups of isolates or any unique features within the genome of the *C. jejuni* isolates associated with GBS. It could not be determined if the lack of hybridization in various regions represents the absence of a particular gene or nucleotide divergence within an existing gene. Additionally, differences due to the presence of genetic elements in either the GBS or enteritis related isolates would not necessarily be detected because of the absence of such elements in the genome of the strain used to construct the microarray.

#### Suppression subtractive hybridization

Suppression subtractive hybridization is a technique designed to identify those regions present in one genome but absent from another (2). This method was first reported in 1996 and applied to the study of *Helicobacter pylori* (4). Genomic DNA extracted from a driver strain is hybridized with DNA extracted from a tester strain with a view to isolating those sequences that are present in the tester strain but absent from the driver strain (23). To facilitate the process, the driver strain DNA must be present in excess. It is important to ensure that tester DNA sequences are short so a digestion by restriction enzymes before hybridization can occur. The tester DNA is then separated into two portions, each of which is subjected to a ligation reaction to attach a different adaptor sequence to the 5' ends. The two portions are then separately hybridized to the driver DNA (in excess). All sequences that hybridize with the driver DNA should be mopped up leaving only tester specific single-stranded sequences. When the two tester portions are mixed and hybridized together, only those sequences unique to the tester strain will have different adaptors present on each strand. PCR is then used to detect these sequences. Only the sequences

that carry both adaptors, one on each end, will amplify. The PCR products are cloned into a vector to produce a subtracted library. Although this procedure is not entirely effective, >50% of clones should be tester-specific (67).

#### Suppression subtraction hybridization experiments

Hepworth et al. (30) looked at the use of suppression subtractive hybridization to extend our knowledge of *C. jejuni* genome diversity. They carried out five subtractions between *C. jejuni* isolates from different sources such as rabbit, cattle and wild birds. They wanted to determine the variability within and between common multilocus sequence type (MLST) clonal complexes. The results of the study showed a correlation between clonal complex and the distribution of the metabolic genes. There was, however, no evidence to support the hypothesis that host preference may have any role in the distribution of such genes. Another study by Ahmed et al. (3) researched genetic differences between two *C. jejuni* strains with different colonization potentials. *C. jejuni* NCTC 11168 has been shown in preliminary studies to be a poor colonizer whereas strain 81116 is a better colonizer in chickens. The technique of subtractive hybridization was used to identify gene fragments of strain 81116 not present in strain 11168. Six clones were found with similarities to restriction-modification enzymes found in other bacteria. Two inserts had similarity to arsenic-resistant genes, and four others had similarities to cytochrome c oxidase III, dTDP-glucose 4,6-dehydratase and an abortive phage-resistance protein. Some of these genes may be involved in colonization potential.



## References

1. Aboytes, K., J. Humphreys, S. Reis, and B. Ward. 2003. Slide coating and DNA immobilization chemistries, p. 1-42. *In* E. Blalock (ed.), *A Beginner's Guide to Microarrays*. Kluwer Academic Publishers, New York.
2. Agron, P. G., M. Macht, L. Tadnedge, E. W. Skowronski, W. Miller, and G. L. Andersen. 2002. Use of subtractive hybridization for comprehensive surveys of prokaryotic genome differences. *FEMS Microbiol. Letters* 211:175-182.
3. Ahmed, I. F., G. Manning, T. M. Wassenaar, S. Cawthraw, and D. G. Newell. 2002. Identification of genetic differences between two *Campylobacter jejuni* strains with different colonization potentials. *Microbiol.* 148:1203-1212.
4. Akopyants, N. S., A. Dradkov, and L. Diatchenko. 1998. PCR-based subtractive hybridization and differences in gene content among strains of *Helicobacter pylori*. *Microbiol.* 95:13108-13113.
5. Allos, B. M. 1997. Association between *Campylobacter* infection and Guillain-Barre Syndrome. *J. Infect. Dis.* 176 (Suppl2):S125-128.
6. Altekruise, S. F., N. J. Stern, P. I. Fields, and D. L. Swerdlow. 1999. *Campylobacter jejuni*- An Emerging Foodborne Pathogen. *Emerging Infectious Diseases* 5.
7. Altekruise, S. F., and D. L. Swerdlow. 2002. *Campylobacter jejuni* and related organisms, p. 103-112. *In* D. O. Cliver and H. P. Riemann (ed.), *Foodborne Diseases*, 2 ed. Academic Press, San Diego.
8. Anonymous, posting date. *Campylobacter jejuni*. [www.cfsan.fda.gov/~mow/chap4.html](http://www.cfsan.fda.gov/~mow/chap4.html). Accessed 16 Sept. 2006.
9. Beier, M. a. J. D. H. 1999. Versatile derivatisation of solid support media for covalent bonding on DNA-microchips. *Nucl. Acids Res.* 27:1970-1977.
10. Berndston, E., M. L. Danielsson-Tham, and A. Engvall. 1996. *Campylobacter* incidence on a broiler farm and the spread of *campylobacter* during the slaughter process. *Int. J. Food Microbiol.* 32:35-47.
11. Berrang, M. E., J. S. Bailey, S. F. Altekruise, B. Patel, W. K. Shaw, R. J. Meinersmann, and P. J. Fedorka-Cray. 2007. Prevalence and numbers of *Campylobacter* on broiler carcasses collected at rehang and postchill in 20 U.S. Processing Plants. *J. Food Prot.* 70:1556-1560.
12. Berrang, M. E., and J. A. Dickens. 2000. Presence and level of *Campylobacter* spp. on broiler carcasses throughout the processing plant. *J. App. Poult. Res.* 9:43-47.

13. Berrang, M. E., D. P. Smith, and A. Hinton. 2006. Application of distilled white vinegar in the cloaca to counter the increase in *Campylobacter* numbers on broiler skin during feather removal. *J. Food Prot.* 69:425-427.
14. Bittner, M., R. Butow, J. deRisi, M. Diehn, J. Eberwine, C. B. Epstein, R. Glynne, S. Grimmond, T. Ideker, J. E. Kacharina, S. Katsabanis, J. Khan, J. Lee, C. L. Liu, P. Marciano, E. M. Marincola, T. McIntosh, D. Monte, J. R. Pollack, V. Rhodius, S. Somerville, E. Tom, E. Wang, J. S. Wei, D. Willhite, and S. Ybarra. 2003. Expression analysis of RNA, p. 101-289. *In* D. Bowtel and J. Sambrook (ed.), *DNA Microarrays: a molecular cloning manual*. Cold Spring Harbor, Cold Spring Harbor.
15. Blaser, M. J. 1997. Epidemiologic and clinical features of *Campylobacter jejuni* infections. *J. Infect. Dis.* 176:S103-105.
16. Buhr, R. J., N. A. Cox, N. J. Stern, M. T. Musgrove, J. L. Wilson, and K. L. Hiatt. 2002. Recovery of *Campylobacter* from segments of the reproductive tract of broiler breeder hens. *Avian Dis.* 46:919-924.
17. Byrne, C. M., M. Clyne, and B. Bourke. 2007. *Campylobacter jejuni* adhere to and invade chicken intestinal epithelial cells in vitro. *Microbiol.* 153:561-569.
18. Carrillo, C. D., E. Taboada, J. H. Nash, P. Lanthier, J. Kelly, P. C. Lau, R. Verhulp, O. Mykytczuk, J. Sy, W. A. Findlay, K. Amoako, S. Gomis, P. Willson, J. W. Austin, A. Potters, L. Babiuk, B. Allans, and C. M. Szymanski. 2004. Genome-wide expression analysis of *Campylobacter jejuni* NCTC11168 reveals coordinate regulation of motility and virulence by *flhA*. *J. Biol. Chem.* 279:20327-20338.
19. Champion, O. L., M. W. Gaunt, O. Gundogdu, A. Elmi, A. A. Witney, J. Hinds, N. Dorrell, and B. W. Wren. 2005. Comparative phylogenomics of the food-borne pathogen *Campylobacter jejuni* reveals genetic markers predictive of infection source. *Proc. Natl. Acad. Sci.* 102:16043-16048.
20. Corry, J. E., and H. I. Atabay. 2001. Poultry as a source of *Campylobacter* and related organisms. *J. Appl. Microbiol.* 90:96S-114S.
21. Cox, N. A., N. J. Stern, K. L. Hiatt, and M. E. Berrang. 2002. Identification of a new source of *Campylobacter* contamination in poultry: Transmission from breeder hens to broiler chickens. *Avian Dis.* 46:535-541.
22. Datta, S., H. Niwa, and K. Itoh. 2003. Prevalence of 11 pathogenic genes of *Campylobacter jejuni* by PCR in strains isolated from humans, poultry meat and broiler and bovine feces. *J. Med. Microbiol.* 52:345-348.
23. Diatchenko, L., Y.-F. C. Lau, A. P. Campbell, A. Chenchik, F. Moqadam, B. Huang, S. Lukyanov, K. Lukyanov, N. Gurskaya, E. D. Sverdlov, and P. D. Siebert. 1996.

- Suppression subtractive hybridization: A method for generating differentially regulated or tissue-specific cDNA probes and libraries. *Proc. Natl. Acad. Sci.* 93:6025-6030.
24. Dudoit, S., Y. Yang, and T. P. Speed. 2000. Statistical methods for identifying differentially expressed genes in replicated cDNA microarray experiments. Berkley Tech Report 578.
  25. Grant, C. R., M. E. Konkel, J. W. Cieplak, and L. S. Tompkins. 1993. Role of flagella in adherence, internalization, and translocation of *Campylobacter jejuni* in nonpolarized and polarized epithelial cell cultures. *Infect. Immun.* 61:1764-1771.
  26. Gregory, E., H. Barnhart, D. W. Dreesen, N. J. Stern, and J. L. Corn. 1997. Epidemiological study of *Campylobacter* spp. in broilers: Source, time of colonization and prevalence. *Avian Dis.* 41:890-898.
  27. Guerry, P., R. Alm, C. Szymanski, and T. J. Trust. 2000. Structure, function, and antigenicity of *Campylobacter* flagella, p. 405-421. In I. Nachamkin and M. J. Blaser (ed.), *Campylobacter* 2nd ed. American Society for Microbiology.
  28. Hanel, I., J. Muller, W. Muller, and F. Schulze. 2004. Correlation between invasion of caco-2 eukaryotic cells and colonization ability in the chick gut in *Campylobacter jejuni*. *Vet. Microbiol.* 101:75-82.
  29. Harvey, S. M. 1980. Hippurate hydrolysis by *Campylobacter fetus*. *J. Clin. Microbiol.* 11:435-437.
  30. Hepworth, P. J., H. Leatherbarrow, C.A. Hart, and C. Winstanley. 2007. Use of suppression subtractive hybridisation to extend our knowledge of genome diversity in *Campylobacter jejuni*. *BMC Genomics* 8:110.
  31. Humphrey, T., S. O'Brien, and M. Madsen. 2007. *Campylobacter* as zoonotic pathogens: A food production perspective. *Intl. J. Food Microbiol.* 117:237-257.
  32. Jacobs-Reitsma, W. F. 2000. *Campylobacter* in the food supply, p. 467-481. In I. Nachamkin and M. J. Blaser (ed.), *Campylobacter*. ASM Press, Washington, D.C.
  33. Jeffrey, J. S., K. H. Tonooka, and J. Lozano. 2001. Prevalence of *Campylobacter* spp. from skin, crop and intestine of commercial broiler chicken carcasses at processing. *Poult Sci.* 80:1390-1392.
  34. Keener, K. M., M. P. Bashor, P. A. Curtis, B. W. Sheldon, and S. Kathariou. 2004. Comprehensive review of *Campylobacter* and poultry processing. *Comprehensive Reviews in Food Science and Food Safety* 3:105-116.
  35. Ketley, J. M. 1997. Pathogenesis of enteric infection by *Campylobacter*. *Microbiol.* 143:5-21.

36. Konkel, M. E., L. A. Joens, and P. F. Mixer. 2000. Molecular characterization of *Campylobacter jejuni* virulence determinants, p. 217-240. In I. N. a. M. J. Blaser (ed.), *Campylobacter*, 2nd ed. American Society for Microbiology.
37. Konkel, M. E., J. D. Klena, V. Rivera-Amill, M. R. Monteville, D. Biswas, B. Raphael, and J. Mickelson. 2004. Secretion of virulence proteins from *Campylobacter jejuni* Is dependent on a functional flagellar export apparatus. *J. Bacteriol.* 186:3296-3303.
38. Lee, M. D., and D. G. Newell. 2006. *Campylobacter* in poultry: filling an ecological niche. *Avian Dis.* 50:1-9.
39. Leonard, E. E., L. S. Tompkins, S. Falkow, and I. Nachamkin. 2004. Comparison of *Campylobacter jejuni* isolates implicated in Guillain-Barre Syndrome and strains that cause enteritis by a DNA microarray. *Infect. Immun.* 72:1199-1203.
40. Monteville, M. R., J. E. Yoon, and M. E. Konkel. 2003. Maximal adherence and invasion of INT 407 cells by *Campylobacter jejuni* requires the CadF outer membrane protein and microfilament reorganization. *Microbiol.* 149:153-165.
41. Murphy, C., C. Carroll, and K. N. Jordan. 2006. Environmental survival mechanisms of the foodborne pathogen *Campylobacter jejuni*. *J. Appl. Microbiol.* 100:623-632.
42. Nachamkin, I. 1995. *Campylobacter* and *Arcobacter*. In E. J. B. P. R. Murray, M. A. Pfaller, F. C. Tenover, and R. H. Tenover (ed.), *Manual of Clinical Microbiology*, 6 ed. ASM Press, Washington, DC.
43. Nesbit, E. G., P. Gibbs, D. W. Dreesen, and M. D. Lee. 2001. Epidemiologic features of *Campylobacter jejuni* isolated from poultry broiler houses and surrounding environments as determined by use of molecular strain typing. *American J. Vet. Res.* 62:190-194.
44. Newell, D. G., and C. Fearnley. 2003. Sources of *Campylobacter* colonization in broiler chickens. *Appl. Environ. Microbiol.* 69:4343-4351.
45. Oyarzabal, O. A., C. Hawk, S. F. Bilgili, C. C. Warf, and G. K. Kemp. 2004. Effects of postchill application of acidified sodium chlorite to control *Campylobacter* spp. and *Escherichia coli* on commercial broiler carcasses. *J Food Prot.* 67:2288-2291.
46. Park, S. F. 2002. The physiology of *Campylobacter* species and its relevance to their role as foodborne pathogens. *Intl. J. Food Microbiol.* 74:177-188.
47. Pearson, B. M., C. Pin, J. Wright, K. I'Anson, T. Humphrey, and J.M. Wells. 2003. Comparative genome analysis of *Campylobacter jejuni* using whole genome DNA microarrays. *FEBS Letters* 554:224-230.

48. Penner, J. L. 1988. The Genus *Campylobacter*: a Decade of Progress. *Clin. Microbiol. Rev.* 1:157-172.
49. Randolph, J. B., and A. S. Waggoner. 1997. Stability, specificity and fluorescence brightness of multiply-labeled fluorescent DNA probes. *Nucl. Acids Res.* 25:2923-2929.
50. Reimers, M. 2005. Statistical analysis of microarray data. *Addict. Biol* 10:10-23.
51. Rinella, E. S., C. D. Eversley, I. M. Carroll, J. M. Andrus, D. W. Threadgill, and D. S. Threadgill. 2006. Human epithelial-specific response to pathogenic *Campylobacter jejuni*. *FEMS Microbiology Letters* 262:236-243.
52. Rollins, D. M., and R. R. Colwell. 1986. Viable but nonculturable stage of *Campylobacter jejuni* and its role in survival in the natural aquatic environment. *Appl. Environ. Microbiol.* 52:531-538.
53. Roszak, D. B., D. J. Grimes, and R. R. Colwell. 1984. Viable but nonrecoverable stage of *Salmonella enteritidis* in aquatic systems. *Can. J. Microbiol.* 30:334-338.
54. Sahin, O., P. Kobalka, and Q. Zhang. 2003. Detection and survival of *Campylobacter* in chicken eggs. *J. Appl. Microbiol.* 95:1070-1079.
55. Sahin, O., T. Y. Morishita, and Q. Zhang. 2002. *Campylobacter* colonization in poultry: sources of infection and modes of transmission. *Ani. Heal. Res. Rev.* 3:95-105.
56. Schena, M., D. Shalon, R. W. Davis, and P. O. Brown. 1995. Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Sci.* 270:467-470.
57. Son, I., M. D. Englen, M. E. Berrang, P. J. Fedorka-Cray, and M. A. Harrison. 2007. Prevalence of *Arcobacter* and *Campylobacter* on broiler carcasses during processing. *Intl. J. Food Microbiol.* 113:16-22.
58. Stern, N. J., P. J. Fedorka-cray, J. S. Bailey, N. A. Cox, S. E. Craven, K. L. Hiatt, M. T. Musgrove, S. R. Ladley, D. Cosby, and G. C. Mead. 2001. Distribution of *Campylobacter* spp in selected U.S. poultry production and processing operations. *J. Food Prot.* 64:1705-1710.
59. Stern, N. J., K. L. Hiatt, G. A. Alfredsson, K. G. Kristinsson, J. Reiersen, H. Hardardottir, H. Briem, E. Gunnarsson, F. Georgsson, R. Lowman, E. Berndtson, A. M. Lammerding, G. M. Paoli, and M. T. Musgrove. 2003. *Campylobacter* spp in Icelandic poultry operations and human disease. *Epidemiol. Infect.* 130:23-32.
60. Thomas, C. J., and T. A. McMeekin. 1980. Contamination of broiler carcass skin during commercial processing procedures: an electron microscopic study. *Appl. Environ. Microbiol.* 40:133-144.

61. van Vliet, A. H. M., and J. M. Ketley. 2001. Pathogenesis of enteric *Campylobacter* infection. *J. Appl. Microbiol.* 90:45S-56S.
62. Vandamme, P. 2000. Taxonomy of the Family Campylobacteraceae, p. 3-26. In I. Nachamkin and M. J. Blaser (ed.), *Campylobacter*, 2 ed. Asm Press.
63. Volokhov, D., V. Chizhikov, K. Chumakov, and A. Rasooly. 2003. Microarray-based identification of thermophilic *Campylobacter jejuni*, *C. coli*, *C. lari*, and *C. upsaliensis*. *J. Clin. Microbiol.* 41:4071-4080.
64. Walker, R. I., M. B. Caldwell, E. C. Lee, P. Guerry, T. J. Trust, and G. M. Ruiz-Palacios. 1986. Pathophysiology of *Campylobacter* enteritis. *Microbiol. Rev.* 50:81-94.
65. Wassenaar, T. M., and M. J. Blaser. 1999. Pathophysiology of *Campylobacter jejuni* infection of humans. *Microbes Infect.* 1:1023-1033.
66. Wildsmith, S. E., and F. J. Elcock. 2001. Microarrays under the microscope. *Mol. Pathol.* 54:8-16.
67. Winstanley, C. 2005. Spot the difference: applications of subtraction hybridisation to the study of bacterial pathogens. *J. Med. Microbiol.* 51:459-467.
68. Zheng, J., J. Meng, S. Zhao, R. Singh, and W. Song. 2006. Adherence to and invasion of human intestinal epithelial cells by *Campylobacter jejuni* and *Campylobacter coli* isolates from retail meat products. *J. Food Prot.* 69:768-774.

Chapter 3  
Simplified Capacitance Monitoring for the Determination  
of *Campylobacter* spp. Growth Rates

---

E. Deann Akins, Patricia L. Rule, Michelle R. Keener, J. Eric Line, Mustafa Simmons, Mark A. Harrison, and Kelli L. Hiatt. Accepted for publication to *Journal of Rapid Methods & Automation in Microbiology*.

## ABSTRACT

Capacitance monitoring is commonly used as an efficient means to measure growth curves of bacterial pathogens. However, the use of capacitance monitoring with *Campylobacter* spp. was previously determined to be difficult due to the complexity of the required media. We investigated capacitance monitoring using a simplified medium for the efficient and reproducible construction of growth curves for *Campylobacter* spp. Initially *Campylobacter* spp. were grown on Mueller Hinton plates under a microaerobic atmosphere (37°C; 24 h), followed by transfer to Mueller Hinton biphasic for 6 h (37°C; microaerobic atmosphere). Serial dilutions were used for inoculation of Bactometer® wells containing 1 mL Mueller Hinton broth plus 0.1M sodium pyruvate for the completion of *Campylobacter* spp. growth curves with the Bactometer®.



## INTRODUCTION

Campylobacteriosis is one of the most common bacterial intestinal disorders of humans in many industrialized countries (Knudsen et al., 2006). There are approximately 20-150 cases per 100,000 reported foodborne illness cases per year in the United States. However, under-reporting is significant and the true incidence could be as high as five to 10 times the reported rate (Lee and Newell, 2006). Epithelial cell invasion is considered to be an essential step in *Campylobacter* spp. infection. Invasion studies using intestinal epithelial cells as well as other cell lines show that the relative ability to invade cultured cells is strain dependent (Poly et al., 2004). When invasion studies are performed, it is required that *Campylobacter* spp. isolates be grown within a range of  $10^6$  to  $10^8$  CFU/ml. Consequently, determination of growth curves for *Campylobacter* spp. can be a time consuming and labor intensive process prior to completing cell invasion or other assays such as gene content or transcriptome analyses using microarrays for this organism.

Microbiological conductimetric methods are often used to monitor bacterial growth. Conductimetric instruments monitor microbial metabolism within a growth medium by the measurement of significant changes in electrical activity including total impedance, conductance, or capacitance (Corry et al., 1995; Wawerla et al., 1999; Line and Pearson, 2003). The Bactometer® microbial monitoring system (bioMerieux, Hazelwood, MO) is one such conductimetric instrument that was previously determined to be efficient for monitoring growth patterns for other bacterial types such as *Salomonella* spp. and *Escherichia coli* (Wawerla et al., 1999). Each Bactometer® module contains 16 wells that contain approximately two mL total volume each, with two electrodes exposed in each well. Due to strict media and atmospheric requirements for growth of *Campylobacter* spp., the use of capacitance monitoring with this

organism was determined to be either labor intensive (requiring complex media formulations) or inefficient and irreproducible with less complex media formulations (personal communications, Patricia Rule). In this investigation, we optimized conditions for use with the Bactometer® such that the efficient and reproducible monitoring of *Campylobacter* spp. for determination of growth curves was achieved in a simple medium.

## MATERIALS AND METHODS

Ten *Campylobacter* spp. isolates (Table 3.1), originally recovered from Iceland, were used for this investigation (Stern et al., 2003). *Campylobacter* spp. require a microaerobic atmosphere (3-5% oxygen, 2-10% carbon dioxide, with a balance of nitrogen) for optimal growth (Altekruse and Swerdlow, 2002). As the Bactometer® does not allow for the manipulation of atmosphere, a 0.5 ml overlay of sterile mineral oil was used in each well to help reduce oxygen from entering wells. Conditions used for optimization are listed in Table 3.2. For the first trial, *Campylobacter* spp. isolates were grown for 24 h on Mueller Hinton agar plates at 37°C under microaerobic conditions followed by transfer to biphasic cultures (25 mL vented capped T-flask containing a 10 mL Mueller Hinton agar base and 5 mL Mueller Hinton broth) for an additional 18 h under the same conditions (Rollins et al., 1983). Serial dilutions, ranging from  $10^{-1}$ - $10^{-6}$  were performed, with 100  $\mu$ L of each dilution added to individual Bactometer® wells containing 1 mL of Mueller Hinton Broth, followed by the addition of mineral oil. Modules were placed into the Bactometer® at 37°C for 48 h following the manufacturer's instructions. Each trial was replicated in triplicate, but no growth was obtained under these initial conditions. Trial 2 was similar with the exception that half of the wells contained 0.2 mL of Mueller Hinton agar to mimic the biphasic culture. After 48 h, no growth was detected in any of the Bactometer® wells, thus biphasic conditions in the wells were not

tested in subsequent experiments. A third trial included the use of 20 mM sodium pyruvate in the primary biphasic cultures during the 18 h of growth. Sodium pyruvate is an organic salt involved with amino acid metabolism and initiates the Krebs cycle where glucose is converted to energy such as ATP. It has also been shown to have protective effects against reactive oxygen species (Corry et al., 1995). Serial dilutions were once again placed into the Bactometer® wells containing 1 mL of Mueller Hinton broth overlaid with a 0.5 mL of sterile mineral oil; again there was no growth of isolates in the Bactometer®. In the fourth trial, the molar concentration of sodium pyruvate was increased to 0.1 M. These conditions resulted in bacterial growth at 48 h, however, the initial bacterial numbers exceeded the Bactometer® threshold value of  $10^6$  CFU/mL. For the fifth trial, the *Campylobacter* spp. isolates were grown in primary biphasic cultures containing a 0.1M concentration of sodium pyruvate, for only 6 h. The Bactometer® wells, containing 1 mL Mueller Hinton broth supplemented with 0.1M sodium pyruvate, were inoculated in triplicate with serial dilutions of the *Campylobacter* spp. isolates. After incubation in the Bactometer® for 48 h, growth curves were successfully obtained as reported in Figure 3.1. A final trial was conducted to determine if 0.1M sodium pyruvate was beneficial when added to both the primary biphasic cultures as well as the Bactometer® wells which only contained Mueller Hinton broth.

## RESULTS AND DISCUSSION

A protocol was developed after all seven trials were completed. Ten *Campylobacter* spp. isolates were grown on Mueller Hinton agar plates for 24 h at 37°C and then inoculated into primary biphasic cultures containing Mueller Hinton broth for an additional 6 h of growth at 37°C under microaerobic conditions. Serial dilutions,  $10^{-1}$ - $10^{-3}$ , were prepared and 100  $\mu$ L of each dilution was placed into wells that contained a 0.1M concentration of sodium pyruvate

along with 1 mL of Mueller Hinton broth. A 0.5 mL overlay of sterile mineral oil was used to help create the microaerobic conditions. All isolates were inoculated in triplicate. Growth curves and threshold times were obtained after incubation in the Bactometer® for 48 h at 37°C (Figure 3.1, Table 3.1).

Our results demonstrate that growth rate can vary greatly among *Campylobacter* isolates, therefore, this method will be useful in studies where establishment of growth curves is critical for subsequent experimental analyses of these bacteria. Our recommendation is that initial *Campylobacter* spp. be grown on Mueller Hinton plates under a microaerobic atmosphere (37°C; 24 h), followed by transfer to Mueller Hinton biphasic cultures for 6 h (37°C; microaerobic atmosphere). Serial dilutions ( $10^{-1}$ - $10^{-3}$ ) should then be used for inoculation of Bactometer® wells containing 1 mL Mueller Hinton broth supplemented with 0.1M sodium pyruvate, followed by an overlay with 0.5 mL of sterile mineral oil.

## References:

1. Altekruze, S.F., D.L. Swerdlow, 2002. *Campylobacter jejuni* and related organisms. In: Foodborne Diseases. D.O. Cliver & H.P. Riemann (eds). San Diego: Academic Press, 103-112.
2. Corry, J.E., D.E. Post, P. Colin, 1995. Culture media for the isolation of *Campylobacters*. *Int. J. Food Microbiol.* 26, 43-76.
3. Kiggens, E.M., W.N. Plastringe, 1956. Effect of gaseous environment on growth and catalase content of *Vibrio fetus* cultures of bovine origin. *J. Bacteriol.* 72, 397-400.
4. Knudsen, K.N., D.D. Bang, L.O. Andresen, M. Madsen, 2006. *Campylobacter jejuni* strains of human and chicken origin are invasive in chickens after oral challenge. *Avian Dis.* 50, 10-14.
5. Lee, M.D., D.G. Newell, 2006. *Campylobacter* in poultry: filling an ecological niche. *Avian Dis.* 50, 1-9.
6. Line, J.E., K.G. Pearson, 2003. Development of a selective broth medium for the detection of injured *Campylobacter jejuni* by capacitance monitoring. *J. Food Prot.* 66, 1752-1755.
7. Poly, F., F. Threadgill, A. Stintzi, 2004. Identification of *Campylobacter jejuni* ATCC 43431-specific genes by whole microbial genome comparison. *J. Bacteriol.* 186, 4781-4795.
8. Rollins, D.M., J.C. Coolbaugh, R.I. Walker, E. Weiss, 1983. Biphasic culture system for rapid *Campylobacter* cultivation. *Appl. Environ. Microbiol.* 45, 284-289.
9. Rule, P.L. bioMerieux, Inc., 595 Anglum Rd, Hazelwood, MO 63042.
10. Stern N.J., K.L. Hiatt, G.A. Alfredsson, K.G. Kristinsson, J. Reiersen, H. Hardardottir, H. Briem, E. Gunnarsson, F. Georgsson, R. Lowman, E. Berndtson, A.M. Lammerding, G.M. Paoli, M.T. Musgrove, 2003. *Campylobacter* spp. in Icelandic poultry operations and human disease. *Epidemiol. Infect.*, 130, 23-32.
11. Wawerla, M., A. Stolle, B. Schalch, H. Eishruher, 1999. Impedance microbiology: applications in food hygiene. *J. Food Prot.* 62, 1488-1496.

Table 3.1: Time required for different strains of *Campylobacter* spp. to reach a detection limit of approximately  $10^6$  CFU/mL by capacitance measurement using the Bactometer<sup>®</sup>.

<b>Isolate</b>	<b>Source</b>	<b>Threshold Time (h)</b>
81-176	human isolate	12
13262	broiler ceca	12
14590	other poultry	12
14131	human isolate	7
14194	poultry production environment	6
4568	domestic species	6
7358	parent breeder fecal	6
12826	broiler ceca	6
8889	parent breeder fecal	6
5069	broiler ceca	6

Table 3.2: Experimental *Campylobacter* spp. growth conditions evaluated for detection of changes in capacitance using the Bactometer<sup>®</sup>

Trials	Primary Biphasic Cultures			Bactometer Well Medium				Growth		
	18 h	6 h + Sodium Pyruvate	6 h	Mueller Hinton		Sodium Pyruvate			Mineral Oil	
				1.0 mL Broth	0.2 mL Agar	20 mM	0.1 M		0.5 mL	
1	x			x				x	No	
2	x			x	x				x	No
3	x			x		x			x	No
4	x			x			x		x	Yes
5		x		x				x	x	Yes
6		x	x	x				x	x	Yes
7			x	x				x	x	Yes




Figure 3.1: Typical mean capacitance response for *Campylobacter* spp. 4568 inoculated into Bactometer® wells at  $10^1$  CFU/mL. Bars represents standard deviation.



## Chapter 4

### Identification of Putative Virulence Factors Utilizing Invasion Assays and Whole Genome Comparisons of *Campylobacter* spp. Isolates Recovered from Iceland

---

E. Deann Akins, Kelli L. Hiett, Bruce S. Seal, Alain Stintzi, Mustafa Simmons, and Mark A. Harrison. To be submitted to BMC Microbiology

## **Abstract**

### Background

*Campylobacter* spp., one of the most common causes of human foodborne bacterial gastroenteritis, is commonly associated with the handling and consumption of raw poultry products and considered to be a major source of *Campylobacter* induced disease in humans. The United States has sought to reduce the number of cases of campylobacteriosis in humans through investigations of the epidemiology and pathogenicity of *Campylobacter* spp. These investigations should be facilitated by employing a library of epidemiologically related *Campylobacter* spp. isolates. The isolates used in this investigation were originally recovered from an extensive epidemiologic investigation conducted in Iceland. We investigated the adherence and invasiveness of 52 *Campylobacter* spp isolates using human colon cells as well as the genetic diversity of four *C. jejuni* isolates that demonstrated a wide range of invasiveness towards human colonic cells.

### Results

*Campylobacter* spp. exhibited a wide range of adhesion and invasion ability, which was determined unrelated to *flaA* short variable region (SVR) allele type. Four isolates comprised of the most invasive isolate (14118), the least invasive (13262), and two in between were selected for comparative genomic analysis. DNA:DNA microarray hybridizations identified genes absent relative to 11168 (PMSRU). Several genes were located in 1 of 7 previously described plasticity regions. There were 372 genes determined present in *C. jejuni* isolates 14118, 5116, 8557, and 13262 as well as *C. jejuni* 11168 (PMSRU). Suppressive subtractive hybridizations identified genes absent from *C. jejuni* 11168 (PMSRU). *C. jejuni* 14118 contained a gene from *C. doylei* 269.97 that encoded for a motility accessory factor and a gene involving transport. *C. jejuni*

13262 contained a cytolethal distending toxin (CDT) operon from *C. lari* as well as a type II restriction modification enzyme unlike isolates 14118, 5116 and 8557 which includes a type I restriction modification enzyme.

### Conclusions

These studies provide further insight into genetic variability of *Campylobacter* spp. The results facilitate the determination of the core *C. jejuni* genome and also provide information regarding putative virulence factors that might explain differences in adhesion and invasiveness. Further investigations of variably present genes, such as hypothetical proteins, should lead way to more knowledge for the development of intervention strategies and biomarkers.

## Background

Infection due to *Campylobacter* spp. exposure to a variety of foods, water, and environmental sources is one of the major causes of human diarrheal disease in industrialized countries [1, 2]. Although *Campylobacter* spp. can asymptotically colonize the intestinal tract of a variety of warm blooded animals, this pathogen often results in human disease ranging from self-limiting gastroenteritis to more serious systemic infections [3]. The majority of *Campylobacter jejuni* cases are enteric, with episodes confined to local acute gastroenteritis characterized by nausea, abdominal pain, diarrhea, and fatigue. Although campylobacteriosis is generally self-limiting, incapacity may last several weeks and up to 10% of cases require medical intervention [4]. *Campylobacter* spp. infections have also been associated with extra-intestinal sequelae such as Guillain-Barré Syndrome [5] and reactive arthritis [6].

The reduction and elimination of the occurrence of foodborne pathogens associated with poultry has received attention in recent years. The increase of commercially processed poultry, which grew from approximately 7.3 to 8.4 billion from 1994-2001 [7], and the subsequent consumption of poultry, the national goal in the U.S., set by governmental agencies, is to reduce the incidence of *Campylobacter* spp. infection from 13.37 per 100,000 people in 2002 to 12.30 per 100,000 people in 2010 [8]. The Center for Disease Control and Prevention (CDC) reported that *Campylobacter* spp. infection decreased 31% between 1996-1998 likely due to the implementation of a new regulation known as FSIS Pathogen Reduction/HACCP Regulation [9]. Since this decrease, the estimated incidence of *Campylobacter* spp. has not changed significantly. In 2007 the number of cases and incidence per 100,000 population were 12.79 for *Campylobacter* spp.[10]. Since the Healthy People 2010 national health target has not been

reached, a need exists to improve understanding of the epidemiology and transmission of *Campylobacter* spp.

Poly et al. [11] found that the ability of different *C. jejuni* isolates to invade cell cultures is strain dependent. Noninvasive strains have been isolated from patients with noninflammatory disease, while invasive strains were isolated from patients with inflammatory diarrhea. These findings suggest that different *Campylobacter* spp. isolates vary in their virulence properties and that these virulence properties are correlated at least in part, with the ability to invade human intestinal epithelial cells [11]. The percent invasion tends to vary based on cell line and specific isolate. Hickey et al. [12] reported that *C. jejuni* 81-176 invaded INT407 cells at a level of 2.1% whereas other campylobacters invaded INT407 cells within a range of 0.001%-0.41%. Hanel et al. [13] investigated 11 *C. jejuni* isolates and determined the range of invasiveness of Caco-2 cells was between 0.00003%-2.14%. The ability of pathogenic bacteria to adhere to host tissues is important for the establishment of an in vivo niche. This binding can be a prerequisite for host cell invasion as with *Campylobacter* spp. [14]. An emerging theme among pathogenic bacteria is their ability to utilize host cell molecules during the infectious process to facilitate their binding and entry into host cells [15].

Investigations into the basis of *Campylobacter* spp. transmission could be facilitated by a library of epidemiologically related *Campylobacter* spp. isolates. A comprehensive epidemiologic investigation was previously conducted in Iceland because the small size allowed for sampling of a well-defined broiler production and processing industry. Iceland's broiler practice is comparable to that of North America and provides the sole source of broiler meat to the Icelandic human population. Additionally, production is on a smaller scale with flock sizes ranging from <1,000 to 13,500 [16]. The limited scale of production and the fact that no broiler

meat products are imported into Iceland enabled an unique total population based epidemiological study that would not be feasible in the U.S. due to the scale of production, complex market distribution, and disperse consuming population.

The epidemiologic analysis included all poultry flocks, other agriculturally important animals, environmental sources, and human clinical cases over a three-year period. *flagellinA* short variable region (*flaA* SVR) DNA sequence analysis of recovered isolates revealed that certain *flaA* SVR subtypes were recovered from chickens as well as from humans. However, there were also *flaA* SVR subtypes that were predominate in poultry, but never recovered from humans. This observation allowed for investigations to facilitate our understanding of the molecular basis and biological consequences of genetic diversity of *C. jejuni* and human disease [17]. Additionally, this information is important to determine if “biomarkers” might exist to allow us to determine if a *Campylobacter* spp. isolate is more likely to result in human disease. To address these goals, cell adhesion/invasion assays, suppressive subtractive hybridizations, and DNA:DNA microarray analyses were utilized to determine genetic differences between invasive and noninvasive *Campylobacter* spp. isolates.

## **Materials and Methods**

### Bacterial Isolates and Growth Conditions

Fifty-two *Campylobacter* spp. isolates, with known spatial and temporal relationships, were used for this investigation (Table 4.1) [16]. These isolates were previously determined to segregate into one of four unique *flaA* SVR allele types. Allele types were selected after comparing the genetic distances between all isolates; the four allele types demonstrating greatest distance were selected (Figure 1). *C. jejuni* 81-176 and *C. jejuni* 11168 (PMSRU) were also included in this study as a control for adhesion/invasion assays, suppressive subtractive

hybridizations, and DNA:DNA microarray analyses respectively. All *Campylobacter* spp. isolates were grown at either 42°C or 37°C for 24 h on Mueller-Hinton (MH) agar (Sigma, St. Louis, MO) under microaerobic conditions (5% O<sub>2</sub>, 10% CO<sub>2</sub>, and 85% N<sub>2</sub>). Isolates were then grown in Mueller-Hinton biphasic cultures for 16 h at 37°C under microaerobic conditions to reach a mid-log phase. *Escherichia coli* DH5 $\alpha$  mcr- and TOP10 (Invitrogen, Carlsbad, CA) cells were cultured aerobically at 37°C on Luria-Bertani (LB) agar plates for 24 h. Transformed TOP10 *E. coli* were grown at 37°C in LB broth supplemented with 50  $\mu$ g/ml of ampicillin.

#### DNA Isolation

For DNA:DNA microarray hybridization and suppressive subtractive hybridization analyses, *Campylobacter* spp. isolates were grown to stationary phase as previously described [18]. Genomic DNA was isolated using a phenol-chloroform extraction. Briefly, cells were pelleted, lysed using an SDS extraction buffer, RNase and proteinase K treated followed by precipitation with 0.2 volume of 10M ammonium acetate. Plasmid DNA was isolate using the Qiagen Plasmid MiniPrep Kit (Valencia, CA), from overnight cultures of transformed *E. coli* TOP10 cells.

#### Cell Culture

Caco-2 cells (provided by Dr. Holly Sellers, PDRC, University of Georgia) were cultured as cell monolayers in modified Eagle medium (MEM) (Cellgro, Herdon, VA) supplemented with nonessential amino acids (Cellgro), sodium pyruvate (Cellgro) and 20% fetal bovine serum (FBS) (Sigma) and incubated at 37°C in a 5% CO<sub>2</sub> humidified incubator [19, 20]. For experimental assays, Caco-2 cell monolayers were seeded at a density of approximately 1 x 10<sup>5</sup> cells into 24 well plates. The plates were incubated at 37°C in a 5% CO<sub>2</sub> humidified incubator

for 16 h prior to infection to allow reformation of the monolayer [21]. Prior to the assay, the cell monolayers were washed with phosphate-buffered saline (PBS) pH 7.2 (Invitrogen) [13].

#### Adherence and Invasion Assay

The bacteria, *Campylobacter* spp. and *E. coli* DH5 $\alpha$  mcr-, were harvested from Mueller Hinton biphasic cultures incubated at 37°C in microaerobic for 12 h and aerobic atmospheres for 12 h, respectively. One mL of cell suspension was removed and centrifuged at 4,000 x rpm for 10 min. Each pellet was resuspended in 1.5 ml of MEM + 1% FBS. The assay was performed by co-incubating mid-log phase isolates in triplicate with semi-confluent Caco-2 cells. Each resuspended isolate (0.5 mL) was inoculated into duplicate wells. Serial dilutions ( $10^1$ -  $10^7$ ) were prepared where 100 $\mu$ l was inoculated onto plates to determine the number of bacteria inoculated into each well [13]. MH agar, incubated at 37°C in a microaerobic atmosphere for 36 h, was utilized for *Campylobacter* spp. isolates while LB agar, incubated at 37°C for 24 h was used for *E. coli*. All plate counts were performed in duplicate. After inoculation, the Caco-2 cells were incubated for 3 h at 37°C in a 5% CO<sub>2</sub> humidified incubator to allow bacterial adhesion and internalization. For determination of adherence, one-half of the cells were washed three times with sterile PBS and the cell monolayer lysed with 0.1% Triton X-100 (Sigma) [14, 22]. The total bacteria associated with the cells (intracellular and extracellular bacteria) were enumerated by plating serial dilutions ( $10^1$ - $10^4$ ), in duplicate on either MH or LB agar. The MH agar plates were incubated at 37°C in a microaerobic atmosphere for 36 h prior to counting. LB agar plates were incubated at 37°C in ambient atmosphere, for 24 h before counting. The remaining wells were measured for bacterial invasion. The infected cells were washed twice with sterile PBS and incubated in fresh culture medium containing 250  $\mu$ g per ml of gentamicin (Sigma) for 3 h to kill remaining viable extracellular bacteria [20]. Quantification of the viable



intracellular bacteria was performed by washing the infected Caco-2 cells with sterile PBS twice followed by subsequent lysis with 0.1% Triton-X 100 [14, 22]. Serial dilutions, ( $10^1$ - $10^3$ ) were plated in duplicate on their respective agars and incubated accordingly as described above.

#### *flaA* SVR DNA Sequence Analysis

*flaA* SVR DNA sequence analysis was performed as previously described [23]. Isolated colonies of *Campylobacter* spp. were suspended in 300  $\mu$ l of sterile H<sub>2</sub>O and placed at 100°C for 10 min. Ten  $\mu$ l of each boiled cell suspension was used as a template for *flaA* SVR PCR with the following primers: FLA242FU: 5' CTA TGG ATG AGC AAT TWA AAA T 3' and FLA625RU: 5'CAA GWC CTG TTC CWA CTG AAG 3'. A 35 cycle reaction was used with 1 min denaturing at 96°C, 1 min annealing at 52°C and 1 min extension at 72°C. The resulting product was approximately 425 bp. Sequence data was generated using either the FLA242 FU primer or the FLA625 RU primer with the Big Dye Dye-Terminator Cycle Sequencing Kit (ABI-PE, Foster City, CA). Data was assembled and edited using Sequencer 4.7 (Genes Codes Corp., Ann Arbor, MI) and aligned using Clustal X (Plate-Forme de Bio-Informatique, France). Aligned sequences were compared and dendrograms were generated using the Neighbor-Joining algorithm with HKY85 distance measurements in PAUP\*4.0 [24].

#### *Campylobacter* spp. Speciation

*Campylobacter* spp. were speciated using a multiplex PCR as previously described by Wang et al. [25].

#### Suppressive Subtractive Hybridization

Suppressive subtractive hybridization was performed [26] using the PCR-select bacterial genome subtraction kit (Clonetech, Palo Alto, CA) with modifications as described below. *C. jejuni* isolate 11168 (PMSRU) was used as the driver while four *C. jejuni* isolates (5116, 14118,

13262, and 8557) recovered from Iceland were used individually as tester samples. Four micrograms of genomic DNA from each isolate was digested with 30 units of AluI (New England Biolabs, Ipswich, MA) and 30 units of DraI (New England Biolabs, Ipswich, MA) for 30 min to produce blunt end fragments ranging approximately 100 to 650 bp in size [17, 27]. Both the first and second subtractions were performed at 59°C. Thermal cycling conditions for enrichment of subtracted sequences were altered to 72°C for 5 min followed by 30 cycles at 94°C for 30 s, 64°C for 30 s and 72°C for 1.5 min.

#### Preparation of the Subtractive-Hybridization Library

Products resulting from PCR amplification were ligated in pCR®2.1 vector using the TA Cloning® Kit Verson V (Invitrogen, Carlsbad, CA) at 14°C. The ligated products were transformed into One Shot TOP10 chemically competent *E. coli* cells using the TA Cloning® Kit Verson V and transformants were selected for kanamycin and ampicillin resistance. White colonies were picked, transferred to LB broth supplemented with ampicillin (50 µg/ml), and grown at 37°C overnight with agitation (200 rpm). Plasmid DNA was isolated as previously stated.

#### DNA Sequence Analysis

Plasmid DNA was digested with 40 units of EcoRI (New England Biolabs) and resolved in a 0.8% agarose gel. Plasmids that released an insert upon digestion were further analyzed by DNA sequence analysis, using the Big Dye Terminator v3.1 Sequencing kit (PE Applied Biosystems, Foster City, CA). DNA sequences were assembled and edited using Sequencher 4.7 (Gene Codes Corporation; Ann Arbor, MI), and similarity searches performed using BLASTN and BLASTX (<http://www.ncbi.nlm.nih.gov/BLAST/>)[28]. Unique DNA sequences obtained

during this investigation were submitted to GenBank (accession numbers provided upon final revision).

#### Microarray Construction, DNA Labeling, DNA:DNA Microarray Hybridization, and Analysis

DNA microarrays used for analysis were prepared as previously described [27]. Primers from the BioPrime Labeling Kit (Invitrogen) were used for random primer labeling of genomic DNA in the presence of amino-allyl dUTP (Ambion, Austin, TX), followed by coupling to either Cy3 or Cy5 (GE Healthcare, Piscataway, NJ) monoreactive flours [18]. Microarrays were hybridized overnight at 42°C and subsequently visualized using a Packard Scan Array Light 2-color scanner with Scan Array Express Software Version 1.1 (Packard BioScience, La Jolla, CA). Three independent hybridization experiments were performed yielding 9 measurements per gene (given that each gene is present thrice on each microarray). Analyses were performed using the software program GACK (genotyping analysis; C.Kim [Stanford University, Stanford, CA]; available at <http://cmgm.stanford.edu/falkow/whatwedo/software/software.html>). This program calculates an idealized normal distribution curve for each array and assigns a binary value to each data point on the microarray based on an estimated probability that a gene is present or absent in a given isolate, relative to 11168 (PMSRU), the genome sequenced isolate.

## **Results**

### *Campylobacter* spp. *flaA* SVR Allele Types

Four *Campylobacter* spp. *flaA* SVR allele groups were chosen for analysis based on comparison of the genetic distances between all *Campylobacter* spp. isolates from the original Iceland investigation (Table 4.1, Figure 4.1). The first *flaA* SVR allele type, DAA, contained 21 isolates ranging in sample origin from human clinical isolates, other poultry isolates (turkey, etc), domestic species, socks (environmental samples), broiler caeca samples, parent breeder fecal

samples, retail product samples, and grandparent fecal samples. *flaA* SVR allele type DAB was comprised of 15 isolates which also included human clinical isolates, broiler caeca samples, other poultry samples, parent breeder fecal samples, and retail product samples. The third *flaA* SVR allele type, DAC, included 5 isolates that were collected from broiler caeca samples, wild bird samples, parent breeder and rearing fecal samples. The last *flaA* SVR allele type, DAD, included 11 isolates from domestic species samples, human clinical samples, other poultry samples, parent breeder fecal samples and retail product samples. Interestingly human clinical isolates were found only in three *flaA* SVR allele types with no human isolates originating from DAC *flaA* SVR allele type.

#### Adhesion of *Campylobacter* spp. with Caco-2 cells

Fifty-two *Campylobacter* spp. isolates (Table 4.1) were used in cell adhesion assays with *E. coli* DH5 $\alpha$  mcr- employed as the negative control, and the highly invasive *C. jejuni* 81-176 [29] employed as the positive control. The average percent adhesion ranged from 0.00008% to 3.4%. *C. jejuni* 11168 (PMSRU) adhered at 0.025% and *C. jejuni* 81-176 adhered at 0.083%. A histogram illustrating the distribution of *flaA* SVR allele groups relative to percent adhesion is presented in Figure 4.2. The level of adhesion was arbitrarily assigned to 3 groups; low (<0.08%), medium (0.08-0.8%), and high (>0.8%). Each *flaA* SVR allele type was represented in each adhesion level. Sixty-six percent of *flaA* SVR allele group DAB was located in the low adhesion range. *flaA* SVR allele group DAD isolates (n=7) were found primarily in the medium range whereas at the high level, allele type DAC had 40% of its isolates (2 out of 5).

There appeared to be no relationship demonstrated between host of recovery source and level of adhesion as illustrated in Figure 4.3. Four of nine human isolates adhered at a percentage of 0.71% or greater. Of the remaining 5 human isolates, 4 fell within the middle

adherence range of 0.1%-0.051% while one *C. jejuni* isolate was present in the low range with an adherence of 0.01%. The poultry isolates were distributed throughout all levels of adhesion.

#### Invasion of *Campylobacter* spp. within Caco-2 cells

Isolates were also tested for their invasiveness of a human Caco-2 cell line. The percent invasion ranged from 0.000003% to 1.2%. *C. jejuni* 11168 (PMSRU) invaded at 0.035% whereas *C. jejuni* 81-176 invaded at 0.75%. Again, percent invasion was arbitrarily divided into 3 levels; low (<0.0025%), medium (0.0025-0.01%), and high (>0.01%) (Figure 4.4). In the high range of invasion, *flaA* SVR allele group DAA occurred the most frequently followed by *flaA* SVR allele groups DAB and DAD. The most invasive isolate, belonging to *flaA* SVR allele DAD, invaded at a higher level than did *C. jejuni* 81-176, the positive control. Interestingly, *flaA* SVR allele group DAC (n=5), which contained no human isolates, was not found in the high invasive range. All five of the *flaA* SVR allele group DAC isolates invaded Caco-2 cells below 0.002%. These isolates were recovered from such sources as parent breeder fecal samples, wild birds, parent rearing fecal samples, and commercial broiler caeca. Isolates belonging to *flaA* SVR allele group DAB, recovered from broiler caeca, retail products, human clinical samples, parent breeder fecal samples, other poultry fecal samples, and parent rearing fecal samples, were found between in all levels of invasion. Four of fifteen isolates (26.6%) within this *flaA* SVR allele group were found to be highly invasive whereas 11 of fifteen isolates (73.3%) were found to have a percent invasion less than 0.0025%. No isolates fell into the medium range of invasiveness with this particular allele group.

Figure 4.5 represents the distribution of isolate source relative to percent invasion. The majority (7 of 9) of the human isolates collected invaded the Caco-2 cells at a rate greater than 0.003%. Four out of the nine human isolates demonstrated invasion levels greater than 0.012%

while the four remaining human isolates invaded within the range of 0.002% to 0.01%. Only one human isolate was placed in the low invasive range at 0.001%. The isolates recovered from wild birds and other domestic animals were not found to be invasive, with invasion rates of Caco-2 cells at <0.0006%. The other poultry isolates were distributed within the three invasion ranges.

#### DNA:DNA Microarray Hybridization Analysis

Four *Campylobacter* spp. isolates, chosen on the basis of the invasiveness of Caco-2 cells, were used for subsequent DNA:DNA microarray hybridization assays. Isolate 14118 demonstrated the highest percent invasion (1.2%) to the Caco-2 cells, isolate 13262 demonstrated low percentage (0.00005%) of invasion while isolates 5116 and 8557 fell in the medium range of the invasion of Caco-2 cells. Isolate 14118 (high invasion) did not have 11 genes relative to *C. jejuni* 11168 (PMSRU) (Table 4.2) while isolate 5116 (medium invasion) did not have 18 genes (Table 4.3). The absent genes from isolate 14118 included 4 genes coding for hypothetical proteins with yet to be defined functions. The remainder of the absent genes coded for integral membrane proteins (*cj0860*), transferases (*cj0407* & *cj1331*), and transport proteins that link inner and outer membranes (*cj0753c*). Isolate 5116 (medium range) missing genes included a putative binding protein (*cj0412*), a bacterioferritin (*cj1534c*), involved in oxidative damage protection, integral membrane proteins, multidrug transporter membrane component (*cj1587c*) and a chemotaxis signal transduction protein (*cj0262c*). There were also six hypothetical proteins with unknown function determined absent. Hybridization analyses demonstrated that of these four isolates, isolate 8557 (medium invasion range) demonstrated the greatest similarity to *C. jejuni* 11168 (PMSRU). There were only three genes of the 1634 tested from the 11168 (PMSRU) genome that were identified as absent in 8557 (Table 4.4). The three genes represent a

hypothetical protein (*cj0056c*), a putative aminotransferase (*cj1436c*) and a *parB* family protein (*cj0101*), which is a predicted transcriptional regulator. The least invasive isolate, 13262 was determined to be the most divergent with 66 genes absent relative to 11168 (PMSRU) (Table 4.5). These genes are related to virulence properties and included cell division proteins, integral membrane proteins, the cytolethal distending toxin, binding proteins and 19 hypothetical proteins.

There were 372 genes determined to be conserved between *C. jejuni* isolate 11168 (PMSRU) and all four *C. jejuni* isolates 14118, 5116, 8557 and 13262 tested using DNA microarray analyses. The common sequences included the core genes responsible for metabolic, cellular, and regulatory processes (Table 4.6) [30]. Fifty-nine of the genes (15.9%) were determined to have unknown function.

#### Identification of Unique *Campylobacter* spp. DNA Fragments Present in *C. jejuni* isolates relative to *C. jejuni* 11168 (PMSRU) by Suppressive Subtractive Hybridization

Suppressive subtractive hybridizations were performed to identify DNA sequences present in *C. jejuni* isolates 14118, 8557, 5116, and 13262 that are absent in 11168 (PMSRU). The initial round of subtractive hybridization at 63°C indicated that the subtraction hybridization had occurred at a low efficiency. This has been recognized as a previous issue with *Campylobacter* spp. [17]. The hybridization temperature was thus lowered to 59°C, which greatly enhanced the annealing of homologous driver and tester DNA, and the subsequent amplification of tester-specific DNA. The number of subtracted clones for all four isolates along with the number of those clones determined to contain inserts and the clones determined to be false positive, is presented in Table 4.7.

Subtracted sequence analysis of isolate *C. jejuni* 14118 (high Caco-2 invasion level) resulted in 68 clones, 2 showing redundancy such that 66 unique clones were analyzed (Table 4.8). The unique *C. jejuni* 14118 clones differing from 11168 (PMSRU) were found to possess significant similarity to *Campylobacter* spp. other than *C. jejuni* 11168 (PMSRU) and the similarities are listed in Table 4.9. The unique subtracted clones that differ from *C. jejuni* demonstrated similarity to *C. doylei* 269.97 and *Blastopirellula marina*.

Subtracted sequence analysis of isolate *C. jejuni* 5116 (medium Caco-2 invasion level) resulted in 24 clones, 8 showing redundancy such that 16 unique clones were analyzed (Table 4.10). The unique clones determined to differ from 11168 (PMSRU), found to possess significant similarity to *Campylobacter* spp. other than *C. jejuni* 11168 (PMSRU), are listed in Table 4.11. Clones determined to differ from *C. jejuni* demonstrated similarity to *C. doylei* 269.97 and *Campylobacter* phage CGC-2007. One clone was also determined to be of hypothetical or unknown function.

Subtracted sequence analysis of isolate *C. jejuni* 8557 (medium Caco-2 invasion level) resulted in 79 clones, 8 showing redundancy such that 71 unique clones were analyzed (Table 4.12). The unique clones differing from 11168 (PMSRU) were found to possess similarity to *Campylobacter* spp. other than *C. jejuni* 11168 (PMSRU), are presented in Table 4.13. The unique clones that differed from *C. jejuni* demonstrated similarity to *C. doylei* 269.97 and *Campylobacter* phage. Nine subtracted clones were determined to be similar to either unknown or hypothetical proteins with unknown function.

Subtracted sequence analysis of isolate *C. jejuni* 13262 (low Caco-2 invasion level) resulted in 59 clones, 4 showing redundancy such that 55 unique clones were analyzed (Table 4.14). The clones determined to differ from 11168 (PMSRU) possessed significant similarity to



*Campylobacter* spp. other than *C. jejuni* 11168 (PMSRU) (Table 4.15). The unique clones that differed from *C. jejuni* were *C. lari*, *C. doylei* 269.97, and *Beggiatoa* spp. Eleven clones were determined to be either hypothetical proteins or have unknown function.

For all four *C. jejuni* isolates, the subtracted sequences grouped into 8 general functional categories as defined by Poly et al. [11]. The first functional group is cell envelope and surface structures while the second category was designated a restriction modification, recombination and repair category. The third category is transport and the fourth category is small molecule metabolism. Bacteriophage sequence comprises another category with the sixth category involving bacterial toxins and chemotaxis is the seventh category. The last category was hypothetical and unknown proteins. Figure 4.6 presents the number of analyzed clones within each functional category for all four *C. jejuni* isolates.

## **Discussion**

It was predicted that cell invasion assays and whole genome comparison would provide insight as to whether certain *flaA* SVR allele types of *Campylobacter* spp. are more pathogenic to humans. The *flaA* short variable region (SVR) is a single locus that has been utilized as a reliable typing method [23, 31, 32]. This typing method has been found to be more discriminatory than serotyping or PCR-restriction fragment length polymorphism of the *flaA* gene. Investigations have demonstrated that *flaA* SVR comparisons can follow the spread of *Campylobacter* spp. populations within the poultry industry [33]. However, Dingle et al. [34] determined that since the *flaA* SVR region had a great allelic diversity at the nucleotide sequence level, this typing technique was not useful for long term population and evolutionary analyses. The *flaA* SVR typing technique is useful for discriminating between related isolates, for example, distinguishing outbreak strains [23]. Knudsen et al. [35] found that phenotypic and

genetic typing methods have proven useful for epidemiological studies of *Campylobacter* spp. infections; however, the prevalence of a specific genotype was not necessarily an indicator for colonizing ability. The results from the current study demonstrated that the *flaA* SVR was not indicative of cell invasiveness and thus there was no correlation between adherence/invasion and the *flaA* SVR allele type. Hanel et al. also reported no association between *flaA* type and invasion into Caco-2 cells when they investigated 11 *C. jejuni* isolates of different origin [13]. Other researchers have demonstrated that the ability to invade human cell lines and the degree to which *Campylobacter* spp. invades eukaryotic cells is dependent on the *Campylobacter* spp. strain. Ketley [36] also stated that clinical isolates appear to be more efficient in the invasion of human cells. This is not in agreement with the results of this investigation as isolates recovered from human clinical cases demonstrated invasion at both high and low levels. Biswas et al. [37] found that the ability to invade cultured cells is also strain dependent but quite variable in efficiency.

Studies show that *C. jejuni* translocation through invaded cells occurs due to a transcellular process rather than via intercellular spaces [38]. The specific function for translocation is not known, but it does depend on functional flagella [20]. Individual host cell type could also play a role with invasion of *Campylobacter* spp. into epithelial cells. The internalization mechanism triggered by *C. jejuni* has been associated with the combined effect of microfilaments (MF) and microtubules (MT) of host cells [22]. Most invasive bacteria including *Salmonella*, *Shigella*, *Listeria* and *Yersinia* spp. [21] trigger microfilament dependent entry pathways. *Neisseria* and *Klebsiella* spp. require both microfilaments and microtubules for invasion. *C. jejuni* has been shown to be less invasive by both MF and MT depolymerization with human intestinal cells [37]. Since the *flaA* SVR is not indicative of virulence, each isolate within an allele type may potentially differ in their genome type.

Whole genome comparison techniques including DNA:DNA microarray hybridization and suppressive subtractive hybridization, were utilized to further identify genetic differences in the four *C. jejuni* isolates that exhibited different cell invasion abilities. Pearson et al., using DNA:DNA microarray hybridizations, [39] characterized seven hypervariable plasticity regions, PR1-7, among *C. jejuni* isolates recovered from diverse origins. PR 1 contains genes encoding the molybdenum transport apparatus and pantothenate biosynthesis genes. PR 2 consists of putative membrane transporters and hypothetical proteins whereas PR 3 consists of ABC transporters and hypothetical proteins. PR 4 contains the N-acetyl neuraminic acid synthase genes, which are involved in the sialylation of lipooligosaccharide (LOS). The LOS biosynthesis genes and post-translational modification of the flagellin genes are located in PR 5. The last two regions code for capsule biosynthesis (PR 6) and membrane proteins (PR 7) [30, 39].

Isolate *C. jejuni* 14118, the most invasive isolate, revealed only one gene missing related to *C. jejuni* isolate within PR 5. This gene, *cj1331*, codes for *pmtB* acylneuraminate cytidyltransferase for flagella modification. Isolate 8557 had one absent gene relative to *C. jejuni* 11168 (PMSRU) from PR 6. Gene *cj1436c* is a putative aminotransferase and is involved in capsular polysaccharide biosynthesis. *cj1724c* was absent relative to *C. jejuni* 11168 (PMSRU) from isolate 5116 and is within PR 7. This gene codes for a 7-cyno-7-deazaguanine reductase. The least invasive isolate, 13262, revealed absence of five genes relative to *C. jejuni* 11168 (PMSRU) within the seven plasticity regions. Gene *cj0303c*, which codes for *modA*, a putative molybdate-binding lipoprotein, belongs to PR 1. Gene *cj0487*, a hypothetical protein, was absent from isolate 13262 and this particular gene falls within the 2nd PR. The third PR includes genes from ABC transporters, membrane and hypothetical proteins. One gene, *cj0737*,

fell within this region. This gene is a putative periplasmic protein and is involved in heme utilization or adhesion. The least invasive isolate, *C. jejuni* 13262, also had two genes absent relative to *C. jejuni* 11168 within PR 6. This region includes genes from the capsular biosynthesis locus. Gene *cj1435c* is a hypothetical protein whereas *cj1436c* is a putative aminotransferase. Both these genes are involved with amino acid transport and metabolism for polysaccharide biosynthesis.

The use of suppressive subtractive hybridization [26] resulted in the recovery of gene sequences determined to be similar to several *Campylobacter* spp. other than *C. jejuni*. Isolates 5116 and 8557, the mid-range invasive isolates, contained clones demonstrating similarity to that of *Campylobacter* spp. phage genes. Previous investigations revealed that up to a total of 89% of the *C. jejuni* isolates and 14% of *C. coli* isolates could be infected by at least one of the bacteriophages [40]. Isolates 5116 and 8557 also had genes similar to those found in *C. doylei* 269.97. These genes are involved in capsular polysaccharide biosynthesis. *C. doylei* can cause both gastritis as well as enteritis and are isolated more often from blood cultures than stool cultures [41]. A five year study conducted in Australia reported that *C. doylei* was isolated from 85.2% of *Campylobacter/Helicobacter* related bacteremia cases [41].

The most invasive isolate, 14118, also contained a gene from *C. doylei* 269.97 that encoded for a motility accessory factor. Flagella-mediated motility is recognized to be one of the factors contributing to *C. jejuni* virulence. The motility accessory factor (*maf*) is a family of flagellin-associated proteins that is involved in flagella glycosylation. This large cluster also contains genes thought to be involved in sugar biosynthesis and transport [42]. An additional *C. doylei* gene involving transport was also found in isolate 14118. This invasive isolate contains a major facilitator superfamily (MSF) protein. MSF is a set of drug efflux proteins that can

contribute to both natural insensitivity to antibiotics and to emerging antibiotic resistance thus these may be potential targets for the development of new antibacterial drugs [43].

The least invasive isolate, 13262, contained a cytolethal distending toxin (CDT) operon from *C. lari*. As previously discussed, the microarray hybridization results determined that this isolate had the *cdtA* gene from *C. jejuni* 11168(PMSRU) absent. The pathogenicity of the species *C. lari*, generally isolated from the intestines of gulls, shellfish, fish and other animals is unknown [44]. Isolate 13262 also has a type II restriction modification enzyme unlike isolates 14118, 5116 and 8557 which include a type I restriction modification enzyme. The role of restriction modification systems in *Campylobacter* spp. is unclear. These specific enzymes might be involved in the breakdown of foreign DNA [45]. These enzymes might also be necessary for stimulating the formation of DNA fragmentation and recombination, resulting in antigenic diversity and variation, such as the homologous recombination observed for the virulence-associated flagellin locus of *C. jejuni* [46]. In a study with *Helicobacter pylori* the presence of restriction modification proteins was associated with the ability of the bacteria to infect its host [47]. This may suggest that these enzymes might affect virulence gene expression. *C. jejuni* could have a similar function where these enzymes control expression of genes involved in colonization.

Lastly, all isolates include multiple unknown and hypothetical proteins, which will require further research into the significance of their contribution to potential virulence. The identification of virulence genes and gene products could help improve diagnostic methods and help determine intervention strategies.

## Conclusion

The determination of virulence factors is important for the reduction of campylobacteriosis in humans. Although the invasion assays did not show that *flaA* SVR subtype was an indicator of virulence, invasion assays did reveal a wide range of invasiveness. DNA:DNA microarray hybridization and suppressive subtractive hybridizations revealed genetic differences between the isolates that should facilitate our understanding of variability in virulence. Future studies should now focus on the putative virulence factors elucidated in this study along with the numerous hypothetical proteins with unknown function. Further knowledge of *Campylobacter* spp. virulence should lead to novel intervention strategies including vaccines that target invasive *Campylobacter* spp.

## References

1. Gaynor EC, Cawthraw S, Manning G, MacKichan JK, Falkow S, Newell DG: **The Genome-Sequenced Variant of *Campylobacter jejuni* NCTC 11168 and the Original Clonal Clinical Isolate Differ Markedly in Colonization, Gene Expression, and Virulence-Associated Phenotypes.** *J Bacteriol* 2004, **186**:503-517.
2. Gormley FJ, MacRae M, Forbes KJ, Ogden ID, Dallas JF, Strachan NJC: **Has Retail Chicken Played a Role in the Decline of Human Campylobacteriosis?** *Appl Environ Microbiol* 2008, **74**:383-390.
3. Hofreuter D, Tsai J, Watson R, Novik V, Altman B, Benitez M, Clark C, Perbost C, Jarvie T, Du L, Galan J: **Unique Features of a Highly Pathogenic *Campylobacter jejuni* Strain.** *Infect Immun* 2006, **74**:4694-4707.
4. Lee MD, Newell DG: ***Campylobacter* in Poultry: Filling an Ecological Niche.** *Avian Dis* 2006, **50**:1-9.
5. Winer JB: **Guillain Barre Syndrome.** *J Clin Pathol* 2001, **54**:381-385.
6. Altekruse SF, Swerdlow DL: ***Campylobacter jejuni* and related organisms.** In *Foodborne Diseases*. 2 edition. Edited by Cliver DO, Riemann HP. San Diego: Academic Press; 2002: 103-112
7. Service GAS: **2002 Georgia Poultry Facts.** *GASS* 2002:29.
8. **Healthy People 2010** [[www.healthypeople.gov](http://www.healthypeople.gov)]. Access 10 June 2008.
9. **General guidelines for implementation of HACCP in a poultry processing plant**
10. **CDC: Preliminary Foodnet data on the incidence of infection with pathogens transmitted commonly through food--10 states, 2007.** *MMWR* 2008, **57**:366-370.
11. Poly F, Threadgill D, Stintzi A: **Identification of *Campylobacter jejuni* ATCC 43431-specific genes by whole microbial genome comparisons.** *J Bacteriol* 2004, **186**:4781-4795.
12. Hickey TE, Baqar S, Bourgeois AL, Ewing CP, Guerry P: ***Campylobacter jejuni*-Stimulated Secretion of Interleukin-8 by INT407 Cells.** *Infect Immun*, 1999, **67**:88-93.
13. Hanel I, Muller J, Muller W, Schulze F: **Correlation between Invasion of Caco-2 Eukaryotic Cells and Colonization Ability in the Chick Gut in *Campylobacter jejuni*.** *Veterinary Microbiology* 2004, **101**:75-82.
14. Monteville MR, Yoon JE, Konkell ME: **Maximal Adherence and Invasion of INT 407 cells by *Campylobacter jejuni* requires the CadF Outermembrane protein and Microfilament Reorganization.** *Microbiol* 2003, **149**:153-165.
15. Watarai M, Funato S, Sasakawa C: **Interaction of Ipa proteins of *Shigella flexneri* with integrin promotes entry of the bacteria into omammalian cells.** *J Exp Med* 1996, **183**:991-999.
16. Stern NJ, Hielt KL, Alfredsson GA, Kristinsson KG, Reiersen J, Hardardottir H, Briem H, Gunnarsson E, Georgsson F, Lowman R, et al: ***Campylobacter* in Icelandic poultry operations and human disease.** *Epidemiol Infect* 2003, **130**:23-32.
17. Ahmed IF, Manning G, Wassenaar TM, Cawthraw S, Newell DG: **Identification of genetic differences between two *Campylobacter jejuni* strains with different colonization potentials.** *Microbiol* 2002, **148**:1203-1212.

18. Poly F, Threadgill D, Stintzi A: **Genomic Diversity in *Campylobacter jejuni*: Identification of *C. jejuni* 81-176-Specific Genes.** *J Clin Microbiol* 2005, **43**:2330-2338.
19. Bras AM, Ketley JM: **Transcellular translocation of *Campylobacter jejuni* across human polarised epithelial monolayers.** *FEMS Microbiology Letters* 1999, **179**:209-215.
20. Grant CR, Konkel ME, W. Cieplak J, Tompkins LS: **Role of Flagella in Adherence, Internalization, and Translocation of *Campylobacter jejuni* in Nonpolarized and Polarized Epithelial Cell Cultures.** *Infect Immun* 1993, **61**:1764-1771.
21. Biswas D, Itoh K, Sasakawa C: **Uptake pathways of clinical and healthy animal isolates of *Campylobacter jejuni* into INT-407 cells.** *FEMS Immunology and Medical Microbiology* 2000, **29**:203-211.
22. Friis LM, Pin C, Pearson BM, Wells JM: **In vitro cell culture methods for investigating *Campylobacter* invasion mechanisms.** *J Microbiol Meth* 2005, **61**:145-160.
23. Meinersmann RJ, Helsel LO, Fields PI, Hiatt KL: **Discrimination of *Campylobacter jejuni* isolates by fla gene sequencing.** *J Clin Microbiol* 1997, **35**:2810-2814.
24. Swofford DL: **PAUP: Phylogenetic Analysis Using Parsimony and other Methods, version 4.0.** 1988.
25. Wang G, Clark CG, Taylor TM, Pucknell C, Barton C, Price L, Woodward DL, Rodgers FG: **Colony Multiplex PCR Assay for Identificaton and DIfferentiation of *Campylobacter jejuni*, *C. coli*, *C. lari*, *C. upsaliensis*, and *C. fetus* subsp. *fetus*.** *J Clin Microbiol* 2002, **40**:4744-4747.
26. Diatchenko L, Lau Y-FC, Campbell AP, Chenchik A, Moqadam F, Huang B, Lukyanov S, Lukyanov K, Gurskaya N, Sverdlov ED, Siebert PD: **Suppression subtractive hybridization: A method for generating differentially regulated or tissue-specific cDNA probes and libraries.** *Proc Nat Acad Sci* 1996, **93**:6025-6030.
27. Hiatt KL, Stintzi A, Andacht TM, Kuntz RL, Seal BS: **Genomic Differences between *Campylobacter jejuni* isolates identify surface membrane and flagellar function gene products potentially important for colonizing the chicken intestine.** *Functional Integrated Genomics* 2008. Accepted for publication.
28. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: **Basic local alignment search tool.** *J Mol Biol* 1990, **215**:403-410.
29. Black RE, Levine MM, Clements ML, Hughes TP, Blaser MJ: **Experimental *Campylobacter jejuni* infections in humans.** *J Infect Dis* 1988, **157**:472-479.
30. Champion OL, Gaunt MW, Gundogdu O, Elmi A, Witney AA, Hinds J, Dorrell N, Wren BW: **Comparative phylogenomics of the food-borne pathogen *Campylobacter jejuni* reveals genetic markers predictive of infection source.** *Proc Nat Acad Sci* 2005, **102**:16043-16048.
31. Meinersmann RJ, Phillips RW, Hiatt KL, Fedorka-Cray P: **Differentiation of *Campylobacter* populations as demonstrated by flagellin short variable region sequences.** *Applied and Environmental Microbiology* 2005, **71**:6368-6374.
32. Hiatt KL, Seal BS, Siragusa GR: ***Campylobacter* spp. subtype analysis using gel-based repetitive extragergenic palindromic-PCR discriminates in parallel fashion to flaA short variable region DNA sequence analysis.** *J Appl Microbiol* 2006, **101**:1249-1258.



33. Cox NA, Stern NJ, Hiatt KL, Berrang ME: **Identification of a new source of *Campylobacter* contamination in poultry: Transmission from breeder hens to broiler chickens.** *Avian Dis* 2002, **46**:535-541.
34. Dingle KE, F.M. Colles, R.Ure, J.A. Wagenaar, B. Duim, F.J. Bolton, A.J. Fox, D. Wareing, M. Maiden: **Molecular Characterization of *Campylobacter jejuni* Clones: A Basis for Epidemiologic Investigation.** *Emer Infect Dis* 2002, **8**:949-955.
35. Knudsen KN, Bang DD, Andresen LO, Madsen M: ***Campylobacter jejuni* strains of human and chicken origin are invasive in chickens after oral challenge.** *Avian Dis* 2006, **50**:10-14.
36. Ketley Jm: **Pathogenesis of enteric infection by *Campylobacter*.** *Microbiol* 1997, **143**:5-21.
37. Biswas D, Itoh K, Sasakawa C: **Role of Microfilaments and Microtubules in the Invasion of INT-407 Cells by *Campylobacter jejuni*.** *Microbiol Immunol* 2003, **47**:469-473.
38. Bras AM, Ketley JM: **Transcellular translocation of *Campylobacter jejuni* across human polarised epithelial monolayers.** *FEMS Microbiol Let* 1999, **179**:209-215.
39. Pearson BM, C. Pin, J. Wright, K. l'Anson, T. Humphrey, and J.M. Wells: **Comparative genome analysis of *Campylobacter jejuni* using whole genome DNA microarrays.** *FEBS Letters* 2003, **554**:224-230.
40. Hansen VM, Rosenquist H, Baggesen DL, Brown S, Christensen BB: **Characterization of *Campylobacter* phages including analysis of host range by selected *Campylobacter* Penner serotypes.** *BMC Microbiology* 2007, **7**:1-9.
41. Parker CT, Miller WG, Horn ST, Lastovica AJ: **Common genomic features of *Campylobacter jejuni* subsp. *doylei* strains distinguish them from *C. jejuni* subsp. *jejuni*.** *BMC Microbiology* 2007, **7**:1-9.
42. Karlyshev AV, D.Linton, Gregson NA, Wren BW: **A novel paralogous gene family involved in phase-variable flagella-mediated motility in *Campylobacter jejuni*.** *Microbiol* 2002, **148**:473-480.
43. Saidijam M, Benedetti G, Ren Q, Xu Z, Hoyle CJ, Palmer SL, Ward A, Bettaney KE, Szakonyi G, Meuller J, et al: **Microbial drug efflux proteins of the major facilitator superfamily.** *Current Drug Targets* 2006, **7**:793-811.
44. Duim B, Vandamme PAR, Rieger A, Laevens S, Dijkstra JR, Wagenaar JA: **Differentiation of *Campylobacter* species by AFLP fingerprinting.** *Gen Mol Biol* 2001, **147**:2729-2737.
45. Stintzi A: **Gene expression profile of *Campylobacter jejuni* in response to growth temperature variation.** *J Bacteriol* 2003, **185**:2009-2016.
46. Harrington CS, Thomson-Carter FM, Carter PE: **Evidence for recombination in the flagellin locus of *Campylobacter jejuni*: implications for the flagellin gene typing scheme.** *J Clin Microbiol* 1997, **35**:2386-2392.
47. Bjorkholm BM, Guruge JL, Oh JD, Syder AJ, Salama N, Guillemin K, Falkow S, Nilsson C, Falk PG, Engstrand L, Gordon JI: **Colonization of germ-free transgenic mice with genotyped *Helicobacter pylori* strains from a case control study of gastric cancer reveals a correlation between host responses and HsdS components of type I restriction-modification systems.** *J Biol Chem* 2002, **277**:34191-34197.

Table 4.1. *flagellinA* short variable region (SVR) allele, sample number, and sample origin of *Campylobacter* spp. isolates employed in this investigation.

<i>FlaA</i> SVR Allele	Sample Number	Sample Origin
DAA	7845	Human Isolate
	14590	Other Poultry
	13759	Human Isolate
	14131	Human Isolate
	4568	Domestic Species
	14194	Socks
	16229	Other Poultry
	5069	Broiler Caeca
	12826	Broiler Caeca
	12578	Broiler Caeca
	8952	Parent Breeder Fecal
	7678	Parent Breeder Fecal
	7486	Parent Breeder Fecal
	7303	Parent Rearing Fecal
	7599	Parent Rearing Fecal
	11393	Parent Rearing Fecal
	12920	Retail Product
	13829	Retail Product
	12620	Retail Product
	10914	Grandparent Fecal
10905	Grandparent Fecal	
DAB	8557	Human Isolate
	8559	Human Isolate
	14078	Human Isolate
	5116	Broiler Caeca
	9801	Broiler Caeca
	14347	Other Poultry
	13719	Broiler Caeca
	11408	Parent Breeder Fecal

Table 4.1 cont. *flagellinA* short variable region (SVR) allele, sample number, and sample origin of *Campylobacter* spp. isolates employed in this investigation.

<i>FlaA</i> SVR Allele	Sample Number	Sample Origin
	11695	Parent Breeder Fecal
	11382	Parent Breeder Fecal
	12435	Parent Breeder Fecal
	12250	Parent Breeder Fecal
	5834	Retail Product
	9465	Retail Product
	13601	Retail Product
DAC	13262	Broiler Caeca
	14641	Wild Birds
	13280	Broiler Caeca
	13931	Parent Breeder Fecal
	13926	Parent Rearing Fecal
DAD	11623	Domestic Species
	13769	Human Isolate
	13783	Human Isolate
	12572	Other Poultry
	14118	Human Isolate
	12537	Other Poultry
	7358	Parent Breeder Fecal
	8889	Parent Breeder Fecal
	7571	Parent Breeder Fecal
	13147	Retail Product
	9522	Retail Product

Table 4.2. Genes identified as absent\* from *C. jejuni* isolate 14118 (high invasion) based on microarray hybridization analysis with *C. jejuni* 11168 (PMSRU).

Gene	Function
<i>cj1273c</i>	<i>rpoZ</i> :DNA-directed RNA polymerase subunit omega, promotes RNA polymerase assembly
<i>cj0860</i>	probable integral membrane protein, permeases of the drug/metabolite transporter (DMT) superfamily
<i>cj0567</i>	hypothetical protein
<i>cj0407</i>	<i>lgt</i> :prolipoprotein diacylglyceryl transferase, cell wall/membrane/envelope biogenesis
<i>cj0424</i>	putative acidic periplasmic protein, uncharacterized protein conserved in bacteria
<i>cj1331</i>	<i>ptmB</i> :acylneuraminate cytidylyltransferase, flagellin modification
<i>cj0668</i>	putative ATP /GTP-binding protein
<i>cj0753c</i>	<i>tonB3</i> :transport protein, links inner and outer membranes
<i>cj0797c</i>	hypothetical protein
<i>cj0436</i>	hypothetical protein
<i>cj0378c</i>	hypothetical protein

\*Gack values were  $\geq -0.05$ , indicative of absence relative to *C. jejuni* 11168

Table 4.3. Genes identified as absent\* from *C. jejuni* isolate 5116 (mid invasion) based on microarray hybridization analysis with *C. jejuni* 11168 (PMSRU).

Name	Function
<i>cj0412</i>	putative ATP /GTP binding protein
<i>cj0010c</i>	<i>rnhB</i> : ribonuclease HII, replication, recombination and repair
<i>cj0087</i>	<i>aspA</i> : aspartate ammonia-lyase,
<i>cj1249</i>	hypothetical protein
<i>cj1534c</i>	possible bacterioferritin, DNA-binding ferritin-like protein (oxidative damage protectant)
<i>cj1449c</i>	hypothetical protein
<i>cj0266c</i>	putative integral membrane protein
<i>cj0073c</i>	hypothetical protein
<i>cj1210</i>	putative integral membrane protein
<i>cj0065c</i>	<i>folk</i> : putative 2-amino-4-hydroxy-6-hydroxymethyl-dihydropteridine pyrophosphokinase
<i>cj1724c</i>	7-cyano-7-deazaguanine reductase
<i>cj0323</i>	hypothetical protein
<i>cj1587c</i>	multidrug transporter membrane component/ATP-binding component
<i>cj0802</i>	<i>cysS</i> : cysteinyl-tRNA synthetase
<i>cj0567</i>	hypothetical protein
<i>cj0262c</i>	putative methyl-accepting chemotaxis signal transduction protein
<i>cj1567c</i>	<i>nuoM</i> : NADH dehydrogenase I chain M
<i>cj1642</i>	hypothetical protein

\*Gack values were  $\geq -0.05$ , indicative of absence relative to *C. jejuni* 11168

Table 4.4. Genes identified as absent\* from *C. jejuni* isolate 8557 (mid invasion) based on microarray hybridization analysis with *C. jejuni* 11168 (PMSRU).

Name	Function
<i>cj0056c</i>	hypothetical protein
<i>cj1436c</i>	putative amino transferase
<i>cj0101</i>	<i>parB</i> : family protein, predicted transcriptional regulators

\*Gack values were  $\geq -0.05$ , indicative of absence relative to *C. jejuni* 11168

Table 4.5- Genes identified as absent\* from *C. jejuni* isolate 13262 (low invasion) based on microarray hybridization analysis with *C. jejuni* 11168 (PMSRU).

Name	Function
<i>cj0192c</i>	<i>clpP</i> : ATP-dependent Clp protease proteolytic subunit
<i>cj1186c</i>	<i>petA</i> : putative ubiquinol-cytochrome C reductase iron-sulfur subunit
<i>cj0894c</i>	<i>ispH</i> 4-hydroxy-3-methylbut-2-enyl diphosphate reductase, penicillin tolerance protein
<i>cj1152c</i>	putative phosphatase
<i>cj0567</i>	hypothetical protein
<i>cj0810</i>	<i>nadE</i> : putative NH(3)-dependent NAD(+) synthetase
<i>cj1060c</i>	small hydrophobic protein
<i>cj0737</i>	putative periplasmic protein, large exoproteins involved in heme utilization or adhesion
<i>cj1038</i>	probable cell division/peptidoglycan biosynthesis protein
<i>cj1435c</i>	hypothetical protein
<i>cj0407</i>	<i>Lgt</i> : prolipoprotein diacylglyceryl transferase
<i>cj1271c</i>	<i>tyrS</i> : tyrosyl-tRNA synthetase
<i>cj1255</i>	putative isomerase
<i>cj0267c</i>	putative integral membrane protein
<i>cj0224</i>	<i>argC</i> : N-acetyl-gamma-glutamyl-phosphate reductase
<i>cj0695</i>	<i>ftsA</i> : cell division protein
<i>cj1016c</i>	<i>livM</i> : putative branched-chain amino-acid ABC transport system permease protein
<i>cj0999c</i>	putative integral membrane protein
<i>cj1409</i>	<i>acpS</i> : 4'-phosphopantetheinyl transferase
<i>cj1567c</i>	<i>nuoM</i> : NADH dehydrogenase I chain M
<i>cj0063c</i>	putative ATP-binding protein
<i>cj0641</i>	hypothetical protein
<i>cj1243</i>	<i>hemE</i> : uroporphyrinogen decarboxylase
<i>cj0079c</i>	<i>cdtA</i> : cytolethal distending toxin
<i>cj0441</i>	<i>acpP</i> : acyl carrier protein
<i>cj0789</i>	putative RNA nucleotidyltransferase
<i>cj1379</i>	<i>selB</i> : putative selenocysteine-specific elongation factor
<i>cj0036</i>	hypothetical protein
<i>cj0366c</i>	transmembrane efflux protein
<i>cj0303c</i>	<i>modA</i> : putative molybdate-binding lipoprotein
<i>cj0825</i>	putative processing peptidase, Type II secretory pathway
<i>cj0188c</i>	hypothetical protein
<i>cj1181c</i>	<i>Tsf</i> : elongation factor Ts

\*Gack values were  $\geq -0.05$ , indicative of absence relative to *C. jejuni* 11168

Table 4.5 cont. Genes identified as absent\* from *C. jejuni* isolate 13262 (low invasion) based on microarray hybridization analysis with *C. jejuni* 11168 (PMSRU).

Name	Function
<i>cj0021c</i>	hypothetical protein
<i>cj0649</i>	hypothetical protein
<i>cj1486c</i>	Putative periplasmic protein
<i>cj1699c</i>	<i>rpmC</i> : 50S ribosomal protein L29
<i>cj1548c</i>	putative NADP-dependent alcohol dehydrogenase
<i>cj0172c</i>	hypothetical protein
<i>cj1133</i>	<i>waaC</i> : putative lipopolysaccharide heptosyltransferase
<i>cj1533c</i>	putative helix-turn-helix containing protein
<i>cj0717</i>	hypothetical protein
<i>cj1254</i>	hypothetical protein
<i>cj0055c</i>	hypothetical protein
<i>cj1689c</i>	<i>rplO</i> : 50S ribosomal protein L15
<i>cj0087</i>	<i>aspA</i> : aspartate ammonia-lyase
<i>cj1436c</i>	putative aminotransferase
<i>cj0017c</i>	putative ATP /GTP binding protein
<i>cj0070c</i>	hypothetical protein
<i>cj0805</i>	putative zinc protease
<i>cj0056c</i>	hypothetical protein
<i>cj1200</i>	putative periplasmic protein
<i>cj1069</i>	hypothetical protein
<i>cj0897c</i>	<i>pheS</i> : phenylalanyl-tRNA synthetase subunit alpha
<i>cj1449c</i>	hypothetical protein
<i>cj1636c</i>	<i>rnhA</i> : ribonuclease H
<i>cj0391c</i>	hypothetical protein
<i>cj0516</i>	<i>plsC</i> : putative 1-acyl-SN-glycerol-3-phosphate acyltransferase
<i>cj1491c</i>	putative two-component regulator
<i>cj0487</i>	hypothetical protein
<i>cj0461c</i>	putative integral membrane protein
<i>cj0724</i>	hypothetical protein
<i>cj0428</i>	hypothetical protein
<i>cj0352</i>	putative transmembrane protein
<i>cj0932c</i>	<i>pckA</i> : phosphoenolpyruvate carboxykinase
<i>cj1021c</i>	putative periplasmic protein

\*Gack values were  $\geq 0.05$ , indicative of absence relative to *C. jejuni* 11168



Table 4.6: Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
Amino acid transport and metabolism	
<i>cj0764c</i>	<i>speA</i> : arginine decarboxylase
<i>cj1018c</i>	branched-chain amino-acid ABC transport system periplasmic binding protein
<i>cj1502c</i>	<i>putP</i> : sodium/proline symporter
<i>cj0931c</i>	<i>argH</i> : argininosuccinate lyase
<i>cj1378</i>	<i>selA</i> : selenocysteine synthase
<i>cj0922c</i>	<i>pebC</i> : ABC-type amino-acid transporter ATP-binding protein
<i>cj1580c</i>	putative peptide ABC-transport system ATP-binding protein
<i>cj1286</i>	<i>upp</i> : uracil phosphoribosyltransferase
<i>cj0762c</i>	<i>aspB</i> : aspartate aminotransferase
<i>cj0481</i>	putative lyase
<i>cj0227</i>	<i>argD</i> : acetylornithine aminotransferase
<i>cj1315c</i>	<i>hisH</i> : imidazole glycerol phosphate synthase subunit HisH
<i>cj0314</i>	<i>lysA</i> : diaminopimelate decarboxylase
<i>cj1202</i>	<i>metF</i> : 5,10-methylenetetrahydrofolate reductase
<i>cj0317</i>	histidinol-phosphate aminotransferase
<i>cj0734c</i>	<i>hisJ</i> : histidine-binding protein precursor
<i>cj0349</i>	<i>trpA</i> : tryptophan synthase subunit alpha
<i>cj0716</i>	putative phospho-2-dehydro-3-deoxyheptonate aldolase
<i>cj1605c</i>	<i>dapD</i> : possible 2,3,4,5-tetrahydropyridine-2-carboxylate N-succinyltransferase
<i>cj0574</i>	<i>ilvI</i> : acetolactate synthase 3 catalytic subunit
<i>cj0197c</i>	<i>dapB</i> : dihydrodipicolinate reductase
<i>cj0817</i>	<i>glnH</i> : glutamine-binding periplasmic protein
<i>cj1014c</i>	<i>livF</i> : branched-chain amino-acid ABC transport system ATP-binding protein
<i>cj0980</i>	putative peptidase
<i>cj1624c</i>	<i>sdaA</i> : L-serine dehydratase
<i>cj1601</i>	<i>hisA</i> : phosphoribosylformimino-5-aminoimidazole carboxamide ribotide isomerase

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj0149c</i>	<i>hom</i> : homoserine dehydrogenase
<i>cj1598</i>	<i>hisD</i> : histidinol dehydrogenase
<i>cj0240c</i>	cysteine desulfurase
<i>cj0940c</i>	<i>glnP</i> : putative glutamine transport system permease
<i>cj0632</i>	<i>ilvC</i> : ketol-acid reductoisomerase
<i>cj0130</i>	<i>tyrA</i> : prephenate dehydrogenase
<i>cj1314c</i>	putative cyclase
<i>cj0921c</i>	bifunctional adhesin/ABC transporter aspartate/glutamate-binding protein
<i>cj0609c</i>	Possible periplasmic protein
<i>cj0665c</i>	<i>argG</i> : argininosuccinate synthase
Nucleotide transport and metabolism	
<i>cj1498c</i>	adenylosuccinate synthetase
<i>cj0353c</i>	phosphatase
<i>cj1195c</i>	<i>pyrC2</i> : dihydroorotase
<i>cj0953c</i>	<i>purH</i> : bifunctional phosphoribosylaminoimidazolecarboxamide formyltransferase/IMP cyclohydrolase
<i>cj0117</i>	pfs 5'-methylthioadenosine/S-adenosylhomocysteine nucleosidase
<i>cj0196c</i>	<i>ppurF</i> : amidophosphoribosyltransferase
<i>cj0419</i>	hypothetical protein
Carbohydrate transport and metabolism	
<i>cj0339</i>	putative transmembrane transport protein
<i>cj1588</i>	putative transmembrane transport protein
<i>cj0250c</i>	putative transmembrane transport protein
<i>cj1174</i>	putative efflux protein
<i>cj1588c</i>	putative transmembrane transport protein
<i>cj1619</i>	<i>kgtP</i> : alpha-ketoglutarate permease
<i>cj1401c</i>	<i>tpiA</i> : triosephosphate isomerase
<i>cj1645</i>	<i>tkt</i> : transketolase
<i>cj0128c</i>	suhB like protein
<i>cj0486</i>	putative sugar transporter

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj0392c</i>	<i>pyk</i> : pyruvate kinase
<i>cj1597</i>	<i>hisG</i> : ATP phosphoribosyltransferase
<i>cj1418c</i>	hypothetical protein
Cell cycle control, cell division, chromosome partitioning	
<i>cj0886c</i>	<i>ftsK</i> : putative cell division protein
<i>cj0696</i>	<i>ftsZ</i> : cell division protein FtsZ
<i>cj1606c</i>	<i>Mrp</i> : putative ATP/GTP-binding protein (mrp protein homolog)
Cell Motility	
<i>cj1729c</i>	<i>flgE</i> : flagellar hook protein
<i>cj0043</i>	<i>FlgE</i> : flagellar hook protein
<i>cj0528c</i>	<i>flgB</i> : flagellar basal body rod protein
<i>cj1338c</i>	<i>flaB</i> : flagellin
<i>cj1675</i>	<i>fliQ</i> : flagellar biosynthesis protein
<i>cj0318</i>	<i>fliF</i> : flagellar MS-ring protein
<i>cj0060c</i>	<i>fliM</i> : flagellar motor switch protein
<i>cj0059c</i>	<i>fliY</i> : flagellar motor switch protein
<i>cj0882c</i>	<i>flhA</i> : flagellar biosynthesis protein A
<i>cj0887c</i>	<i>flaD</i> : flagellar hook-associated protein
<i>cj0526c</i>	<i>fliE</i> : flagellar hook-basal body protein
<i>cj0283c</i>	<i>cheW</i> : chemotaxis protein
<i>cj0064c</i>	<i>flhF</i> : flagellar biosynthesis regulator FlhF
<i>cj0697</i>	<i>flgG2</i> : putative flagellar basal-body rod protein
<i>cj1190c</i>	putative MCP-domain signal transduction protein
<i>cj0924c</i>	<i>cheB</i> : putative MCP protein-glutamate methyltransferase
<i>cj1474c</i>	putative type II protein secretion system D protein
<i>cj1471c</i>	putative type II protein secretion system E protein
<i>cj0246c</i>	putative MCP-domain signal transduction protein

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj1343c</i>	putative periplasmic protein
Cell Wall/membrane/envelope biogenesis	
<i>cj0910</i>	putative periplasmic protein
<i>cj0131</i>	putative periplasmic protein
<i>cj0735</i>	putative periplasmic protein
<i>cj1406c</i>	putative periplasmic protein
<i>cj0129c</i>	outer membrane protein
<i>cj1670c</i>	putative periplasmic protein
<i>cj1621</i>	putative periplasmic protein
<i>cj1513c</i>	possible periplasmic protein
<i>cj0365c</i>	putative outer membrane channel protein
<i>cj0367c</i>	putative membrane fusion component of efflux system
<i>cj0162c</i>	putative periplasmic protein
<i>cj1416c</i>	putative sugar nucleotidyltransferase
<i>cj1289</i>	possible periplasmic protein
<i>cj1485c</i>	putative periplasmic protein
<i>cj0413</i>	putative periplasmic protein
<i>cj0770c</i>	putative periplasmic protein
<i>cj1428c</i>	<i>fcl</i> : putative fucose synthetase
<i>cj1142</i>	<i>neuC1</i> : putative N-acetylglucosamine-6-phosphate 2-epimerase/N-acetylglucosamine-6-phosphatase
<i>cj0511</i>	putative secreted protease
<i>cj1151c</i>	<i>waaD</i> : ADP-L-glycero-D-manno-heptose-6-epimerase
<i>cj1131c</i>	UDP-glucose 4-epimerase
<i>cj1317</i>	N-acetylneuraminic acid synthetase
<i>cj0821</i>	glmU UDP-N-acetylglucosamine pyrophosphorylase
<i>cj0645</i>	putative secreted transglycosylase

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj0576</i>	<i>lpxD</i> : UDP-3-O-[3-hydroxymyristoyl] glucosamine N-acyltransferase
<i>cj1294</i>	putative aminotransferase (degT family)
<i>cj1128</i>	<i>wlaD</i> : putative glycosyltransferase
<i>cj1311</i>	acylneuraminate cytidyltransferase
<i>cj0611c</i>	putative transmembrane transport protein
<i>cj1055c</i>	putative integral membrane protein
<i>cj0238</i>	putative integral membrane protein
<i>cj0946</i>	putative lipoprotein
<i>cj0361</i>	<i>lspA</i> : lipoprotein signal peptidase
<i>cj1030c</i>	<i>lepA</i> : GTP-binding protein LepA
<i>cj1086c</i>	hypothetical protein
Coenzyme transport and metabolism	
<i>cj0542</i>	<i>hemA</i> : glutamyl-tRNA reductase
<i>cj0857c</i>	<i>moeA</i> : putative molybdopterin biosynthesis protein
<i>cj0580c</i>	coproporphyrinogen III oxidase
<i>cj0580c</i>	coproporphyrinogen III oxidase
<i>cj0853c</i>	<i>hemL</i> : glutamate-1-semialdehyde aminotransferase
<i>cj0725c</i>	<i>mogA</i> : molybdenum cofactor biosynthesis protein
<i>cj1218c</i>	<i>ribA</i> : riboflavin synthase subunit alpha
<i>cj1046c</i>	thiamine biosynthesis protein ThiF
<i>cj1239</i>	<i>pdxA</i> : 4-hydroxythreonine-4-phosphate dehydrogenase
<i>cj0306c</i>	<i>bioF</i> : 8-amino-7-oxononanoate synthase
<i>cj0589</i>	<i>ribF</i> : bifunctional riboflavin kinase/FMN adenylyltransferase
<i>cj0308c</i>	<i>bioD</i> : putative dethiobiotin synthetase

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj0585</i>	<i>folP</i> : putative dihydropteroate synthase
<i>cj0453</i>	<i>thiC</i> : thiamine biosynthesis protein ThiC
<i>cj0383c</i>	<i>ribH</i> : riboflavin synthase subunit beta
<i>cj1458c</i>	<i>thiL</i> : thiamine monophosphate kinase
<i>cj0230c</i>	nicotinate phosphoribosyltransferase
<i>cj1368</i>	hypothetical protein
<i>cj1404</i>	hypothetical protein
<i>cj1047c</i>	hypothetical protein
Defense mechanism	
<i>cj0077c</i>	<i>cdtC</i> : cytolethal distending toxin
<i>cj0205</i>	<i>bacA</i> : putative undecaprenol kinase
<i>cj0619</i>	putative integral membrane protein
<i>cj0140</i>	hypothetical protein
Energy production and conversion	
<i>cj1382c</i>	<i>fldA</i> : flavodoxin FldA
<i>cj1265c</i>	<i>hydC</i> : Ni/Fe-hydrogenase B-type cytochrome subunit
<i>cj1066</i>	<i>rdxA</i> : nitroreductase
<i>cj0537</i>	<i>oorB</i> : 2-oxoglutarate-acceptor oxidoreductase subunit OorB
<i>cj0439</i>	<i>sdhC</i> : putative succinate dehydrogenase subunit C
<i>cj0076c</i>	L-lactate permease
<i>cj0333c</i>	<i>fdxA</i> : ferredoxin
<i>cj0409</i>	fumarate reductase
<i>cj1586</i>	putative bacterial haemoglobin
<i>cj0074c</i>	putative iron-sulfur protein
<i>cj0991c</i>	putative oxidoreductase ferredoxin-type electron transport protein
<i>cj1488c</i>	<i>ccoQ</i> : cb-type cytochrome C oxidase subunit IV
<i>cj0781</i>	<i>napG</i> : quinol dehydrogenase periplasmic component
<i>cj1570c</i>	<i>nuoJ</i> : NADH dehydrogenase subunit J

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj1578c</i>	<i>nuoB</i> : NADH dehydrogenase subunit B
<i>cj0783</i>	<i>napB</i> : periplasmic nitrate reductase small subunit
<i>cj0265c</i>	putative cytochrome C-type haem-binding periplasmic protein
<i>cj1267c</i>	Ni/Fe-hydrogenase small chain
<i>cj0104</i>	<i>atpH</i> : F0F1 ATP synthase subunit delta
<i>cj1192</i>	putative C4-dicarboxylate transport protein
<i>cj1167</i>	<i>ldh</i> : putative L-lactate dehydrogenase
<i>cj0532</i>	malate dehydrogenase
<i>cj0780</i>	<i>napA</i> : periplasmic nitrate reductase
<i>cj1399c</i>	putative Ni/Fe-hydrogenase small subunit
<i>cj1153</i>	putative periplasmic cytochrome C
<i>cj0203</i>	putative transmembrane protein
Intercellular trafficking, secretion and vesicular transport	
<i>cj0986c</i>	putative integral membrane protein
<i>cj0530</i>	putative periplasmic protein
<i>cj0851c</i>	putative integral membrane protein
<i>cj0852c</i>	putative integral membrane protein
<i>cj0587</i>	putative integral membrane protein
<i>cj1092c</i>	<i>secF</i> : preprotein translocase subunit <i>SecF</i>
<i>cj1206c</i>	<i>ftsY</i> : putative signal recognition particle protein
<i>cj0110</i>	<i>exbD</i> : tolR family transport protein
<i>cj0579c</i>	sec-independent translocase
<i>cj0472</i>	<i>secE</i> preprotein translocase subunit <i>SecE</i>
<i>cj0578c</i>	sec-independent protein translocase
Lipid transport and metabolism	
<i>cj0375</i>	putative lipoprotein

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj1090c</i>	putative lipoprotein
<i>cj0176c</i>	putative lipoprotein
<i>cj0158c</i>	putative haem-binding lipoprotein
<i>cj0978c</i>	putative lipoprotein
<i>cj1299</i>	putative acyl carrier protein
<i>cj1029c</i>	<i>mapA</i> : putative lipoprotein
<i>cj1279c</i>	putative fibronectin domain-containing lipoprotein
<i>cj0842</i>	putative lipoprotein
<i>cj1346c</i>	<i>dxr</i> : 1-deoxy-D-xylulose 5-phosphate reductoisomerase
<i>cj1665</i>	possible lipoprotein thioredoxin
<i>cj1026c</i>	putative lipoprotein
<i>cj1104</i>	4-diphosphocytidyl-2-C-methyl-D-erythritol kinase
<i>cj1347c</i>	<i>cdsA</i> : phosphatidate cytidyltransferase
<i>cj0329c</i>	<i>plsX</i> : fatty acid/phospholipid synthesis protein
<i>cj1037c</i>	<i>pycA</i> : acetyl-CoA carboxylase
<i>cj0182</i>	putative transporter
Posttranslational modification, protein turnover, and chaperones	
<i>cj0759</i>	<i>dnaK</i> : molecular chaperone DnaK
<i>cj1207c</i>	putative lipoprotein thioredoxin
<i>cj0193c</i>	<i>tig</i> : trigger factor
<i>cj1112c</i>	methionine sulfoxide reductase B
<i>cj0623</i>	<i>hypB</i> : hydrogenase isoenzyme formation protein
<i>cj0596</i>	<i>peb4:cbf2</i> peptidyl-prolyl cis-trans isomerase
<i>cj1106</i>	possible periplasmic thioredoxin
<i>cj0954c</i>	putative dnaJ-like protein
<i>cj0622</i>	<i>hypF</i> : transcriptional regulatory protein hypF



Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj1639</i>	<i>nifU</i> : protein homolog
<i>cj0701</i>	putative protease
<i>cj0133</i>	hypothetical
<i>cj0625</i>	<i>hypD</i> : hydrogenase isoenzyme formation protein
<i>cj1035c</i>	arginyl-tRNA-protein transferase
<i>cj1034c</i>	possible dnaJ-like protein
<i>cj1289</i>	possible periplasmic protein
<i>cj0950c</i>	putative lipoprotein
Signal transduction mechanisms	
<i>cj1024c</i>	signal-transduction regulatory protein
<i>cj0890c</i>	putative sensory transduction transcriptional regulator
<i>cj0248</i>	hypothetical protein
<i>cj0643</i>	putative two-component response regulator
<i>cj1261</i>	two-component regulator
<i>cj1222c</i>	putative two-component sensor
Replication, recombination and repair	
<i>cj0718</i>	<i>dnaE</i> : DNA polymerase III subunit alpha
<i>cj0002</i>	<i>dnaN</i> : DNA polymerase III subunit beta
<i>cj0464</i>	<i>recG</i> : ATP-dependent DNA helicase
<i>cj1157</i>	DNA polymerase III subunits gamma and tau
<i>cj0003</i>	<i>gyrB</i> : DNA gyrase subunit B
<i>cj0001</i>	<i>dnaA</i> : chromosomal replication initiation protein
<i>cj0338c</i>	<i>polA</i> : DNA polymerase I
<i>cj0198c</i>	recombination factor protein <i>RarA</i>
<i>cj0836</i>	<i>ogt</i> : methylated-DNA--protein-cysteine methyltransferase
<i>cj0595c</i>	endonuclease III
<i>cj0680c</i>	<i>uvrB</i> : excinuclease ABC subunit B

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<b>Transcription</b>	
<i>cj1156</i>	<i>rho</i> : transcription termination factor Rho
<i>cj0368c</i>	transcriptional regulatory protein
<i>cj1563c</i>	putative transcriptional regulator
<i>cj1230</i>	<i>hspR</i> : putative heat shock transcriptional regulator
<i>cj0394c</i>	pantothenate kinase
<i>cj1635c</i>	<i>rnc</i> : ribonuclease III
<i>cj0478</i>	<i>rpoB</i> : DNA-directed RNA polymerase subunit beta
<i>cj0883c</i>	hypothetical protein
<b>Translation ribosomal structure and biogenesis</b>	
<i>cj0207</i>	<i>infC</i> : translation initiation factor IF-3
<i>cj0094</i>	<i>rplU</i> : 50S ribosomal protein L21
<i>cj0474</i>	<i>rplK</i> : 50S ribosomal protein L11
<i>cj1182c</i>	<i>rpsB</i> : 30S ribosomal protein S2
<i>cj1592</i>	<i>rpsM</i> : 30S ribosomal protein S13
<i>cj1694c</i>	<i>rpsN</i> : 30S ribosomal protein S14
<i>cj1701c</i>	<i>rpsC</i> : 30S ribosomal protein S3
<i>cj0884</i>	<i>rpsO</i> : 30S ribosomal protein S15
<i>cj1702c</i>	<i>rplV</i> : 50S ribosomal protein L22
<i>cj1696c</i>	<i>rplX</i> : 50S ribosomal protein L24
<i>cj0370</i>	<i>rpsU</i> : 30S ribosomal protein S21
<i>cj1692c</i>	<i>rplF</i> : 50S ribosomal protein L6
<i>cj0476</i>	<i>rplJ</i> : 50S ribosomal protein L10
<i>cj1697c</i>	<i>rplN</i> : 50S ribosomal protein L14
<i>cj0640c</i>	<i>aspS</i> : aspartyl-tRNA synthetase
<i>cj0765c</i>	<i>hisS</i> : histidyl-tRNA synthetase
<i>cj0845c</i>	<i>gltX</i> : glutamyl-tRNA synthetase
<i>cj0577c</i>	<i>queA</i> : S-adenosylmethionine:tRNA ribosyltransferase-isomerase
<i>cj0153c</i>	putative rRNA methylase

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj0636</i>	NOL1\NOP2\sun family protein
<i>cj0166</i>	<i>miaA</i> : tRNA delta(2)-isopentenylpyrophosphate transferase
<i>cj0930</i>	translation-associated GTPase
<i>cj0588</i>	<i>tlyA</i> : putative haemolysin
<i>cj0879c</i>	putative periplasmic protein
<b>Inorganic ion transport and metabolism</b>	
<i>cj0755</i>	<i>cfrA</i> : putative iron uptake protein
<i>cj1354</i>	enterochelin uptake ATP-binding protein
<i>cj0237</i>	carbonic anhydrase
<i>cj1194</i>	possible phosphate permease
<i>cj0614</i>	<i>pstC</i> : putative phosphate transport system permease protein
<i>cj0785</i>	possible <i>napD</i> protein homolog
<i>cj1284</i>	<i>ktrA</i> : putative K <sup>+</sup> uptake protein
<i>cj1283</i>	<i>ktrB</i> : putative K <sup>+</sup> uptake protein
<i>cj1398</i>	<i>feoB</i> : ferrous iron transport protein
<i>cj0612c</i>	<i>cft</i> : ferritin
<i>cj1615</i>	putative haemin uptake system permease protein
<i>cj0616</i>	putative phosphate transport ATP-binding protein
<i>cj0263</i>	zinc transporter <i>ZupT</i>
<i>cj0020c</i>	cytochrome C551 peroxidase
<i>cj0045c</i>	putative iron-binding protein
<i>cj1617</i>	<i>chuD</i> : putative haemin uptake system periplasmic haemin-binding protein
<i>cj1614</i>	haemin uptake system outer membrane receptor
<i>cj0142c</i>	ABC transporter ATP-binding protein
<i>cj0141c</i>	ABC transporter integral membrane protein
<i>cj0169</i>	<i>sodB</i> : superoxide dismutase
<i>cj0772c</i>	putative periplasmic protein
<i>cj0613</i>	<i>pstS</i> : possible periplasmic phosphate binding protein
<i>cj0770c</i>	putative periplasmic protein
<i>cj1658</i>	putative integral membrane protein

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj0241c</i>	putative iron-binding protein
Secondary metabolite biosynthesis and transport	
<i>cj0261c</i>	hypothetical protein
<i>cj0977</i>	hypothetical protein
<i>cj0590</i>	hypothetical protein
General Function prediction only	
<i>cj0572</i>	<i>ribA</i> : bifunctional 3,4-dihydroxy-2-butanone 4-phosphate synthase/GTP cyclohydrolase II protein
<i>cj0431</i>	putative periplasmic ATP /GTP-binding protein
<i>cj1041c</i>	putative periplasmic ATP/GTP-binding protein
<i>cj1159c</i>	small hydrophobic protein
<i>cj1545c</i>	<i>MdaB</i> : protein homolog
<i>cj0834c</i>	ankyrin repeat-containing possible periplasmic protein
<i>cj0154c</i>	putative methylase
<i>cj1278c</i>	<i>trmB</i> : tRNA (guanine-N(7))-methyltransferase
<i>cj1321</i>	putative transferase
<i>cj0778</i>	major antigenic peptide PEB2
<i>cj0947c</i>	putative hydrolase
<i>cj0985c</i>	hippurate hydrolase
<i>cj0900c</i>	small hydrophobic protein
<i>cj0251c</i>	highly acidic protein
<i>cj0556</i>	hypothetical protein
<i>cj1270c</i>	hypothetical protein
<i>cj0760</i>	hypothetical protein
<i>cj0465c</i>	hypothetical protein
<i>cj1225</i>	hypothetical protein
<i>cj0647</i>	hypothetical protein
<i>cj1307</i>	putative amino acid activating enzyme
<i>cj0935c</i>	putative transmembrane transport protein
<i>cj0183</i>	putative integral membrane protein with haemolysin domain

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj1373</i>	putative integral membrane protein
<i>cj0846</i>	putative integral membrane protein
<i>cj0091</i>	putative lipoprotein
<i>cj0413</i>	putative periplasmic protein
Unknown function	
<i>cj0993c</i>	hypothetical protein
<i>cj0254</i>	hypothetical protein
<i>cj0794</i>	hypothetical protein
<i>cj1475c</i>	hypothetical protein
<i>cj1209</i>	hypothetical protein
<i>cj0138</i>	hypothetical protein
<i>cj0796c</i>	hypothetical protein
<i>cj0569</i>	hypothetical protein
<i>cj1144c</i>	hypothetical protein
<i>cj1162c</i>	hypothetical protein
<i>cj0877c</i>	hypothetical protein
<i>cj0494</i>	hypothetical protein
<i>cj0403</i>	hypothetical protein
<i>cj0286c</i>	hypothetical protein
<i>cj1232</i>	hypothetical protein
<i>cj1484c</i>	hypothetical protein
<i>cj1214c</i>	hypothetical protein
<i>cj1245c</i>	hypothetical protein
<i>cj1176c</i>	hypothetical protein
<i>cj1562</i>	hypothetical protein
<i>cj0522</i>	hypothetical protein
<i>cj0959c</i>	hypothetical protein
<i>cj1012c</i>	hypothetical protein
<i>cj1631c</i>	hypothetical protein
<i>cj0963</i>	hypothetical protein

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj1006c</i>	hypothetical protein
<i>cj0700</i>	hypothetical protein
<i>cj1467</i>	hypothetical protein
<i>cj0189c</i>	hypothetical protein
<i>cj1453c</i>	hypothetical protein
<i>cj0247c</i>	hypothetical protein
<i>cj0815</i>	hypothetical protein
<i>cj0598</i>	hypothetical protein
<i>cj1405</i>	hypothetical protein
<i>cj1575c</i>	hypothetical protein
<i>cj1465</i>	hypothetical protein
<i>cj0800c</i>	hypothetical protein
<i>cj0041</i>	hypothetical protein
<i>cj1236</i>	hypothetical protein
<i>cj0418</i>	hypothetical protein
<i>cj0455c</i>	hypothetical protein
<i>cj1656c</i>	hypothetical protein
<i>cj0583</i>	hypothetical protein
<i>cj1384c</i>	hypothetical protein
<i>cj0849c</i>	hypothetical protein
<i>cj0563</i>	hypothetical protein
<i>cj1089c</i>	hypothetical protein
<i>cj0550</i>	hypothetical protein
<i>cj0302c</i>	hypothetical protein
<i>cj0873c</i>	hypothetical protein
<i>cj1443c</i>	<i>KpsF</i> : protein
<i>cj0552</i>	hyprophobic protein

Table 4.6 cont. Genes determined to be commonly distributed throughout *C. jejuni* isolates 14118, 5116, 8557 and 13262 based on microarray hybridization with *C. jejuni* 11168 (PMSRU).

Gene ID	Description
<i>cj0610c</i>	putative periplasmic protein
<i>cj0593c</i>	putative integral membrane protein
<i>cj0204</i>	putative integral membrane protein
<i>cj0553</i>	putative integral membrane protein
<i>cj1166c</i>	putative integral membrane protein
<i>cj0014c</i>	putative integral membrane protein
<i>cj1022c</i>	putative integral membrane protein

Table 4.7. Total number of inserts and clones provided for sequence analysis along with percentage of clones with sequences absent from *C. jejuni* 11168 (PMSRU) based on suppressive subtractive hybridizations.

Isolate	# of inserts analyzed	# of subtracted clones provided for DNA sequence analysis	% of clones with sequences determined unique relative to 11168 (PMSRU)
13262	94	59	63%
8557	141	79	56%
14118	103	68	66%
5116	33	24	72%



Table 4.8. Isolate 14118 (Caco-2 high invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
Cell Envelop & Surface structures						
37	343	32.4	a) LOS gene locus, partial sequence b) unknown	<i>C. jejuni</i> strain LC <i>C. jejuni</i>	266/266 (100%) 68/68 (100%)	gb DQ535892.1 gb ABZ79836.1
45	221	33.5	a) putative outer-membrane protein b) putative outer-membrane protein	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	130/131 (99%) 43/48 (89%)	gb CP000814.1 ref YP_001482042.1
48	492	36.4	a) no significant similarity to any nucleic acid b) putative sugar transferase,	<i>C. jejuni</i>	48/73 (65%)	emb CAI38725.1
57	247	35.2	a) class H lipooligosaccharide biosynthesis gene locus, partial sequence b) unknown	<i>C. jejuni</i> strain RM1553	165/165 (100%)	gb EU404106.1
92	322	42.8	a) <i>flaA</i> and <i>flaB</i> genes  b) <i>FlaB</i>	<i>C. jejuni</i> <i>C. jejuni</i> TGH9011(ATCC43431) <i>C. jejuni</i>	54/54 (100%) 221/241 (91%)  79/80 (98%)	gb ABZ79829.1 Z29327.1  gb ABS89177.1
130	470	38.1	a) putative integral membrane protein b) putative integral membrane protein	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	209/210 (99%) 46/47 (97%)	gb CP000814.1 ref YP_001481584.1
132	308	35.7	a) no significant similarity to any nucleic acid b) putative glycosyltransferase,	<i>C. jejuni</i>	17/70 (24%)	b AAR98510.1
150	253	33.5	a) no significant similarity to any nucleic acid b) motility accessory factor	<i>C. doylei</i> 269.97	39/51 (76%)	ref YP_001397577.1
163	297	37.3	a) putative integral membrane protein, b) putative integral membrane protein,	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	221/221 (100%) 73/74 (98%)	gb CP000814.1 ref YP_001481894.1
179	297	36.3	a) 324 bp at 5' side: transformation system protein, 738 bp at 3' side: motility accessory factor  b) hypothetical protein C8J_1258	<i>C. doylei</i> 269.97  <i>C. jejuni</i> 81116	185/199 (92%)  62/70 (88%)	gb CP000768.1  ref YP_001482834.1
191	375	33.0	a) class O lipooligosaccharide biosynthesis gene locus, partial sequence b) putative dTDP-glucose 4,6-dehydratase	<i>C. jejuni</i> RM3423  <i>C. jejuni</i>	294/298 (98%)  76/79 (96%)	gb EF143352.1  gb ABN41486.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.8 cont. Isolate 14118 (Caco-2 high invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
197	527	40.9	a) class J lipooligosaccharide biosynthesis gene locus	<i>C. jejuni</i> RM1508	224/236 (94%)	gb EU404104.1
209	249	39.7	b) hypothetical protein C8J_1345 a) class S lipooligosaccharide biosynthesis gene locus, partial sequence	<i>C. jejuni</i> 81116 <i>C. jejuni</i> RM3419	78/85 (91%) 156/158 (98%)	YP_001482920.1 gb EU404110.1
229	315	32.6	b) unknown a) class H lipooligosaccharide biosynthesis gene locus	<i>C. jejuni</i> <i>C. jejuni</i> RM1553	40/44 (90%) 237/239 (99%)	gb ABZ79851.1 gb EU404106.1
241	281	33.4	b) unknown a) LOS biosynthesis cluster	<i>C. jejuni</i> <i>C. jejuni</i> 11828	78/80 (97%) 191/191 (100%)	gb ABZ79837.1 gb AF343914.1
249	274	43.5	b) hypothetical protein C8J_1094 a) flagellin A ( <i>flaA</i> ) gene	<i>C. jejuni</i> 81116 <i>C. jejuni</i> D5477	63/63 (100%) 148/163 (90%)	YP_001482670.1 gb AF369587.1
271	168	48.8	b) flagellin A a) putative periplasmic protein	<i>C. jejuni</i> HB93-13 <i>C. jejuni</i> 81-176	57/61 (93%) 89/92 (96%)	ZP_01071151.1 gb CP000538.1
290	268	37.6	b) putative periplasmic protein a) integral membrane protein gene b) integral membrane protein	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> <i>C. jejuni</i>	29/32 (90%) 190/190 (100%) 62/67 (92%)	YP_001000654.1 gb AF273109.1 gb AAF82114.1
Restriction-modification, recombination & repair						
26	354	33.6	a) type I restriction-modification system, M subunit b) type I restriction-modification system, M subunit	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	261/264 (98%) 87/91 (95%)	gb CP000538.1 YP_001000444.1
91	294	33.3	a) hypothetical protein b) <i>RloA</i>	<i>C. jejuni</i> 81116 <i>C. jejuni</i>	202/204 (99%) 40/42 (95%)	gb CP000814.1 gb AAN33168.1
135	392	33.2	a) <i>RloB</i> b) <i>RloB</i>	<i>C. jejuni</i> 81116 <i>C. jejuni</i> CG8486	316/317 (99%) 105/106 (99%)	gb CP000814.1 ZP_01809391.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.8 cont. Isolate 14118 (Caco-2 high invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
173	361	34.9	a) putative McrBC restriction endonuclease system, McrB subunit b) McrBC restriction endonuclease system, McrB subunit, putative	<i>C. doylei</i> 269.97 <i>C. jejuni</i> HB93-13	211/224 (94%) 82/95 (86%)	gb CP000768.1 ZP_01072052.1
215	320	36.3	a) <i>HsdR</i> pseudogene, hsdR-1 allele, complete sequence; <i>RloG</i> gene, <i>HsdS</i> pseudogene, hsdS-5 allele, and <i>HsdM</i> gene, hsdM-1 allele b) <i>HsdM</i>	<i>C. jejuni</i> RM1167 <i>C. jejuni</i>	230/233 (98%) 77/77 (100%)	gb AF486547.1 gb AAM00833.1
224	451	35.6	a) hypothetical protein b) <i>RloA</i>	<i>C. jejuni</i> 81116 <i>C. jejuni</i> CG8486	279/294 (94%) 87/99 (87%)	gb CP000814.1 ZP_01809390.1
291	399	33.3	a) <i>HsdR</i> gene, hsdR-1 allele, <i>RloA</i> and <i>RloB</i> genes, <i>HsdS</i> gene, hsdS-1 allele, and <i>HsdM</i> gene, hsdM-1 allele b) putative type I specificity subunit <i>HsdS</i>	<i>C. jejuni</i> RM1861 <i>C. jejuni</i> CG8486	262/262 (100%) 65/65 (100%)	gb AF486553.1 ZP_01809392.1
Transport						
136	403	40.2	a) di-/tripeptide transporter b) di-/tripeptide transporter	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	312/313 (99%) 106/115 (92%)	gb CP000814.1 YP_001482189.1
225	173	41.6	a) di-/tripeptide transporter b) di-/tripeptide transporter	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	118/118 (100%) 39/39 (100%)	gb CP000814.1 YP_001482189.1
240	119	36.1	a) Na <sup>+</sup> /H <sup>+</sup> antiporter b) no significant similarity to any protein	<i>C. jejuni</i> 81116	41/41 (100%)	gb CP000814.1
281	102	39.2	a) major facilitator superfamily protein b) major facilitator superfamily protein,	<i>C. doylei</i> 269.97 <i>C. doylei</i> 269.97	46/46 (100%) 15/15 (100%)	gb CP000768.1 YP_001397475.1
Chemotaxis						
145	412	37.1	a) methyl-accepting chemotaxis protein, b) putative MCP-type signal transduction protein	<i>C. jejuni</i> 81116 <i>C. jejuni</i>	334/336 (99%) 102/104 (98%)	gb CP000814.1 ZP_01809677.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.8 cont. Isolate 14118 (Caco-2 high invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
159	432	35.4	a) methyl-accepting chemotaxis protein, b) methyl-accepting chemotaxis protein	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	355/355 (100%) 110/110 (100%)	gb CP000814.1 YP_001482984.1
Other (bacteriophage sequence)						
234	343	33.8	a) prophage Lp2 protein 6 b) prophage Lp2 protein 6	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	255/256 (99%) 85/87 (97%)	gb CP000538.1 ZP_02271034.1
Small molecular metabolism						
82	390	36.9	a) hypothetical protein b) cytochrome c biogenesis protein	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81-176	185/185 (100%) 61/61 (100%)	gb CP000814.1 YP_999754.1
30	613	33.5	a), hypothetical protein b) lipoprotein, putative	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	371/371 (100%) 47/48 (97%)	gb CP000814.YP_001481975.1
148	644	34.2	a) hypothetical protein b) putative subunit of dimethyl sulfoxide reductase	<i>C. jejuni</i> 81-176 <i>C. jejuni</i>	561/569 (98%) 113/116 (97%)	gb CP000538.1 gb AAY53800.1
154	228	41.6	a) Ser/Thr protein phosphatase family protein b) Ser/Thr protein phosphatase family protein	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	139/140 (99%) 45/47 (95%)	gb CP000814.1 YP_001482369.1
172	347	39.5	a) anaerobic dimethyl sulfoxide reductase chain A b) hypothetical protein C8J_1482	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81116	256/259 (98%) 86/86 (100%)	gb CP000538.1 YP_001483057.1
203	351	34.1	a) hydrolase, carbon-nitrogen family b) hydrolase, carbon-nitrogen family	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> RM1221	261/261 (100%) 72/75 (96%)	gb CP000025.1 YP_179189.1
264	467	32.3	a) oxidoreductase, molybdopterin binding, putative orotidine 5'-phosphate decarboxylase b) orotidine 5'-phosphate decarboxylase,	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	380/380 (100%) 91/92 (98%)	gb CP000538.1 gb EAQ73091.1
267	486	37.0	a) arylsulfate sulfotransferase b) arylsulfate sulfotransferase, degenerate	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	401/405 (99%) 134/138 (97%)	gb CP000538.1 YP_001000550.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.8 cont. Isolate 14118 (Caco-2 high invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
272	209	40.6	a) 2-isopropylmalate synthase b) 2-isopropylmalate synthase	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	113/113 (100%) 37/37 (100%)	gb CP000814.1 YP_001483199.1
279	97	40.0	a) <i>RlfA</i> b) no significant similarity to any protein	<i>C. jejuni</i> 81116	41/41 (100%)	gb CP000814.1
Hypothetical & unknown						
25	239	33.9	a) hypothetical protein b) hypothetical protein C8J_0526	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	161/162 (99%) 52/55 (94%)	gb CP000814.1 YP_001482102.1
28	281	36.2	a) hypothetical protein b) hypothetical protein C8J_0400	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	189/190 (99%) 60/66 (90%)	gb CP000814.1 YP_001481976.1
40	401	31.7	a) 318 bp at 5' side: ATP synthase F0 sector C subunit, 798 bp at 3' side: hypothetical protein b) hypothetical protein <i>C. jejuni</i> _04900	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81-176	311/312 (99%) 70/70 (100%)	gb CP000814.1 ZP_02271300.1
47	695	29.7	a) hypothetical protein b) hypothetical protein C8J_0035	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	277/277 (100%) 92/92 (100%)	gb CP000814.1 YP_001481613.1
54	232	43.1	a), hypothetical protein b) hypothetical protein C8J_1589	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	155/156 (99%) 50/52 (96%)	gb CP000814.1 YP_001483163.1
67	381	35.6	a) hypothetical protein b) hypothetical protein C8J_0648	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	291/291 (100%) 86/87 (98%)	gb CP000814.1 YP_001482224.1
78	438	38.1	a) hypothetical protein b) hypothetical protein C8J_0034	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	314/328 (95%) 97/109 (88%)	gb CP000814.1 YP_001481612.1
122	307	35.1	a) hypothetical protein b) hypothetical protein C8J_0036	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	231/231 (100%) 77/79 (97%)	b CP000814.1 YP_001481614.1
134	716	34.9	a) hypothetical protein b) hypothetical protein C8J_0988	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	640/641 (99%) 213/213 (100%)	gb CP000814.1 YP_001482564.1
147	408	45.8	a) hypothetical protein b) hypothetical protein C8J_0878	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	284/287 (98%) 92/95 (96%)	gb CP000814.1 YP_001482454.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.8 cont. Isolate 14118 (Caco-2 high invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
153	386	38.8	a) hypothetical protein	<i>C. jejuni</i> 81-176	259/304 (85%)	gb CP000538.1
			b) hypothetical protein CJE1531	<i>C. jejuni</i> RM1221	82/99 (82%)	YP_179516.1
155	264	42.4	a) hypothetical protein	<i>C. jejuni</i> 81116	171/171 (100%)	gb CP000814.1
			b) hypothetical protein C8J_0986	<i>C. jejuni</i> 81116	56/57 (98%)	YP_001482562.1
170	268	41.0	a) hypothetical protein	<i>C. jejuni</i> 81116	172/172 (100%)	gb CP000814.1
			b) hypothetical protein C8J_0988	<i>C. jejuni</i> 81116	60/64 (93%)	YP_001482564.1
187	329	41.3	a) hypothetical protein	<i>C. jejuni</i> 81116	216/253 (85%)	gb CP000814.1
			b) hypothetical protein Cjejjeuni_07040	<i>C. jejuni</i> 81-176	76/84 (90%)	ZP_02271659.1
188	320	33.1	a) hypothetical protein	<i>C. jejuni</i> 81116	230/234 (98%)	gb CP000814.1
			b) hypothetical protein C8J_1619	<i>C. jejuni</i> 81116	62/62 (100%)	YP_001483193.1
200	362	38.7	a) hypothetical protein	<i>C. jejuni</i> 81116	283/285 (99%)	Gb CP000814.1
			b) hypothetical protein C8J_0142	<i>C. jejuni</i> 81116	79/81 (97%)	YP_001481718.1
208	507	39.2	a) hypothetical protein	<i>C. jejuni</i> 81116	416/417 (99%)	gb CP000814.1
			b) hypothetical protein C8J_0140	<i>C. jejuni</i> 81116	138/139 (99%)	YP_001481716.1
231	435	35.1	a) hypothetical protein	<i>C. jejuni</i> 81-176	296/300 (98%)	gb CP000538.1
			b) hypothetical protein cju10	<i>C. jejuni</i> 81-176	81/83 (97%)	gb ABF83701.1
244	399	35.8	a) conserved hypothetical protein, & DNA gyrase, A subunit	<i>C. jejuni</i> 81-176	194/195 (99%), 125/126 (99%)	gb CP000538.1
			b) conserved hypothetical protein	<i>C. coli</i> RM2228	55/55 (100%)	ZP_00370899.1
253	268	38.8	a) hypothetical protein,	<i>C. jejuni</i> 81116	190/190 (100%)	gb CP000814.1
			b) hypothetical protein C8J_0400	<i>C. jejuni</i> 81116	62/67 (92%)	YP_001481976.1
277	340	32.9	a) hypothetical protein	<i>C. jejuni</i> 81116	249/250 (99%)	gb CP000814.1
			b) hypothetical protein C8J_0065	<i>C. jejuni</i> 81116	74/74 (100%)	YP_001481642.1
278	340	32.9	a) hypothetical protein	<i>C. jejuni</i> 81116	249/250 (99%)	gb CP000814.1
			b) hypothetical protein C8J_0065	<i>C. jejuni</i> 81116	74/74 (100%)	YP_001481642.1
286	285	33.5	a) hypothetical protein	<i>C. jejuni</i> 81116	196/196 (100%)	Gb CP000814.1
			b) hypothetical protein CJE0273	<i>C. jejuni</i> RM1221	40/48 (83%)	YP_178296.1
287	343	40.6	a) no significant similarity to any nucleic acid			
			b) CDP-abequose synthase	<i>Blastopirellula marina</i> DSM 3645	41/94 (43%)	ZP_01090940.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.9. Unique clones, recovered from suppressive subtractive hybridizations of *C. jejuni* isolate 14118 (Caco-2 high invasion) determined to possess significant similarity to *Campylobacter* spp. other than *C. jejuni* 11168 (PMSRU).

Isolate	Percentage of Clones (n=68)
<i>C. jejuni</i> 81116	52.3%
<i>C. jejuni</i> LC	2.0%
<i>C. jejuni</i> RM1553	3.0%
<i>C. jejuni</i> 43431	2.0%
<i>C. jejuni</i> RM3423	2.0%
<i>C. jejuni</i> RM1508	2.0%
<i>C. jejuni</i> RM3419	2.0%
<i>C. jejuni</i> 11828	2.0%
<i>C. jejuni</i> D5477	2.0%
<i>C. jejuni</i> 81-176	11.0%
<i>C. jejuni</i> RM1167	2.0%
<i>C. jejuni</i> RM1861	2.0%
<i>C. jejuni</i> RM1221	2.0%
<i>C. jejuni</i>	5.0%

Table 4.10. Isolate 5116 (Caco-2 mid invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
Cell Envelop & surface structures						
4	524	29.7	a) lipooligosaccharide biosynthesis gene locus b) putative glycosyltransferase	<i>C. jejuni</i> RM3423 <i>C. jejuni</i>	446/448 (99%) 108/114 (94%)	gb EF143352.1 gb ABN41491.
37	261	42.9	a) hypothetical protein b) chimeric flagellin A/B	<i>C. jejuni</i> 81116 <i>C. jejuni</i>	183/183 (100%) 63/71 (88%)	gb CP000814.1 gb AAF25214.1 AF202168_1
59	573	25.0	a) no significant similarities to any nucleic acid b) capsular polysaccharide synthesis	<i>C. coli</i> RM2228	66/158 (41%)	ZP_00368108.1
65	395	31.9	a) no significant similarities to any nucleic acid b) capsular polysaccharide synthesis	<i>C. coli</i> RM2228	85/106 (80%)	ZP_00368108.1
84	275	38.5	a) cell division protein FtsK, putative b) cell division protein FtsK, putative,	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	195/198 (98%) 50/50 (100%)	Gb CP000538.1 ZP_02271252.1
88	511	34.4	a) no significant similarities to any nucleic acid b) putative periplasmic protein,	<i>C. jejuni</i> HB93-13	145/146 (99%)	ZP_01071241.1
90	338	38.5	a) capsular polysaccharide biosynthesis protein b) capsular polysaccharide biosynthesis protein	<i>C. doylei</i> 269.97 <i>C. jejuni</i> HB93-13	240/261 (91%) 84/97 (86%)	gb CP000768.1 ZP_01071340.1
Restriction – modification, recombination and repair						
55	498	30.5	a) type I restriction modification b) type I restriction modification DNA specificity domain protein	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	421/422 (99%) 140/155 (90%)	gb CP000538.1 YP_001000445.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit



Table 4.10.cont: Isolate 5116 (Caco-2 mid invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
16	253	39.1	a) 1062 bp at 5' side: MATE efflux family protein 176 bp at 3' side: phosphate ABC transporter, ATP-binding protein	<i>C. doylei</i> 269.97	165/177 (93%)	gb CP000768.1
34	257	43.1	b) conserved hypothetical protein, a) baseplate assembly protein V, putative	<i>C. jejuni</i> 260.94 <i>C. jejuni</i> RM1221	57/59 (96%) 179/180 (99%)	ZP_01069942.1 gb CP000025.1
77	253	39.5	b) baseplate assembly protein V, putative a) 1062 bp at 5' side: MATE efflux family protein 176 bp at 3' side: phosphate ABC transporter, ATP-binding protein	<i>C. jejuni</i> 260.94 <i>C. doylei</i> 269.97	60/64 (93%) 165/177 (93%)	ZP_01070038.1 gb CP000768.1
			b) conserved hypothetical protein	<i>C. jejuni</i> 260.94	57/59 (96%)	ZP_01069942.1
Small Molecule Metabolism						
29	660	41.5	a) rRNA-23S ribosomal RNA	<i>C. jejuni</i> RM1221	582/591 (98%)	gb CP000025.1
			b) conserved hypothetical protein,	<i>C. jejuni</i> CF93-6	35/35 (100%)	ZP_01067405.1
64	415	32.5	a), phosphoribosylglycinamide formyltransferase	<i>C. jejuni</i> 81-176	338/339 (99%)	gb CP000538.1
			b) phosphoribosylglycinamide formyltransferase	<i>C. jejuni</i> 81-176	112/116 (96%)	YP_999906.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.10.cont. Isolate 5116 (Caco-2 mid invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
Other (bacteriophage sequence)						
12	463	42.3	a) <i>Campylobacter</i> phage CGC-2007 isolate Cj00-2544 <i>cje0217</i> gene, partial cds; <i>cje0218</i> , <i>cje0219</i> , <i>cje0220</i> , <i>cje0221</i> , <i>cje0222</i> , <i>cje0223</i> , <i>cje0224</i> , <i>cje0225</i> , <i>cje0226</i> , <i>cje0227</i> , <i>cje0228</i> , <i>cje0229</i> , <i>cje0230</i> , and <i>cje0231</i> genes, complete cds; and <i>cje0232</i> gene, partial cds b) <i>cje0227</i> , <i>Campylobacter</i> phage CGC-2007]	<i>Campylobacter</i>	373/384 (97%)	gb EF694687.1
Hypothetical and unknown proteins						
3	570	27.5	a) no significant similarities to any nucleic acid b) conserved hypothetical protein		81/164 (49%)	ZP_01067542.1
10	373	30.7	a) hypothetical protein, b) hypothetical protein C8J_1252	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	290/290 (100%) 96/97 (98%)	gb CP000814.1 YP_001482828.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.11. Unique clones, recovered from suppressive subtractive hybridizations, of *C. jejuni* isolate 5116 (Caco-2 mid invasion) determined to possess significant similarity to *Campylobacter* spp. other than *C. jejuni* 11168 (PMSRU).

Isolate	Percentage of Clones (%) (n=16)
<i>C. jejuni</i> RM3423	6.0%
<i>C. jejuni</i> 81116	13.0%
<i>C. jejuni</i> RM2228	13.0%
<i>C. jejuni</i> 81-176	19.0%
<i>C. jejuni</i> HP93-13	6.0%
<i>C. jejuni</i> RM1221	13.0%

Table 4.12. Isolate 8557 (Caco-2 mid invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
Cell Envelop & Surface structures						
31	268	44	a) tail fiber protein H, putative b) tail fiber protein H, putative	<i>C. jejuni</i> RM 1221 <i>C. jejuni</i> 260.94	188/191 (98%) 63/64 (98%)	CP000025.1 YP_178254.1
49	336	34.2	a) No Significant Similarities to any nucleic acid b) putative periplasmic protein	<i>C. jejuni</i> HB93-13	85/86 (98%)	ZP_01071241.1
52	415	38.0	a) putative peptide chain release factor 2 ( <i>prfB</i> ) gene, partial cds; hypothetical protein, <i>CysD</i> ( <i>cysD</i> ), <i>CysN</i> ( <i>cysN</i> ), putative sodium/sulfate symporter, putative adenylylsulfate kinase ( <i>cysC</i> ), hypothetical protein, and putative glycosyltransferase genes; and <i>cj1457c</i> gene, partial cds b) <i>CysN</i>	<i>C. jejuni</i> ATCC 43432 <i>C. jejuni</i>	329/339 (97%) 102/105 (97%)	AY791516.1 AAX33831.1
63	556	30.5	a) LOS biosynthesis cluster b) unknown,	<i>C. jejuni</i> 11828 <i>C. jejuni</i>	476/480 (99%) 159/163 (97%)	AF343914.1 AAK12964.1
69	244	28.6	a) capsular polysaccharide biosynthesis protein, b) capsular polysaccharide biosynthesis protein	<i>C. doylei</i> 269.97 <i>C. jejuni</i> HB93-13	292 bits (158%) 58/63 (92%)	CP000768.1 ZP_01071278.1
86	566	31.9	a) baseplate assembly protein W b) putative baseplate assembly protein W	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> RM1221	483/490 (98%) 91/93 (97%)	CP000025.1 CAB94938.1
134	396	29.0	a) motility accessory factor b) motility accessory factor	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	319/319 (100%) 90/91 (98%)	gb CP000538.1 YP_001000998.1
135	266	36.5	a) putative peptide chain release factor 2 ( <i>prfB</i> ) gene, partial cds; hypothetical protein, <i>CysD</i> , <i>CysN</i> , putative sodium/sulfate symporter, putative adenylylsulfate kinase, <i>cysC</i> , hypothetical protein, and putative glycosyltransferase genes; and <i>Cj1457c</i> gene, partial cds b) putative sodium/sulfate symporter,	<i>C. jejuni</i> ATCC 43432 <i>C. jejuni</i>	187/189 (98%) 63/66 (95%)	gb AY791516.1 gb AAX33832.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.12 cont. Isolate 8557 (Caco-2 mid invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
194	250	42.4	a) class O lipooligosaccharide biosynthesis gene locus	<i>C. jejuni</i> RM3423	174/174 (100%)	gb EF143352.1
200	111	41.4	b) putative aminotransferase, a) flagellar hook protein	<i>C. jejuni</i> <i>C. doylei</i> 269.97	60/70 (85%) 52/53 (98%)	gb ABN41492.1 gb CP000768.1
204	381	35.7	b) flagellar hook subunit protein	<i>C. jejuni</i> CG8486	18/18 (100%)	ZP_01810497.1
213	238	39.5	a) capsular polysaccharide biosynthesis protein	<i>C. doylei</i> 269.97	282/303 (93%)	gb CP000768.1
			b) capsular polysaccharide biosynthesis protein	<i>C. jejuni</i> HB93-13	95/102 (93%)	ZP_01071340.1
			a) class O lipooligosaccharide biosynthesis gene locus	<i>C. jejuni</i> RM3423	160/160 (100%)	gb CP000814.1
			b) putative glucose-1-phosphate thymidyltransferase,	<i>C. jejuni</i>	53/53 (100%)	gb ABN41485.1
Restriction-modification, recombination & repair						
34	241	34	a) no significant similarities to any nucleic acid b) adenine-specific methyltransferase	<i>Fusobacterium nucleatum</i>	34/52 (65%)	NP_602723.1
74	269	39.7	a) prophage MuSo1, F protein, putative	<i>C. jejuni</i> RM1221	192/192 (100%)	CP000025.1
191	493	34.9	b) prophage MuSo1, F protein, putative a) type I restriction-modification system, M subunit	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> 81116	30/30 (100%) 408/411 (99%)	YP_178274.1 gb CP000814.1
			b) type I restriction-modification system, M subunit	<i>C. jejuni</i> 81116	137/137 (100%)	YP_001483027.1
220	291	34.0	a) HsdR pseudogene, hsdR-1 allele, complete sequence; RloG gene, complete cds; HsdS pseudogene, hsdS-5 allele, and HsdM gene, hsdM-1 allele	<i>C. jejuni</i> RM1167	130/135 (96%)	gb AF486547.1
			b) no significant similarities to any protein		236/244 (96%)	gb CP000814.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.12 cont. Isolate 8557 (Caco-2 mid invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
236	791	34.4	a) <i>HsdR</i> gene, <i>hsdR</i> -1 allele, complete cds; <i>RloD</i> gene, complete cds; <i>HsdS</i> gene, <i>hsdS</i> -3 allele, complete cds; <i>MloA</i> gene, complete cds; and <i>HsdS</i> gene, <i>hsdM</i> -1 allele, complete cds	<i>C. jejuni</i> RM2240	708/715 (99%)	gb AF486556.1
285	411	31.3	b) <i>HsdM</i>	<i>C. jejuni</i>	236/238 (99%)	gb AAM00874.1
			a) type I restriction-modification system, M subunit	<i>C. jejuni</i> 81-176	306/338 (90%)	gb CP000538.1
			b) putative restriction enzyme subunit S	<i>C. jejuni</i> 260.94	110/111 (99%)	ZP_01070278.1
Transport						
46	343	39	a) Na/Pi-cotransporter, putative	<i>C. jejuni</i> 81-176	252/255 (98%)	CP000538.1
			b) Na/Pi-cotransporter, putative	<i>C. jejuni</i> HB93-13	51/53 (96%)	ZP_01071640.1
81	725	30.0	a) Na/Pi-cotransporter, putative	<i>C. jejuni</i> 81-176	120/121 (99%)	CP000538.1
			b) predicted ATP-dependent endonuclease of the OLD family	<i>C. jejuni</i> 260.94	153/155 (98%)	ZP_01070305.1
126	402	30.8	a) permease, putative	<i>C. jejuni</i> RM1221	324/327 (99%)	gb CP000025.1
			b) hypothetical protein Cj8486_1595c	<i>C. jejuni</i> CG8486	24/25 (96%)	ZP_01809396.1
214	726	37.2	a) Na/Pi-cotransporter, putative	<i>C. jejuni</i> 81-176	632/634 (99%)	gb CP000538.1
			b) putative penicillin-binding protein	<i>C. jejuni</i> 81116	131/138 (94%)	YP_001482062.1
219	382	35.6	a) putative peptide chain release factor 2 ( <i>prfB</i> ) gene, partial cds; hypothetical protein, <i>CysD</i> , <i>CysN</i> , putative sodium/sulfate symporter, putative adenylylsulfate kinase ( <i>cysC</i> ), hypothetical protein, and putative glycosyltransferase genes, complete cds; and <i>Cj1457c</i> gene, partial cds	<i>C. jejuni</i> ATCC 43432	284/284 (100%)	gb AY791516.1
			b) putative sodium/sulfate symporter	<i>C. jejuni</i>	94/94 (100%)	gb AAX33832.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.12 cont. Isolate 8557 (Caco-2 mid invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
234	356	31.1	a) transporter, putative	<i>C. jejuni</i> RM1221	271/279 (97%)	gb CP000025.1
			b) conserved hypothetical protein	<i>C. jejuni</i> 81-176	25/27 (92%)	YP_001001196.1
Small Molecule Metabolism						
41	239	34.3	a) oxidoreductase, putative	<i>C. jejuni</i> 81-176	144/150 (96%)	CP000025.1
			b) oxidoreductase, putative	<i>C. coli</i> RM2228	49/51 (96%)	YP_179670.1
98	307	27.4	a) lysyl-tRNA synthetase	<i>C. jejuni</i> 81116	207/207 (100%)	gb CP000814.1
			b) lysyl-tRNA synthetase	<i>C. jejuni</i> 81116	69/75 (92%)	gb ABV51975.1
102	445	48.4	a) dipeptidyl-peptidase	<i>C. jejuni</i> 81-176	359/362 (99%)	CP000538.1
			b) X-Pro dipeptidyl-peptidase family protein	<i>C. jejuni</i> HB93-13	120/121 (99%)	ZP_01071387.1
166	347	40.6	a) dipeptidyl-peptidase,	<i>C. jejuni</i> 81-176	257/258 (99%)	gb CP000538.1
			b) X-Pro dipeptidyl-peptidase family protein	<i>C. jejuni</i> HB93-13	79/81 (97%)	ZP_01071387.1
168	233	41.6	a) arylsulfate sulfotransferase	<i>C. jejuni</i> 81116	143/144 (99%)	b CP000814.1
			b) arylsulfate sulfotransferase	<i>C. jejuni</i> 81116	47/50 (94%)	YP_001482389.1
181	520	32.8	a) TPR domain protein	<i>C. jejuni</i> 81-176	297/301 (98%)	gb CP000538.1
			b) putative transmembrane protein	<i>C. jejuni</i> 81116	76/89 (85%)	YP_001481941.1
183	457	32.0	a) hypothetical protein	<i>C. jejuni</i> 81116	353/363 (97%)	b CP000814.1
			b) lectin C-type domain protein	<i>C. doylei</i> 269.97	32/34 (94%)	YP_001398053.1
185	401	38.9	a) histidyl-tRNA synthetase	<i>C. jejuni</i> 81-176	300/302 (99%)	b CP000538.1
			b) histidyl-tRNA synthetase	<i>C. jejuni</i> 81116	96/105 (91%)	YP_001482292.1
196	467	40.0	a) molybdopterin-guanine dinucleotide biosynthesis protein MobB	<i>C. jejuni</i> 81-176	199/199 (100%)	gb CP000538.1
			b) molybdopterin-guanine dinucleotide biosynthesis protein MobB	<i>C. jejuni</i> 260.94	62/63 (98%)	ZP_01070343.1
208	900	30.3	a) putative aminotransferase (DegT family)	<i>C. jejuni</i> 81116	237/246 (96%)	gb CP000814.1
			b) probable aminotransferase (degT family)	<i>C. coli</i> RM2228	44/44 (100%)	ZP_00367343.1
222	425	34.1	a) CrcB heat shock protein Htp	<i>C. jejuni</i> 81116	236/244 (96%)	gb CP000814.1
			b) CRCB protein like protein,	<i>C. jejuni</i> CG8486	42/42 (100%)	ZP_01809576.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.12 cont. Isolate 8557 (Caco-2 mid invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
229	387	36.2	a) putative pyrazinamidase/nicotinamidase b) hypothetical protein CJJ81176_0155	<i>C. doylei</i> 269.97 <i>C. jejuni</i> 81-176	310/310 (100%) 96/96 (100%)	b CP000768.1 YP_999844.1
231	844	33.4	a) no significant similarities to any nucleic acid b) ATP/GTP-binding protein	<i>C. jejuni</i> 260.94	208/216 (96%)	ZP_01070279.1
260	288	34.0	a) no significant similarities to any nucleic acid b) CAAX amino terminal protease family protein	<i>C. lari</i> Rm2100	7/37 (100%) 35/40 (87%)	gb AY725194.1 ZP_00369366.1
261	320	43.1	a) MmgE/PrpD family protein b) MmgE/PrpD family protein	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	243/243 (100%) 80/81 (98%)	gb CP000538.1 YP_001482885.1
266	299	37.1	a) host-nuclease inhibitor protein, putative b) host-nuclease inhibitor protein, putative	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> 260.94	182/207 (87%) 69/70 (98%)	gb CP000025.1 ZP_01069888.1
Other (bacteriophage sequence)						
2	263	37.6	a) bacteriophage DNA transposition protein A, putative b) bacteriophage DNA transposition protein A, putative	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> CF93-6	183/184 (99%) 61/62 (98%)	CP000025.1 ZP_01068156.1
56	228	33.9	a) phage tail protein, putative b) phage tail protein	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> RM1221	138/139 (99%) 41/42 (97%)	CP000025.1 YP_178275.1
123	507	36.8	a) <i>Campylobacter</i> phage CGC-2007 isolate Cj00-3477 <i>cje0217</i> -like gene, complete sequence; <i>cje0218</i> , <i>cje0219</i> , <i>cje0220</i> , <i>cje0221</i> , <i>cje0222</i> , <i>cje0223</i> , <i>cje0224</i> , <i>cje0225</i> , <i>cje0226</i> , <i>cje0227</i> , <i>cje0228</i> , <i>cje0229</i> , <i>cje0230</i> , and <i>cje0231</i> genes, complete cds; and <i>cje0232</i> gene, partial cds b) <i>cje0231</i> , <i>Campylobacter</i> phage CGC-2007	<i>Campylobacter</i> <i>Campylobacter</i>	429/429 (100%) 142/143 (99%)	gb EF694689.1 gb ABU53798.1
139	488	37.5	a) downstream insertion site of CMLP1-like temperate bacteriophage b) bacteriophage DNA transposition protein A	<i>C. jejuni</i> <i>C. jejuni</i> 260.94	408/410 (98%) 133/137 (97%)	B EF092316.1 ZP_01069915.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit



Table 4.12 cont. Isolate 8557 (Caco-2 mid invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
162	320	41.8	a) <i>Campylobacter</i> phage CGC-2007 isolate Cj00-3477 <i>cje0217</i> -like gene, complete sequence; <i>cje0218</i> , <i>cje0219</i> , <i>cje0220</i> , <i>cje0221</i> , <i>cje0222</i> , <i>cje0223</i> , <i>cje0224</i> , <i>cje0225</i> , <i>cje0226</i> , <i>cje0227</i> , <i>cje0228</i> , <i>cje0229</i> , <i>cje0230</i> , and <i>cje0231</i> genes, complete cds; and <i>cje0232</i> gene, partial cds	<i>Campylobacter</i>	237/238 (99%)	gb EF694689.1
177	306	42.4	b) <i>cje0222</i> , <i>Campylobacter</i> phage CGC-2007	<i>Campylobacter</i>	42/43 (97%)	gb ABU53725.1
			a) <i>Campylobacter</i> phage CGC-2007 isolate Cj00-3477 <i>cje0217</i> -like gene, complete sequence; <i>cje0218</i> , <i>cje0219</i> , <i>cje0220</i> , <i>cje0221</i> , <i>cje0222</i> , <i>cje0223</i> , <i>cje0224</i> , <i>cje0225</i> , <i>cje0226</i> , <i>cje0227</i> , <i>cje0228</i> , <i>cje0229</i> , <i>cje0230</i> , and <i>cje0231</i> genes, complete cds; and <i>cje0232</i> gene, partial cds	<i>Campylobacter</i>	215/215 (100%)	gb EF694689.1
189	383	37.0	b) <i>cje0231</i> , <i>Campylobacter</i> phage CGC-2007	<i>Campylobacter</i>	73/76 (96%)	gb ABU53798.1
			a) <i>Campylobacter</i> phage CGC-2007 isolate Cj00-3477 <i>cje0217</i> -like gene, complete sequence; <i>cje0218</i> , <i>cje0219</i> , <i>cje0220</i> , <i>cje0221</i> , <i>cje0222</i> , <i>cje0223</i> , <i>cje0224</i> , <i>cje0225</i> , <i>cje0226</i> , <i>cje0227</i> , <i>cje0228</i> , <i>cje0229</i> , <i>cje0230</i> , and <i>cje0231</i> genes, complete cds; and <i>cje0232</i> gene, partial cds	<i>Campylobacter</i>	305/307 (99%)	gb EF694689.1
192	250	42.4	b) <i>cje0231</i> , <i>Campylobacter</i> phage CGC-2007	<i>Campylobacter</i>	70/70 (100%)	gb ABU53865.1
			a) phage uncharacterized protein	<i>C. jejuni</i> RM1221	179/181 (98%)	gb CP000025.1
195	440	32.5	b) phage uncharacterized protein	<i>C. jejuni</i> RM1221	25/25 (100%)	YP_178272.1
			a) <i>Campylobacter</i> phage CGC-2007 isolate CjNC13256 <i>cje0217</i> gene, partial cds; <i>cje0218</i> , <i>cje0219</i> , <i>cje0220</i> , <i>cje0221</i> , <i>cje0222</i> , <i>cje0223</i> , <i>cje0224</i> , <i>cje0225</i> , <i>cje0226</i> , <i>cje0227</i> , <i>cje0228</i> , <i>cje0229</i> , <i>cje0230</i> , and <i>cje0231</i> genes, complete cds; and <i>cje0232</i> gene, partial cds	<i>Campylobacter</i>	355/356 (99%)	gb EF694693.1
202	311	40.3	b) <i>cje0221</i> , <i>Campylobacter</i> phage CGC-2007	<i>Campylobacter</i>	99/100 (99%)	gb ABU53788.1
			a) Mu-like prophage I protein	<i>C. jejuni</i> RM1221	235/235 (100%)	gb CP000025.1
			b) Mu-like prophage I protein, putative	<i>C. jejuni</i> 260.94	77/77 (100%)	ZP_01069769.1

Table 4.12 cont. Isolate 8557 (Caco-2 mid invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
215	461	41.6	a) <i>Campylobacter</i> phage CGC-2007 isolate Cj00-0949 <i>cje0215</i> gene, partial cds; <i>cje0216</i> , <i>cje0217</i> , <i>cje0218</i> , <i>cje0219</i> , <i>cje0220</i> , <i>cje0221</i> , <i>cje0222</i> , <i>cje0223</i> , <i>cje0224</i> , <i>cje0225</i> , <i>cje0226</i> , <i>cje0227</i> , <i>cje0228</i> , <i>cje0229</i> , <i>cje0230</i> , and <i>cje0231</i> genes, complete cds; and <i>cje0232</i> gene, partial cds	<i>Campylobacter</i>	375/384 (97%)	gb EF694684.1
250	132	47.7	b) <i>cje0227</i> , <i>Campylobacter</i> phage CGC-2007	<i>Campylobacter</i>	126/128 (98%)	gb ABU53861.1
			a) <i>Campylobacter</i> phage CGC-2007 isolate Cj00-2818 <i>cje0215</i> gene	<i>Campylobacter</i>	75/76 (98%)	gb EF694688.1
			b) <i>cje0227</i> , <i>Campylobacter</i> phage CGC-2007	<i>Campylobacter</i>	25/25 (100%)	gb ABU53861.1
Hypothetical & unknown proteins						
1	68	34.5	a) no significant similarities to any nucleic acid b) no significant similarities to any protein			
50	402	37.0	a) hypothetical protein b) hypothetical protein CJJ26094_1718	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 260.94	308/325 (94%) 100/104 (96%)	CP000814.1 ZP_01070483.1
82	78	48.7	a) no significant similarities to any nucleic acid b) no significant similarities to any protein			
89	644	36.2	a) hypothetical protein b) conserved hypothetical protein		480/493 (97%) 98/100 (98%)	CP000025.1 ZP_01069745.1
114	746	38.0	a) conserved hypothetical protein, b) hypothetical protein CJJ81176_1327	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	504/510 (98%) 169/171 (98%)	ZP_01071387.1 gb CP000538.1
122	214	36.0	a) hypothetical protein b) hypothetical protein CJJ26094_1718	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 260.94	166/175 (94%) 55/58 (94%)	b CP000814.1 ZP_01070483.1
128	505	34.5	a) hypothetical protein b) conserved hypothetical protein	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> 260.94	425/427 (99%) 141/142 (99%)	Gb CP000025.1 ZP_01069964.1
144	395	38.7	a) conserved hypothetical protein, b) hypothetical protein C8J_0876	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81116	306/306 (100%) 101/102 (99%)	gb CP000538.1 YP_001482452.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.12 cont. Isolate 8557 (Caco-2 mid invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
147	188	47	a) conserved hypothetical protein, b) hypothetical protein C8J_0878	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81116	111/111 (100%) 36/37 (97%)	gb CP000538.1 YP_001482454.1
161	379	39	a) conserved hypothetical protein, b) hypothetical protein CJJ81176_0761	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	221/222 (99%) 74/79 (93%)	gb CP000538.1 YP_001000434.1
172	105	49.5	a) no significant similarities to any nucleic acid b) no significant similarities to any protein			
176	302	37.4	a) hypothetical protein b) hypothetical protein C8J_1093	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	216/220 (98%) 69/72 (95%)	gb CP000814.1 YP_001482669.1
201	362	34.5	a) hypothetical protein b) hypothetical protein CJJ26094_0512	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> 260.94	271/273 (99%) 91/91 (100%)	Gb CP000025.1 ZP_01070032.1
203	265	40.3	a) conserved hypothetical protein b) hypothetical protein C8J_0877	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81116	187/188 (99%) 59/62 (95%)	gb CP000538.1 YP_001482453.1
218	215	41.7	a) domain of unknown function (DUF955) superfamily b) domain of unknown function (DUF955) superfamily	<i>C. jejuni</i> 81116 <i>C. jejuni</i> 81116	137/139 (98%) 47/48 (97%)	Gb CP000814.1 YP_001482223.1
226	567	39.8	a) conserved hypothetical protein b) hypothetical protein CJE0246	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> RM1221	483/485 (99%) 159/163 (97%)	gb CP000025.1 YP_178269.1
242	334	38.0	a) conserved domain protein b) conserved domain protein	<i>C. jejuni</i> RM1221	249/259 (96%) 59/59 (100%)	gb CP000025.1 ZP_01070050.1
255	211	40.3	a) hypothetical protein b) hypothetical protein CJJ26094_0512		134/134 (100%) 44/44 (100%)	gb CP000025.1 ZP_01070032.1
264	290	37.1	a) conserved hypothetical protein b) conserved hypothetical protein		213/213 (100%) 74/82 (90%)	gb CP000538.1 ZP_01071231.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.13. Unique clones, recovered from suppressive subtractive hybridizations, of *C. jejuni* isolate 8557 (Caco-2 mid invasion) determined to possess significant similarity to *Campylobacter* spp. other than *C. jejuni* 11168 (PMSRU).

Isolate	Percentage of Clones (%) (n=71)
<i>C. jejuni</i> RM1221	20.0%
<i>C. jejuni</i> 81-176	24.0%
<i>C. jejuni</i> HB93-13	1.4%
<i>C. jejuni</i> 43432	4.0%
<i>C. jejuni</i> 11828	1.4%
<i>C. jejuni</i> RM3423	3.0%
<i>C. jejuni</i> 81116	14.0%
<i>C. jejuni</i> RM1167	1.4%
<i>C. jejuni</i> RM224	1.4%

Table 4.14. Isolate 13262 (Caco-2 low invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
Cell Envelop & Surface structures						
1	352	32.0	a) no significant similarities to any nucleic acid b) putative beta-N-acetylgalactosaminyltransferase	<i>C. jejuni</i>	83/105 (79%)	ABN41494.1
3	842	33.8	a) co-chaperone protein DnaJ b) chaperone protein dnaJ	<i>C. jejuni</i> 81-176 <i>C. doylei</i> 269.97	211/228 (92%) 76/90 (84%)	CP000538.1 ZP_01070931.1
97	244	42.6	a) cell division protein FtsK, putative b) dna translocase spoiiiie	<i>C. jejuni</i> 81-176 <i>C. doylei</i> 269.97	151/162 (93%) 53/63 (84%)	CP000538.1 ZP_01070300.1
100	497	32.1	a) no significant similarities to any nucleic acid b) possible sugar transferase	<i>C. jejuni</i> CG8486	113/125 (90%)	ZP_01810450.1
173	375	34.6	a) peptidase family protein b) flagellar motor switch protein	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> CG8486	289/305 (94%) 63/72 (87%)	CP000538.1 ZP_01810543.1
174	251	37.8	a) flagellar biosynthetic protein FlhB b) ATP synthase F0 sector B subunit	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> CG8486	106/110 (96%) 28/30 (93%)	CP000538.1 ZP_01810581.1
181	345	32.7	a) DNA polymerase III, beta subunit b) DNA polymerase III, beta subunit	<i>C. jejuni</i> Rm1221 <i>C. jejuni</i> Rm1221	290/299 (98%) 97/101 (96%)	gb CP000025.1 ZP_01068757.1
237	553	35.2	a) flagellar hook assembly protein b) ABC transporter, ATP-binding protein	<i>C. doylei</i> 269.97 <i>C. doylei</i> 269.97	122/135 (90%) 26/27 (96%)	CP000025.1 ABS43505.1
Bacterial Toxin						
273	232	41.8	a) CDT operon ( <i>cdtA</i> , <i>cdtB</i> , <i>cdtC</i> genes) b) cytolethal distending toxin C	<i>C. lari</i> <i>C. lari</i>	95/109 (87%) 31/39 (79%)	AB292356.1 dbj BAF48048.1
Restriction- modification, recombination & repair						
9	429	35.2	a) DNA-binding protein Roi b) DNA-binding protein Roi	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> RM1221	192/201 (95%) 67/69 (97%)	CP000025.1 YP_179421.1
22	351	35.8	a) terminase B protein, putative b) terminase B protein, putative	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> RM1221	182/185 (98%) 43/45 (95%)	CP000025.1  CP000025.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.14 cont. 13262 (Caco-2 low invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
45	321	36.1	a) no significant Similarities to any nucleic acid b) phenylalanyl-tRNA synthetase, beta subunit	<i>C. jejuni</i> 260.94	65/83 (78%)	ZP_01070273.1
98	735	35.5	a) type II restriction-modification enzyme b) type II restriction-modification enzyme	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> RM1221	409/415 (98%) 140/150 (93%)	CP000025.1 YP_178058.1
106	230	39.5	a) single-stranded-DNA-specific exonuclease <i>RecJ</i> b) putative single-stranded-DNA-specific exonuclease	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> CG8486	151/157 (96%) 51/54 (94%)	CP000538.1 ZP_01810523.1
143	664	41.3	a) rRNA-23S ribosomal RNA b) conserved hypothetical protein	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> CF93-6	582/593 (98%) 48/51 (94%)	CP000025.1 ZP_01067405.1
199	263	33.8	a) possible polysaccharide modification protein b) hypothetical protein Cj8486_1461c		173/187 (92%) 59/63 (93%)	AY332625.1 ZP_01810447.1
275	390	38.9	a) DNA-binding protein Roi b) conserved domain protein	<i>C. doylei</i> 269.97 <i>C. jejuni</i> RM1221	263/287 (91%) 52/58 (89%)	CP000768.1 AAW35141.1
Transport 13	377	38.5	a) trkA domain protein b) hypothetical protein Cj8486_1071	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> CG8486	250/258 (96%) 86/86 (100%)	CP000538.1 ZP_01809767.1
111	265	34.3	a) di-/tripeptide transporter b) di- and tri-peptide transporter	<i>C. jejuni</i> 81-176 <i>C. doylei</i> 269.97	184/189 (97%) 43/45 (95%)	gb CP000538.1 AAV30680.1
129	492	34.3	a) CTP synthase b) CTP synthase	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	99/99 (100%) 99/99 (100%)	CP000538.1
190	145	37.2	a) putative peptide ABC-transport system periplasmic b) anaerobic dimethyl sulfoxide reductase chain A	<i>C. jejuni</i> 81-176 <i>C. jejuni</i> 81-176	86/87 (98%) 28/29 (97%)	DQ493924.1 AAV53798.1
191	412	33.4	a) GlnD family protein b) GlnD family protein	<i>C. jejuni</i> RM1221 <i>C. jejuni</i> RM1221	258/261 (98%) 85/88 (96%)	CP000025.1 YP_179542.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.14 cont. 13262 (Caco-2 low invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
213	511	30.2	a) no significant similarities to any nucleic acid			
			b) ABC transporter	<i>Beggiatoa</i> sp. PS	45/146 (30%)	EDN71435.1
279	269	37.2	a) macrolide-specific efflux protein macA	<i>C. jejuni</i> 81-176	209/213 (98%)	CP000538.1
			b) macrolide-specific efflux protein macA	<i>C. jejuni</i> 81-176	71/71 (100%)	YP_001398157.1
Other (bacteriophage sequence)						
125	406	30.8	a) site-specific recombinase, phage integrase family	<i>C. jejuni</i> RM1221	328/330 (99%)	CP000025.1
			b) site-specific recombinase, phage integrase family	<i>C. jejuni</i> RM1221	98/100 (98%)	YP_178560.1
Small molecular metabolism						
2	438	32.4	a) HAD-superfamily phosphatase, subfamily IIIC	<i>C. jejuni</i> 81-176	90/90 (100%)	CP000538.1
			b) hypothetical protein Cjejd_02000147	<i>C. doylei</i> 269.97	71/72 (92%)	ZP_01807491.1
16	796	31.4	a) D12 class N6 adenine-specific DNA methyltransferase	<i>C. jejuni</i> RM1221	274/279 (98%)	CP000025.1
			b) D12 class N6 adenine-specific DNA methyltransferase	<i>C. jejuni</i> . CG8486	118/153 (77%)	ZP_01810087.1
30	251	43.8	a) carbamoyl-phosphate synthase, large subunit	<i>C. jejuni</i> 81-176	99/105 (94%)	CP000538.1
			b) carbamoyl-phosphate synthase large chain	<i>C. jejuni</i> CG8486	33/34 (97%)	ZP_01810658.1
55	145	47.5	a) anaerobic dimethyl sulfoxide reductase chain A	<i>C. jejuni</i> 81-176	83/84 (98%)	CP000538.1
			b) anaerobic dimethyl sulfoxide reductase chain A	<i>C. jejuni</i>	27/27 (100%)	AAY53798.1
59	244	45.9	a) polyphosphate kinase	<i>C. jejuni</i> RM1221	144/148 (98%)	CP000025.1
			b) conserved hypothetical protein	<i>C. jejuni</i> 260.94	44/44 (100%)	ZP_01069225.1
115	194	40.7	a) 3-dehydroquinate synthase	<i>C. jejuni</i> 81-176	111/117 (94%)	CP000538.1
			b) 3-dehydroquinate synthase	<i>C. doylei</i> 269.97	36/40 (90%)	ZP_01069050.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.14 cont. 13262 (Caco-2 low invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
140	109	37.6	a) anaerobic dimethyl sulfoxide reductase chain A	<i>C. jejuni</i> 81-176	54/54 (100%)	CP000538.1
			b) anaerobic dimethyl sulfoxide reductase chain A	<i>C. jejuni</i>	18/19 (94%)	AAY53798.1
179	354	35.6	a) no significant similarities to any nucleic acid			
			b) GDP-L-fucose synthetase co-enzyme binding	<i>C. jejuni</i> 84-25	70/102 (68%)	ZP_01099795.1
187	599	38.8	a) dihydroorotase, homodimeric type D12 class N6 adenine-specific DNA methyltransferase	<i>C. jejuni</i> RM1221	376/409 (91%)	gb CP000025.1
			b) dihydroorotase	<i>C. jejuni</i> RM1221	101/107 (94%)	YP_178329.1
236	323	39.6	a) no significant similarities to any nucleic acid			
			b) biotin sulfoxide reductase ( <i>bisC</i> )	<i>C. lari</i> RM2100	74/82 (90%)	ZP_00368912.1
245	238	36.1	a) hydrogenase, (NiFe)/(NiFeSe) small subunit family	<i>C. jejuni</i> RM1221	157/192 (96%)	CP000025.1
			b) hydrogenase, (NiFe)/(NiFeSe) small subunit family	<i>C. jejuni</i> RM1221	51/54 (94%)	ZP_01070165.1
277	269	36.1	a) hydrogenase assembly chaperone HypC/HupF	<i>C. jejuni</i> 81-176	140/152 (92%)	CP000538.1
			b) hydrogenase assembly chaperone HypC/HupF	<i>C. coli</i> RM2228	51/61 (83%)	ZP_00366792.1
Hypothetical & unknown						
31	117	58.0	a) no significant similarities to any nucleic acid			
			b) no significant similarities to any protein			
60	75	50.1	a) no significant similarities to any nucleic acid			
			b) no significant similarities to any protein			

<sup>a</sup> a) BLASTN hit and b) BLASTX hit



Table 4.14 cont. 13262 (Caco-2 low invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
68	597	30.0	a) no significant similarities to any nucleic acid			
69	85	57.6	b) possible sugar transferase a) uncultured bacterium gene for 16S rRNA	<i>C. jejuni</i> CG8486	159/166 (95%) 44/44 (100%)	ZP_01810450.1 AB177205.1
124	378	30.9	b) no similarity to any protein a) hypothetical protein	<i>C. jejuni</i> RM1221	170/172 (98%)	CP000025.1
136	430	35.1	b) hypothetical protein CJE0592 a) polyribonucleotide	<i>C. jejuni</i> RM1221	38/40 (95%) 347/353 (98%)	YP_178608.1 CP000025.1
			nucleotidyltransferase, conserved hypothetical protein,	<i>C. jejuni</i> 260.94	96/98 (97%)	ZP_01069129.1
146	200	43.5	b) conserved hypothetical protein a) hypothetical protein	<i>C. jejuni</i> CF93-6	121/123 (98%)	ZP_01068327.1
150	297	34.3	b) hypothetical protein	<i>C. jejuni</i> CF93-6	40/41 (97%)	ZP_01068327.1
175	383	32.6	a) conserved hypothetical protein b) hypothetical protein CJJ81176_0772,	<i>C. jejuni</i> 81-176	211/221 (95%) 19/19 (100%)	CP000538.1 YP_001000440.1
192	83	48.2	a) no significant similarities to any nucleic acid b) hypothetical protein	<i>C. jejuni</i> RM1221	45/106 (45%)	ZP_01834321.1
201	498	32.5	a) no significant similarities to any nucleic acid b) no significant similarities to protein			
211	250	37.2	a) conserved domain protein b) conserved domain protein	<i>C. jejuni</i> RM1221	396/399 (99%) 132/136 (97%)	CP000025.1 AAW35931.1
218	166	51.2	a) conserved domain protein b) hypothetical protein CJE0556	<i>C. jejuni</i> RM1221	173/174 (99%) 59/64 (92%)	CP000025.1 YP_178572.1
222	321	38.6	a) hypothetical protein b) hypothetical protein CJE0590	<i>C. jejuni</i> RM1221	187/189 (98%) 62/63 (98%)	P000025.1 YP_178606.1
240	248	38.7	a) hypothetical protein b) hypothetical protein CJJ26094_1412	<i>C. jejuni</i> <i>C. jejuni</i> 260.94	227/245 (91%) 69/80 (86%)	CP000025.1 ZP_01069043.1
			a) no significant similarities to any nucleic acid b) no significant similarities to any protein			

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.14 cont. 13262 (Caco-2 low invasion) unique clones relative to *C. jejuni* 11168 (PMSRU).

Category & Clone #	Length (bp)	% G+C	BLASTN and BLASTX results <sup>a</sup>	Organism	% Similarity % Identity	Accession no.
253	289	35.6	a) hypothetical protein	<i>C. doylei</i> 269.97	205/213 (96%)	CP000768.1
			b) hypothetical protein JJD26997_1925	<i>C. doylei</i> 269.97	63/64 (98%)	YP_001398856.1
271	401	31.6	a) hypothetical protein	<i>C. jejuni</i> RM1221	219/226 (96%)	CP000025.1
			b) conserved hypothetical protein	<i>C. jejuni</i> RM1221	60/81 (74%)	ABS43420.1

<sup>a</sup> a) BLASTN hit and b) BLASTX hit

Table 4.15. Unique clones, recovered from suppressive subtractive hybridizations, of *C. jejuni* isolate 13262 (Caco-2 low invasion) determined to possess significant similarity to *Campylobacter* spp. other than *C. jejuni* 11168 (PMSRU).

Isolate	Percentage of Clones (n=59)
<i>C. jejuni</i> 81-176	31.0%
<i>C. jejuni</i>	2.0%
<i>C. jejuni</i> CH8486	4.0%
<i>C. jejuni</i> RM1221	27.0%
<i>C. jejuni</i> 260.94	2.0%
<i>C. jejuni</i> HS:23	2.0%
<i>C. jejuni</i> 84-25	2.0%
<i>C. jejuni</i> CF93-6	2.0%




Figure 4.1: Dendrogram of representative *Campylobacter* spp. *flaA* SVR alleles recovered during the Icelandic Epidemiology Investigation.




Figure 4.2. Distribution of *Campylobacter* spp. *flaA* SVR allele groups by percent adhesion to Caco-2 cells.

### Distribution of Host Type by % Adhesion




Figure 4.3 Distribution of *Campylobacter* spp. percent adhesion of Caco-2 cells relative to original host.




Figure 4.4. Distribution of *Campylobacter* spp. *flaA* SVR allele groups by percent invasion of Caco-2 cells.

### Distribution of % Invasion of Caco-2 Cells Relative to host Type




Figure 4.5. Distribution of *Campylobacter* spp. host type by percent invasion of Caco-2 cells.



### Number of Clones Belonging to Specific Functional Categories




Figure 4.6. Functional categories of clones recovered using suppressive subtractive hybridization on all 4 *C. jejuni* isolates.

Chapter 5  
Summary and Conclusions

The studies presented were based on three primary objectives: 1) To optimize conditions for use with the Bactometer® such that efficient and reproducible monitoring of *Campylobacter* spp. for determination of growth curves was achieved in a simple medium, 2) To investigate the adherence and invasiveness of 52 *Campylobacter* spp. isolates using Caco-2 cells, and 3) To investigate the genetic diversity of four *C. jejuni* isolates that demonstrated a wide range of invasiveness towards human colonic cells.

For the first objective, conditions for use with the Bactometer® were investigated to determine efficient and reproducible monitoring of *Campylobacter* spp. growth curves in a simple medium. Results suggested that isolates be grown on Mueller Hinton plates under a microaerobic atmosphere (37°C; 24 h), followed by transfer to Mueller Hinton biphasic cultures for 6 h (37°C; microaerobic atmosphere). Serial dilutions were inoculated in Bactometer® wells containing 1 mL Mueller Hinton broth plus 0.1M sodium pyruvate. Utilizing the Bactometer® is important since determining growth curves by hand is a time consuming and labor intensive process.

For the second objective, adhesion and invasion assays were performed on 52 *Campylobacter* spp. isolates using human Caco-2 cells. *Campylobacter* spp. exhibited a wide distribution of adhesion and invasion ability, which was determined unrelated to *flaA* SVR allele type. There also appeared to be no relationship between host of recovery source and level of adhesion or invasion. This objective was important in determining if the *flaA* SVR allele type could predict potential virulence.

The last objective investigated the genetic diversity of four *C. jejuni* isolate that demonstrated a wide range of invasiveness towards human colonic cells. Four isolates comprised of the most invasive isolate (14118), the least invasive isolate (13262), and two

isolates in between were selected for DNA: DNA microarray hybridizations and suppressive subtractive hybridizations. DNA:DNA microarray hybridizations identified genes absent relative to 11168 (PMSRU) and also determined 372 genes present in *C. jejuni* isolates 14118, 5116, 8557, and 13262 as well as *C. jejuni* 11168 (PMSRU). Suppressive subtractive hybridizations identified genes absent from *C. jejuni* 11168 (PMSRU). *C. jejuni* 14118 contained a gene from *C. doylei* 269.97 that encoded for a motility accessory factor and a gene involving transport. *C. jejuni* 13262 contained a CDT operon from *C. lari* as well as included a type II restriction system. These results provide further insight into the genetic variability of *Campylobacter* spp. The results facilitated determination of the core *C. jejuni* genome, and also provides information regarding putative virulence factors.

These experiments will hopefully serve as a basis for future investigations, which will help us better understand *Campylobacter* spp. Further studies should investigate genes that code for hypothetical proteins. Understanding *Campylobacter* spp. will potentially help develop better strategies for reducing poultry associated *Campylobacter* spp. and potentially reduce human exposure to the organism so that public health is improved.