Template for Written Asthma Action Plan

Symptoms	Medication
WHEN WELL • No asthma symptoms	Regular Controller Treatment EVERYDAY:
Before exercise	1. 2. 3. Reliever puffs ONLY when necessary
CAUTION	
 Wake at night due to asthma symptoms Have day time asthma symptoms more than 2 times Used reliever more then 2 times Have limited activity or exercise Have flu like symptoms 	STEP UP TREATMENT 1 puffs times/day for next 7-14 days. If improved go back to regular treatment. 2. Reliever puffs 4-6 hourly x 3 days If on Symbicort® 2-4 puffs at a time Do not exceed 12 puffs/day If improved go back to regular treatment.
• If NO improvement at anytime with the above treatment then add	Prednisolone 30 mg per day x 5-7 days. (for Adults) (Children should consult Dr. first)
DANGER GET HELP WHEN • Severe shortness of breath • Reliever medicine is not helping • Can only speak in short sentence • Feeling frightened	SEE YOUR DOCTOR DO NOT WAIT CALL 995 FOR AN AMBULANCE Reliever puffs at 10 minutes interval till you get to the nearest Dr. or hospital. Prednisolone 30 mg immediately.
Affix Patient Stickers	Reinforced by: Date:

Disclaimer:

All information contained herein is intended for your general information only and is not a substitute for medical advice for treatment of asthma. If you have specific questions, consult your doctor.