

Unique Plan Description: SURG Pre-Operative SCIP VC

Plan Selection Display: SURG Pre-Operative SCIP VC

PlanType: Medical

Version: 1

Begin Effective Date: 9/11/2013 15:31

End Effective Date: Current

Available at: **Ascension Health**

VC Rehab Hosp

VCH Behav Hlth

VCH HH Hospice

VCH Select

VCH St Francis

VCH St Joseph

VCH St Teresa

VCHSJ Assess

VCHSJ SBH

VCHST IP Rehab

SURG Pre-Operative SCIP VC

Patient Care

- Sequential Compression Device
T;N, Constant Order

Medications

***Dosing based upon weight for the following agents

1. Cefazolin - < 80 kg - 1 gm
80 kg - 120 kg - 2 gm
> 120 kg - 3 gm
2. Clindamycin - <= 100 kg - 600 mg
> 100 kg - 900 mg
3. Cefotetan - <= 80 kg - 1 gm
> 80 kg - 2 gm
4. Vancomycin - <= 100 kg - 1 gm
> 100 kg - 1.5 gm*** <i>(NOTE)*</i>

Give alternative antibiotic if patient has either: - Cephalosporin allergy - Severe penicillin allergy*(NOTE)**

Cardiothoracic Surgery

Drug of Choice*(NOTE)**

- ceFAZolin
*1 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe (DEF)**
Comments: Give within 60 minutes of the incision
2 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe
Comments: Give within 60 minutes of the incision
3 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe
Comments: Give within 60 minutes of the incision

Alternative Antibiotic or give vancomycin if patient is MRSA colonized*(NOTE)**

- vancomycin
*1 gm, Infuse Over: 90 minutes, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV (DEF)**
Comments: Give within 120 minutes of incision
1.5 gm, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV
Comments: Give within 120 minutes of incision

Vascular Surgery

Drug of Choice*(NOTE)**

- ceFAZolin
*1 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe (DEF)**
Comments: Give within 60 minutes of the incision
2 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe
Comments: Give within 60 minutes of the incision
3 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe
Comments: Give within 60 minutes of the incision

Alternative Antibiotic or give vancomycin if patient is MRSA colonized*(NOTE)**

- vancomycin
 - 1 gm, Infuse Over: 90 minutes, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV (DEF)**
 - Comments: Give within 120 minutes of incision*
 - 1.5 gm, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV*
 - Comments: Give within 120 minutes of incision*

Obstetric Surgery

Drug of Choice (NOTE)*

- ceFAZolin
 - 1 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe (DEF)**
 - Comments: Give within 60 minutes of the incision*
 - 2 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe*
 - Comments: Give within 60 minutes of the incision*
 - 3 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe*
 - Comments: Give within 60 minutes of the incision*

Alternative Antibiotic (NOTE)*

- Cleocin Phosphate
 - 600 mg, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, infuse over 30 minutes, Form: Soln-IV (DEF)**
 - Comments: Give within 120 minutes of incision*
 - 900 mg, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, infuse over 60 minutes, Form: Soln-IV*
 - Comments: Give within 120 minutes of incision*

Give vancomycin in addition to selected antibiotic if patient is MRSA colonized (NOTE)*

- vancomycin
 - 1 gm, Infuse Over: 90 minutes, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV (DEF)**
 - Comments: Give within 120 minutes of incision*
 - 1.5 gm, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV*
 - Comments: Give within 120 minutes of incision*

Neurosurgery

Antibiotic of Choice for a Clean Procedure (NOTE)*

- ceFAZolin
 - 1 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe (DEF)**
 - Comments: Give within 60 minutes of the incision*
 - 2 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe*
 - Comments: Give within 60 minutes of the incision*
 - 3 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe*
 - Comments: Give within 60 minutes of the incision*

Alternative Antibiotic for a Clean Procedure (NOTE)*

- vancomycin
 - 1 gm, Infuse Over: 90 minutes, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV (DEF)**
 - Comments: Give within 120 minutes of incision*
 - 1.5 gm, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV*
 - Comments: Give within 120 minutes of incision*

Antibiotic of Choice for a Clean-Contaminated Procedure (NOTE)*

- Cleocin Phosphate
 - 600 mg, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, infuse over 30 minutes, Form: Soln-IV (DEF)**
 - Comments: Give within 120 minutes of incision*
 - 900 mg, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, infuse over 60 minutes, Form: Soln-IV*
 - Comments: Give within 120 minutes of incision*

Alternative Antibiotic for a Clean-Contaminated Procedure**(NOTE)***

- Unasyn
3 gm, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Soln-IV

Comments: Give within 60 minutes of incision

Give vancomycin in addition to selected antibiotic if patient is MRSA colonized**(NOTE)***

- vancomycin
*1 gm, Infuse Over: 90 minutes, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV (DEF)**

Comments: Give within 120 minutes of incision

1.5 gm, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV

Comments: Give within 120 minutes of incision

Head and Neck Surgery

****Antibiotic of Choice*****(NOTE)*****

- ceFAZolin
*1 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe (DEF)**

Comments: Give within 60 minutes of the incision

2 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe

Comments: Give within 60 minutes of the incision

3 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe

Comments: Give within 60 minutes of the incision

Alternative Antibiotics - Order both clindamycin and gentamicin**(NOTE)***

- Cleocin Phosphate
*600 mg, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, infuse over 30 minutes, Form: Soln-IV (DEF)**

Comments: Give within 120 minutes of incision

900 mg, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, infuse over 60 minutes, Form: Soln-IV

Comments: Give within 120 minutes of incision

- gentamicin
80 mg, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Soln-IV

Comments: Give within 60 minutes of incision

Give vancomycin in addition to selected antibiotic if patient is MRSA colonized**(NOTE)***

- vancomycin
*1 gm, Infuse Over: 90 minutes, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV (DEF)**

Comments: Give within 120 minutes of incision

1.5 gm, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV

Comments: Give within 120 minutes of incision

Orthopedic Surgery

*****Drug of Choice*****(NOTE)*****

- ceFAZolin
*1 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe (DEF)**

Comments: Give within 60 minutes of the incision

2 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe

Comments: Give within 60 minutes of the incision

3 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe

Comments: Give within 60 minutes of the incision

Alternative Antibiotic**(NOTE)***

- Cleocin Phosphate
*600 mg, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, infuse over 30 minutes, Form: Soln-IV (DEF)**

Comments: Give within 120 minutes of incision

900 mg, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, infuse over 60 minutes, Form: Soln-IV

Comments: Give within 120 minutes of incision

Give vancomycin in addition to selected antibiotic if patient is MRSA colonized (NOTE)*

vancomycin

1 gm, Infuse Over: 90 minutes, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV (DEF)*

Comments: Give within 120 minutes of incision

1.5 gm, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV

Comments: Give within 120 minutes of incision

Abdominal/Colorectal Surgery

Drug of Choice (NOTE)*

cefoTEtan

1 g, IV Piggyback, Once, PRN Other (See Comment), Form: Injection (DEF)*

2 g, IV Piggyback, Once, PRN Other (See Comment), Form: Injection

Alternative Antibiotics - Give both Ciprofloxacin and Metronidazole (NOTE)*

Cipro

400 mg =, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Soln-IV

Comments: Give within 120 minutes of incision

Flagyl

500 mg, IV Piggyback, Once, PRN Other (See Comment), infuse over 1 hr, Form: Soln-IV

Comments: Give within 120 minutes of incision

Give vancomycin in addition to selected antibiotic if patient is MRSA colonized (NOTE)*

vancomycin

1 gm, Infuse Over: 90 minutes, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV (DEF)*

Comments: Give within 120 minutes of incision

1.5 gm, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV

Comments: Give within 120 minutes of incision

Gynecologic Surgery

Drug of Choice (NOTE)*

cefoTEtan

1 g, IV Piggyback, Once, PRN Other (See Comment), Form: Injection (DEF)*

2 g, IV Piggyback, Once, PRN Other (See Comment), Form: Injection

Alternative Antibiotics - Give both Ciprofloxacin and Clindamycin (NOTE)*

Cleocin Phosphate

600 mg, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, infuse over 30 minutes, Form: Soln-IV (DEF)*

Comments: Give within 120 minutes of incision

900 mg, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, infuse over 60 minutes, Form: Soln-IV

Comments: Give within 120 minutes of incision

Cipro

400 mg =, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Soln-IV

Comments: Give within 120 minutes of incision

Give vancomycin in addition to selected antibiotic if patient is MRSA colonized (NOTE)*

vancomycin

1 gm, Infuse Over: 90 minutes, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV (DEF)*

Comments: Give within 120 minutes of incision

1.5 gm, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV

Comments: Give within 120 minutes of incision

Urologic/Genitourinary Surgery

Drug of Choice (NOTE)*

- ceFAZolin
 - 1 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe (DEF)**
Comments: Give within 60 minutes of the incision
 - 2 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe*
Comments: Give within 60 minutes of the incision
 - 3 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe*
Comments: Give within 60 minutes of the incision

Alternative Antibiotic (NOTE)*

- Cipro
 - 400 mg =, IV Piggyback, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Soln-IV*
Comments: Give within 120 minutes of incision

Give vancomycin in addition to selected antibiotic if patient is MRSA colonized (NOTE)*

- vancomycin
 - 1 gm, Infuse Over: 90 minutes, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV (DEF)**
Comments: Give within 120 minutes of incision
 - 1.5 gm, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV*
Comments: Give within 120 minutes of incision

Procedure

Drug of Choice (NOTE)*

- ceFAZolin
 - 1 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe (DEF)**
Comments: Give within 60 minutes of the incision
 - 2 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe*
Comments: Give within 60 minutes of the incision
 - 3 gm, IV Push, Once, PRN Other (See Comment), Prophylaxis of infection: Surgical, Form: Syringe*
Comments: Give within 60 minutes of the incision

Alternative Antibiotic or give vancomycin if patient is MRSA colonized (NOTE)*

- vancomycin
 - 1 gm, Infuse Over: 90 minutes, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV (DEF)**
Comments: Give within 120 minutes of incision
 - 1.5 gm, IV Piggyback, Once, PRN Other (See Comment), Form: Soln-IV*
Comments: Give within 120 minutes of incision

***Report Legend:**

DEF - This order sentence is the default for the selected order

GOAL - This component is a goal

IND - This component is an indicator

INT - This component is an intervention

IVS - This component is an IV Set

NOTE - This component is a note

Rx - This component is a prescription

SUB - This component is a sub phase